Table 1: Cycling competition history according to presence of fibrosis.

	Cyclist LGE-	Cyclist LGE+	P value
N	20	21	
Active Years	$24.4 \pm 11.3$	$28.4 \pm 14.5$	0.34
Training per week (hrs)	$11.5 \pm 1.8$	$11.5 \pm 2.1$	0.91
<b>Total no. Competitions</b>	$622.5 \pm 587.7$	$702.1 \pm 567.4$	0.67
Competitions per year	$22.8 \pm 13.4$	$31.3 \pm 27.7$	0.22
<b>Total distance (miles)</b>	$22\ 853.7 \pm 33\ 546.7$	$33\ 595.0 \pm 38\ 063.1$	0.35
Average distance per race (miles)	$40.4 \pm 59.9$	$49.0 \pm 43.3$	0.61
Total time (hrs)	1194.1 ± 1456.3	$1633.2 \pm 1610.2$	0.37

Values are mean  $\pm$  standard deviation.

LGE- = Fibrosis absent; LGE+ = Fibrosis present.

Table 2: Triathlon competition history according to presence of fibrosis.

	Triathlete LGE-	Triathlete	P value	
	LGE+			
N	6	3		
<b>Active Years</b>	$25.5 \pm 9.4$	$23.0 \pm 13.0$	0.75	
Training per week (hrs)	$11.3 \pm 1.7$	$10.7 \pm 1.2$	0.61	
<b>Total no. Competitions</b>	$215.3 \pm 195.7$	$81.7 \pm 89.8$	0.31	
Competitions per year	$7.5 \pm 5.3$	$3.0 \pm 2.6$	0.22	
<b>Total swimming distance</b>	$247.4 \pm 175.6$	$102.3 \pm 99.6$	0.23	
(km)				
Total cycling distance (km)	$7343.3 \pm 4596.2$	$3073.3 \pm 2731.0$	0.19	
<b>Total running distance (km)</b>	$1807.3 \pm 1149.6$	$759.3 \pm 672.0$	0.20	
Total distance (km)	$9398.0 \pm 5918.1$	$3935.0 \pm 3500.6$	0.19	
Total time (hrs)	$465.7 \pm 270.6$	$185.7 \pm 153.7$	0.15	
No of sprint distances	$103.0 \pm 134.5$	$37.3 \pm 46.5$	0.45	
No of Olympic distances	$316.0 \pm 588.9$	$35.7 \pm 47.5$	0.45	
No of middle distances	$11.0 \pm 8.7$	$2.7 \pm 3.1$	0.16	
No of iron man distances	$4.0 \pm 7.5$	$3.3 \pm 4.9$	0.90	

Values are mean  $\pm$  standard deviation.

LGE- = Fibrosis absent; LGE+ = Fibrosis present.