



# Weekly Influenza and COVID-19 Surveillance graphs

UKHSA publishes a weekly national influenza and COVID-19 surveillance report which summarises the information from the surveillance systems which are used to monitor influenza, COVID-19 and other seasonal respiratory viruses in England.

Additional figures based on these surveillance systems are included in this slide set.

The figures presented in this slide set are based on data from week 15 (between 11 April and 17 April 2022).



# Contents

- 1) [COVID-19 Pandemic Overview](#)
- 2) [Confirmed COVID-19 episodes in England](#)
- 3) [Respiratory Datamart system \(England\)](#)
- 4) [Second generation surveillance system \(SGSS\)](#)
- 5) [Community surveillance](#)
- 6) [Surveillance in 'educational-age' cohorts](#)
- 7) [Secondary Care surveillance](#)
- 8) [Mortality surveillance](#)
- 9) [Possible reinfections in England](#)
- 10) [Co/secondary infections with COVID-19](#)




UK Health  
Security  
Agency

# COVID-19 Pandemic Overview




# Confirmed COVID-19 episodes tested under Pillar 1, by sample week, since week 5 2020






# Confirmed COVID-19 episodes tested under Pillar 2, by sample week, since week 5 2020






# Weekly overall hospital and ICU/HDU admission rates per 100,000 of new COVID-19 cases reported through SARI Watch, England since week 12 2020





# Number of deaths since week 10 2020 by week of death and time since laboratory confirmation of COVID-19, England



Since 1 April 2022, free universal symptomatic and asymptomatic testing for the general public in England is no longer available, as outlined in the plan for [living with COVID-19](#). Data should be interpreted in the context of this change to testing.



UK Health  
Security  
Agency

# Confirmed COVID-19 episodes in England






## Data Information

- From the week 32 report onwards, case rates have been updated to use the latest ONS population estimates for mid-2020. Previously case rates were calculated using the mid-2019 population estimates
- From 11 January 2022 the requirement for [confirmatory PCR testing in individuals who test positive using a lateral flow device was temporarily removed](#).
- Rates by ethnicity and IMD quantile will continue to be presented using the mid-2019 estimates, until the mid-2020 estimates become available.
- From 31 January 2022, UKHSA moved all COVID-19 case reporting in England to use a new episode-based definition which includes possible reinfections. Each infection episode is counted separately if there are at least 91 days between positive test results (PCR or LFD). Each infection episode begins with the earliest positive specimen date. Further information can be found on the [UK COVID-19 dashboard](#).
- Since 1 April 2022, free universal symptomatic and asymptomatic testing for the general public in England is no longer available, as outlined in the plan for [living with COVID-19](#). As such, there will be a reduction in the reporting of data obtained through Pillar 2 from April 2022 onwards. Data in this report should be interpreted in the context of this change to testing. [Public health guidance](#) remains in place for cases and their close contacts




# Confirmed COVID-19 episodes tested under Pillar 2, based on sample week with overall weekly PCR positivity for Pillar 2 (%)






# Weekly confirmed COVID-19 case rates per 100,000, by episode, tested under Pillar 2, by sex






# Weekly confirmed COVID-19 case rates per 100,000, by episode, tested under Pillar 2, by age group






# Weekly PCR positivity (%) of confirmed COVID-19 cases tested overall and by sex under Pillar 2






# Weekly PCR positivity (%) of confirmed COVID-19 cases tested under Pillar 2, by male and age group






# Weekly PCR positivity (%) of confirmed COVID-19 cases tested under Pillar 2, by female and age group






# Weekly confirmed COVID-19 case rates by episode, per 100,000 population (Pillar 2), by UKHSA Centres and sample week








# Weekly PCR positivity of confirmed COVID-19 cases tested under Pillar 2 (%) by UKHSA Centres and sample week






# Weekly incidence per 100,000 population by ethnicity (Pillar 2), England






# Weekly COVID-19 episodes tested under Pillar 1, per 100,000 population by age group and region, weeks 6 to 15






# Weekly COVID-19 episodes tested under Pillar 2, per 100,000 population by age group and region, weeks 6 to 15





# Weekly COVID-19 episodes tested under Pillar 1, per 100,000 population by ethnicity and region, weeks 6 to 15




- Black/African/Caribbean/Black British
- Indian (Asian or British)
- Mixed/Multiple Ethnic Groups
- Other Asian/Asian British
- Other ethnic group
- Pakistani (Asian or British)
- White

\*these incidence rates have been calculated using the mid-2019 ONS population estimates




# Weekly COVID-19 episodes tested under Pillar 2 per 100,000 population by ethnicity and region, weeks 6 to 15






# Weekly COVID-19 rate tested under Pillar 1, per 100,000 population by IMD quintile (1 being the most deprived and 5 being the least deprived)



\*these incidence rates have been calculated using the mid-2019 ONS population estimates




# Weekly COVID-19 rate tested under Pillar 2, per 100,000 population by IMD quintile (1 being the most deprived and 5 being the least deprived)








# Weekly rate of COVID-19 episodes per 100,000 population (Pillar 2), by upper-tier local authority, England (box shows enlarged map of London area)






# Cumulative rate of COVID-19 episodes per 100,000 population tested under Pillar 1, by upper-tier local authority, England (box shows enlarged map of London area)





# Cumulative rate of COVID-19 episodes per 100,000 population tested under Pillar 2, by upper-tier local authority, England (box shows enlarged map of London area)









UK Health  
Security  
Agency

# Respiratory Datamart system (England)




# Respiratory DataMart – Influenza subtypes






# Respiratory DataMart – Respiratory syncytial virus (RSV)









# Respiratory DataMart – Respiratory syncytial virus (RSV) weekly positivity by UKHSA region







# Respiratory DataMart – other respiratory viruses








UK Health  
Security  
Agency

# Second generation surveillance system (SGSS)


# SGSS reported Influenza A cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.

Previously, this data was presented by report date however is now presented by specimen date.

## SGSS reported Influenza B cases by region (all ages)




The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.

Previously, this data was presented by report date however is now presented by specimen date.




# SGSS reported RSV cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution. Previously, this data was presented by report date however is now presented by specimen date.




# SGSS reported Adenovirus cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.




# SGSS reported Parainfluenza cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.




# SGSS reported Rhinovirus cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.



# SGSS reported hMPV cases by region (all ages)



The presented figures are based on laboratory reports through SGSS. Testing and reporting procedures vary by virus, UKHSA Centre and over time, including short-term trends in testing. Therefore comparisons should be done with caution.





UK Health  
Security  
Agency

# Community surveillance



# COVID-19 clusters or outbreaks in educational settings

UK Health  
Security  
Agency

## Data Information

- we report on new acute respiratory infection (ARI) incidents reported to Health Protection Teams (HPTs) and entered on HPZone in the previous reporting week in educational settings by locality
- individual case notes are reviewed by an epidemiologist and an assessment made about whether the criteria for a confirmed COVID-19 cluster or outbreak are met. See definitions below
- the incidents captured on HPZone represent a subset of all ongoing clusters and outbreaks in England. A variety of arrangements are in place with local authorities and other stakeholders supporting HPTs, however data may not routinely be documented on HPZone. As a result, the number of outbreaks reported for some of the regions are underestimates
- For the 2021-2022 academic year the thresholds for reporting an outbreak in an educational setting to HPTs and HPZone have been revised, therefore comparisons with the 2020 to 2021 season should be interpreted with caution. Please see the next slide for the updated thresholds.

## Caveats

- National Schools and Universities helplines remain in place to support educational settings to manage cases and outbreaks that may not require HPT input
- From Monday 19 July 2021, schools, colleges and nurseries no longer carry out routine contact tracing. Close contacts are now identified and contacted by NHS Test and Trace.



## Thresholds for reporting

For the 2021-2022 academic year the thresholds for reporting an outbreak in an educational setting to HPZone have been revised, therefore when comparing with the 2020-2021 season, please interpret with caution.

Clusters and outbreaks are now reported to HPZone if either of the two following criteria are met:

- 5 cases or 10% (whichever is reached first) test-confirmed cases of COVID-19 (either PCR testing or LFD Ag testing with follow-up PCR) within 10 days, among students or staff clustered in a consistent group or cohort. Dates should be calculated based on illness onset, or test date if asymptomatic

Or

- Evidence of severe illness e.g. students or staff members admitted to hospital or a death as a result of a COVID-19 infection (PCR or LFD Ag with follow up PCR) as the setting may require advice on risk assessment and communication.

## Definitions


**Cluster:** two or more test-confirmed cases of COVID-19 among individuals associated with a specific non-residential setting with illness onset dates within a 14-day period (in the absence of detailed information about the type of contact between the cases).

**Outbreak:** two or more test-confirmed cases of COVID-19 among individuals associated with a specific non-residential setting with illness onset dates within 14 days, and one of:

- identified direct exposure between at least 2 of the test-confirmed cases in that setting (for example under one metre face to face, or spending more than 15 minutes within 2 metres) during the infectious period of one of the cases
- When there is no sustained local community transmission - absence of an alternative source of infection outside the setting for the initially identified cases



# Number of COVID-19 confirmed clusters or outbreaks by type of educational setting, England





# Number of COVID-19 confirmed clusters or outbreaks by type of educational setting, England

End of academic year total

Week 36 2020- 34 2021

Centres	Cumulative number of confirmed COVID-19 clusters or outbreaks by type of educational setting for the 2020/21 academic year Week 36 2020- 34 2021						
	Nursery	Primary School	Secondary School	Combined	Special Educational Needs (SEN) schools	College University	Total
<b>Total</b>	<b>846</b>	<b>2125</b>	<b>2122</b>	<b>40</b>	<b>666</b>	<b>268</b>	<b>6067</b>

Week 15 2022

Main table

PHE Centres	Cumulative number of confirmed COVID-19 clusters or outbreaks by type of educational setting for the 2021/22 academic year from Week 35 2021						
	Nursery	Primary School	Secondary School	Combined	Special Educational Needs (SEN)	College University	Total
East Midlands Centre	72 (0)	57 (0)	30 (0)	14 (0)	163 (0)	6 (1)	<b>342 (1)</b>
East of England Centre	0 (0)	12 (0)	8 (0)	3 (0)	11 (0)	2 (0)	<b>36 (0)</b>
London Centre	363 (0)	1084 (0)	257 (0)	59 (0)	208 (0)	29 (0)	<b>2000 (0)</b>
North East Centre	0 (0)	2 (0)	0 (0)	0 (0)	3 (1)	0 (0)	<b>5 (1)</b>
North West Center	10 (0)	31 (0)	13 (0)	4 (0)	120 (0)	7 (0)	<b>185 (0)</b>
South East Centre	42 (0)	388 (0)	127 (0)	34 (0)	283 (1)	7 (0)	<b>881 (1)</b>
South West Centre	4 (0)	64 (0)	79 (0)	35 (1)	246 (1)	1 (0)	<b>429 (2)</b>
West Midlands Centre	17 (0)	73 (0)	52 (0)	7 (0)	139 (0)	6 (0)	<b>294 (0)</b>
Yorkshire & the Humber	17 (0)	36 (0)	27 (0)	5 (0)	81 (0)	0 (0)	<b>166 (0)</b>
<b>Total</b>	<b>525 (0)</b>	<b>1747 (0)</b>	<b>593 (0)</b>	<b>161 (1)</b>	<b>1254 (3)</b>	<b>58 (1)</b>	<b>4338 (5)</b>



\* Number of clusters or outbreaks for the most recent week in brackets



UK Health  
Security  
Agency

# Secondary Care surveillance

# Weekly admission rates for hospital and ICU/HDU laboratory confirmed COVID-19 cases reported through SARI Watch, week 15




Source: PHE SARI-Watch (Severe Acute Respiratory Infection-Watch, formerly CHES).

\*Only NHS Acute trusts that have reported  $\geq 1$  day in the past week; excludes Specialist trusts. Acute NHS trusts (including Specialist trusts) reporting into SARI-Watch COVID-19 hospitalisation surveillance are typically around 100 per week. This was 105 for the hospitalisation (all levels of care) indicator in week 11 to 17 April 2022 inclusive and 96 trusts for the ICU/HDU indicator. For the maps, as Specialist trusts are excluded, the number of trusts providing data on COVID-19 hospitalisations in week ending 17 April 2022 was 95 and 86 for ICU/HDU admissions for COVID-19.




# Age/sex pyramid of hospitalisations (all levels of care) for COVID-19, data from sentinel acute NHS trusts, England

(a) Peak of 2<sup>nd</sup> wave (week 53 2020 to week 3 2021) n= 6,359



Reporting trusts=22

(b) Most recent 4 weeks (week 12 2022 to 15 2022) n= 2,991



Reporting trusts=14

This figure is based on individual patient level data which are provided to SARI Watch from a subset of NHS Acute Trusts, therefore the data should be interpreted with caution as the distribution of age, sex and ethnic group may not be representative of all hospitalised patients.






# Age/sex pyramid for admissions to ICU/HDU for COVID-19, mandatory case level data, acute NHS trusts, England


(a) Peak of 2<sup>nd</sup> wave (week 53 2020 to week 3 2021) n= 3,349

(b) Most recent 4 weeks (week 12 2022 to 15 2022) n=266

■ Male ■ Female



Reporting trusts=70



Reporting trusts=39


This figure is based on individual patient level data which are provided to SARI Watch from a subset of NHS Acute Trusts, therefore the data should be interpreted with caution as the distribution of age, sex and ethnic group may not be representative of all hospitalised patients.



# Laboratory confirmed admissions for COVID-19, to acute NHS trusts, by level


## UK Health Security Agency of care and ethnicity

### (a) Peak of 2<sup>nd</sup> wave (week 53 2020 to week 3 2021)



Reporting trusts  
 Lower level of care=5932  
 ICU/HDU=3060

### (b) Most recent 4 weeks (week 12 2022 to 15 2022)




Reporting trusts  
 Lower level of care=2731  
 ICU/HDU=231

This figure is based on individual patient level data which are provided to SARI Watch from a subset of NHS Acute Trusts, therefore the data should be interpreted with caution as the distribution of age, sex and ethnicity may not be representative of all hospitalised patients.


**Caveat:** From week 24 the ethnicity analysis is based on a new method for assigning ethnicity, developed by UKHSA. The previous method used the most recent ethnicity recorded through linkage to Hospital Episode Statistics. However, this method led to unfeasibly high rates in the 'Other' ethnic group when applied to COVID-19 cases, hospitalisation or mortality. The new method uses the most frequent ethnicity recorded through linkage to Hospital Episode Statistics, unless the most frequent was 'Other' when the second most frequent was chosen.



# Weekly COVID-19 hospitalisation rate per 100,000 trust catchment population by age group and region, weeks 6 to 15




# Hospital admission rate (excluding ICU/HDU) by ethnicity per 100,000 trust catchment population



Caveat: From week 24 the ethnicity analysis is based on a new method for assigning ethnicity, developed by UKHSA. The previous method used the most recent ethnicity recorded through linkage to Hospital Episode Statistics. However, this method led to unfeasibly high rates in the 'Other' ethnic group when applied to COVID-19 cases, hospitalisation or mortality. The new method uses the most frequent ethnicity recorded through linkage to Hospital Episode Statistics, unless the most frequent was 'Other' when the second most frequent was chosen.




# Rate of admission to ICU/HDU by ethnicity, per 100,000 trust catchment population



Caveat: From week 24 the ethnicity analysis is based on a new method for assigning ethnicity, developed by UKHSA. The previous method used the most recent ethnicity recorded through linkage to Hospital Episode Statistics. However, this method led to unfeasibly high rates in the 'Other' ethnic group when applied to COVID-19 cases, hospitalisation or mortality. The new method uses the most frequent ethnicity recorded through linkage to Hospital Episode Statistics, unless the most frequent was 'Other' when the second most frequent was chosen.



# COVID-19 as primary reason for admission among SARS-CoV-2 positive patient by month of admission



### Notes

- 1) Case-level sentinel data from week 35 2021 (commencing 30 August 2021) to week 15 2022 (ending 17 April 2022) inclusive
- 2) 21% (4224/19906) of total records in this period have missing data on the 'Admission due to COVID-19' indicator – these are excluded from analysis
- 3) London trusts under-represented




UK Health  
Security  
Agency

# Mortality surveillance




# Cumulative mortality rate of COVID-19 cases per 100,000 population tested under Pillar 1 and 2 since the beginning of the pandemic by (a) 28 day definition and (b) 60 day definition

(a)




(b)







# Age-adjusted mortality rate\*\* (per 100,000 population) in confirmed cases of COVID-19 by IMD quintile, by week using the 60 day definition



\*\*Rates are time-adjusted: a weekly population denominator has been used to calculate the mortality rate



UK Health  
Security  
Agency

# Co/secondary infections with COVID-19



## Co/secondary infections with COVID-19

- Caveat - undertesting for other pathogens may result in an underestimate of co/secondary infection cases.
- Preceding/co-/secondary infections refers to when a patient has a COVID-19 infection with one or more other pathogen (Please see Appendix 1 – Pre-/co-/secondary infection with COVID-19 definitions.)
  - Preceding infection: COVID-19 acquired after another pathogen
  - Co-infection: COVID-19 and other pathogen acquired at the same time
  - Secondary infection: COVID-19 acquired before another pathogen
- Numbers of pre-/co-/secondary infection remain low across UKHSA surveillance systems.
- For patients with severe respiratory failure requiring Extra Corporeal Membrane Oxygenation (ECMO), analysis of data from five adult ECMO centres in England indicates that among patients with severe respiratory failure due to COVID-19, almost a third of these have co/secondary infections.
- Published data analysis from pandemic wave 1 (W-1) indicates increased mortality associated with COVID-19 and [influenza](#), [key bacterial and fungal infections](#) and [invasive pneumococcal disease \(IPD\)](#) in comparison to patients without co/secondary infection.
- [Data analysis](#) from W-1 indicates that *Aspergillus* and *candidemia* cases have increased risk of mortality in comparison to patients without co/secondary infection.



## UK Health Security Agency Co/secondary infections among patients with severe respiratory failure requiring Extra Corporeal Membrane Oxygenation (ECMO)

Analysis is based on cumulative data from five adult ECMO centres in England. Data for the current and previous seasons are presented. Each season commences around October (ISO week 40) ending in September in the following year (ISO week 39).

Data for the current season (2021-22) is from 4 October 2021 to 6 March 2022 inclusive (week 40 2021 to week 09 2022). This period includes effects from the Delta and Omicron waves of the pandemic. The 2020-21 season is from 28 September 2020 to 3 October 2021 inclusive and includes effects from the Alpha and Delta waves. The 2019-20 season is from 30 September 2019 to 27 September 2020 inclusive and includes effects from the original Wuhan strain.

- In the 2021-22 season, 32% (18/56) of ECMO patients admitted for severe respiratory failure due to laboratory confirmed COVID-19 had clinically significant co/secondary infections. In the previous season (2020-21) this proportion was 30% (116/381). In the 2019-20 season this proportion was 33% (79/236).
- In all three seasons the majority of clinically significant co/secondary infections among respiratory failure COVID-19 cases comprised Gram-negative bacilli from the order Enterobacterales:
  - 39% (7/18) in the current season 2021-22
  - 29% (34/116) in 2020-21 and 46% (36/79) in 2019-20. The proportion decreased significantly in 2020-21 compared to 2019-20 and no change was detected in other key pathogens between these two periods.



UK Health  
Security  
Agency

# Surveillance of bacterial, fungal and viral infections, in COVID-19 patients in England, Jan 2020 – Apr 2022

HCAI, Fungal, AMR, AMU & Sepsis Division



# Updates

- From 31 January 2022, UKHSA has changed the COVID-19 case definition to include multiple infection episodes. Reported co-/secondary/preceding infections in England now use the new definition, revising all cases back to the beginning of the pandemic.
- The Unified Infection Dataset (UID) project has been extended to incorporate the Co- and Secondary infections with COVID-19 datasets
- The following outputs included in this section have been produced via the UID, combining previously separate data pipelines
  - Key HCAI bacterial and fungal specimens reported to SGSS and HCAI data capture system
  - Respiratory viral specimens reported to SGSS and Respiratory Datamart
  - Fungal specimens reported to mycology reference lab (MRL)
- The Co- and secondary infections team have undertaken an extensive data validation exercise which has identified additional respiratory viral specimens from Respiratory Datamart and allowed us to make improvements to the methodologies. Preceding infections for all pipelines (other pathogen infections occurring before COVID-19 specimen) are now included. Please note, all cases since January 2020 have been revised in line with this validation.
- Data are provisional and subject to change due to possible delayed reporting of microbiological samples



# Number of COVID-19 patient-episodes with bacterial, fungal or viral infections in COVID-19 patients diagnosed in England during wave 1\*, by infection type and timing of diagnosis

Bacterial/ fungal/ viral infection by specimen type	COVID-19 patient-episodes with bacterial/ fungal/ viral infection		Timing of bacterial/fungal/viral diagnosis in relation to COVID-19 diagnosis								
			Preceding infection			Coinfection			Secondary infection		
	n	% of COVID cases	n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases
Bacterial/fungal bloodstream & lower respiratory infection	129	0.05	9	6.98	<0.01	2	1.55	<0.01	118	91.47	0.05
Bacterial/fungal bloodstream infection	3,077	1.26	1,282	41.66	0.53	801	26.03	0.33	994	32.30	0.41
Bacterial/fungal lower respiratory infection	647	0.27	109	16.85	0.04	94	14.53	0.04	444	68.62	0.18
<i>Clostridioides difficile</i> infection	308	0.13	100	32.47	0.04	43	13.96	0.02	165	53.57	0.07
Fungal respiratory/bloodstream infection (MRL) ‡	93	0.04	10	10.75	<0.01	2	1.83	<0.01	81	74.31	0.03
Other respiratory virus infection	373	0.15	136	36.46	0.06	193	51.74	0.08	44	11.80	0.02
Any site†	<b>4,636</b>	<b>1.90</b>	<b>1,647</b>	<b>35.53</b>	<b>0.68</b>	<b>1,136</b>	<b>24.42</b>	<b>0.47</b>	<b>1,853</b>	<b>39.83</b>	<b>0.76</b>

## Key findings:

- 1.90% of COVID-19 patient-episodes had a bacterial, fungal or other respiratory viral infection detected in either the 28 days prior or 28 days following (60 days for MRL) their COVID-19 diagnosis
  - 1.26% had a key bacterial/fungal bloodstream infection
- The majority of infections with key organisms were categorised as secondary infections (39.83%).

Data correction as of 19 Apr 2022

Please see appendix 1 for Co- and secondary infection definitions with COVID-19  
Please note patients can have multiple COVID-19 infection-episodes, numbers here do not reflect the number of patients. Testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.

\*SARS-CoV2 specimen dates from 30 Jan 2020 to 28 Jun 2020 (N=243,902). Data for wave 1 remains static (last updated 11 Apr 2022).

‡ Definition for preceding infection differs for MRL specimens - detection within 60 days

† includes the combination Bacterial/fungal bloodstream & *Clostridioides difficile* infection (1 coinfection, 5 secondary & 1 preceding), Bacterial/fungal bloodstream, lower respiratory & *Clostridioides difficile* infection (1 secondary), & Bacterial/fungal lower respiratory & *Clostridioides difficile* infection (1 secondary) .



# Number of COVID-19 patient-episodes with bacterial, fungal or viral infections in COVID-19 patients diagnosed in England during wave 2\*, by infection type and timing of diagnosis

Bacterial/ fungal/ viral infection by specimen type	COVID-19 patient-episodes with bacterial/ fungal/ viral infection		Timing of bacterial/fungal/viral diagnosis in relation to COVID-19 diagnosis								
	n	% of COVID cases	Preceding infection			Coinfection			Secondary infection		
			n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases
Bacterial/fungal bloodstream & lower respiratory infection	438	0.01	10	2.28	<0.01	5	1.14	<0.01	423	96.58	0.01
Bacterial/fungal bloodstream infection	7,829	0.21	3,515	44.90	0.10	1,032	13.18	0.03	3,282	41.92	0.09
Bacterial/fungal lower respiratory infection	2,166	0.06	335	15.47	0.01	145	6.69	<0.01	1,686	77.84	0.05
<i>Clostridioides difficile</i> infection	789	0.02	311	39.42	0.01	77	9.76	<0.01	401	50.82	0.01
Fungal respiratory/bloodstream infection (MRL) ‡	215	0.01	26	12.09	<0.01	3	1.32	<0.01	186	81.94	0.01
Other respiratory virus infection	361	0.01	84	23.27	<0.01	140	38.78	<0.01	137	37.95	<0.01
Any site†	11,830	0.32	4,293	36.29	0.12	1,404	11.86	0.04	6,133	51.79	0.17

## Key findings:

- 0.32% of COVID-19 patient-episodes had a bacterial, fungal or other respiratory viral infection detected in either the 28 days prior or 28 days following (60 days for MRL) their COVID-19 diagnosis
- While the prevalence is lower in W2 than W1 (1.91%), the number of patient-episodes with a pre-/co-/secondary infection was much greater in W2 than W1 (11,830 vs. 4,636, respectively)
- The majority of infections with key organisms were secondary infections (51.79%).

Please see appendix 1 for Co- and secondary infection definitions with COVID-19

Please note patients can have multiple COVID-19 infection-episodes, numbers here do not reflect the number of patients.

\*SARS-CoV2 specimen dates from 29 Jun 2020 to 26 Apr 2021 (N=3,697,677). Data for wave 2 remains static (last updated 11 Apr 2022).

‡ Definition for preceding infection differs for MRL specimens - detection within 60 days

† includes the combination Bacterial/fungal bloodstream & *Clostridioides difficile* infection (2 coinfection, 12 secondary & 10 preceding), Bacterial/fungal bloodstream, lower respiratory & *Clostridioides difficile* infection (2 secondary), & Bacterial/fungal lower respiratory & *Clostridioides difficile* infection (4 secondary & 2 preceding)

Data correction as of 19 Apr 2022





UK Health  
Security  
Agency

Number of COVID-19 patient-episodes with bacterial, fungal or viral infections in COVID-19 patients diagnosed in England during wave 3\*, by infection type and timing of diagnosis

Bacterial/ fungal/ viral infection by specimen type	COVID-19 patient-episodes with bacterial/ fungal/ viral infection		Timing of bacterial/fungal/viral diagnosis in relation to COVID-19 diagnosis								
	n	% of COVID cases	Preceding infection			Coinfection			Secondary infection		
			n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases	n	% infections by site	% of COVID cases
Bacterial/fungal bloodstream & lower respiratory infection	217	<0.01	13	5.99	<0.01	6	2.76	<0.01	198	91.24	<0.01
Bacterial/fungal bloodstream infection	4,753	0.04	1,799	37.85	0.02	877	18.45	0.01	2,077	43.70	0.02
Bacterial/fungal lower respiratory infection	1,940	0.02	362	18.66	<0.01	166	8.56	<0.01	1,412	72.78	0.01
<i>Clostridioides difficile</i> infection	609	0.01	225	36.95	<0.01	62	10.18	<0.01	322	52.87	<0.01
Fungal respiratory/bloodstream infection (MRL) ‡	195	<0.01	12	6.15	<0.01	3	1.54	<0.01	180	92.31	<0.01
Other respiratory virus infection	2,046	0.02	604	29.52	0.01	912	44.57	0.01	530	25.90	<0.01
Any site†	9,788	0.08	3,022	30.87	0.03	2,029	20.73	0.02	4,737	48.40	0.04

Key findings:

- 0.08% of COVID-19 patient-episodes had a bacterial, fungal or other respiratory viral infection detected in either the 28 days prior or 28 days following (60 days for MRL) their COVID-19 diagnosis
- Prevalence in W3 lower than W2 and W1; however, patient-episodes of COVID-19 and another key infection are still higher in W3 than W1 (9,788 vs 4,636, respectively)
- The majority of infections with key organisms were categorised as secondary infections (48.40%).

Data correction as of 19 Apr 2022

Please see appendix 1 for Co- and secondary infection definitions with COVID-19

Please note patients can have multiple COVID-19 infection-episodes, numbers here do not reflect the number of patients.


\*SARS-CoV2 specimen dates from 27 Apr 2021 to 13 Feb 2022 (N=11,733,771). Last updated 18 Apr 2022.

‡ Definition for preceding infection differs for MRL specimens - detection within 60 days

† includes the combination Bacterial/fungal bloodstream & *Clostridioides difficile* infection (15 secondary & 7 preceding), Bacterial/fungal bloodstream, lower respiratory & *Clostridioides difficile* infection (1 coinfection & 1 secondary), & Bacterial/fungal lower respiratory & *Clostridioides difficile* infection (2 coinfection & 2 secondary)



# Most frequent bacterial/fungal species in blood specimens, by timing of diagnosis, in COVID-19 patients diagnosed in England




## Key findings:

In all three waves, the most frequent bacterial/fungal organisms identified from blood specimens were *Escherichia coli*, *Staphylococcus aureus* and *Klebsiella pneumoniae*.

Please note, testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.



# Most frequent bacterial/fungal species in lower respiratory tract specimens, by timing of diagnosis, in COVID-19 patients diagnosed in England



Please note different y-axes


### Key findings:

In all three waves, the most frequent bacterial/fungal organisms identified from respiratory specimens were *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*.

Please note, testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.



# Most frequent viral specimens, by timing of diagnosis, in COVID-19 patients diagnosed in England



Please note different y-axes


## Key findings:

The most frequent viral organisms identified from respiratory specimens changed from wave to wave, rhinovirus was identified within the top three pre-/co-/secondary infection causative organisms for all waves and most prevalent organism for W1 and W2, with RSV accounting for the most pre-/co-/secondary infections in W3

Please note, testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.



## Most frequent fungal species (MRL), by timing of diagnosis, in COVID-19 patients diagnosed in England



Please note different y-axes

### Key findings:

In all three waves, the most frequent fungal organisms identified were *Aspergillus fumigatus complex* and *Candida albicans*.

Please note, testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.



## COVID-19 Co/secondary infection with fungi and vaccine preventable bacteria

Bacteria/Fungi	First Wave	Second Wave	Third Wave	Total Cases
	(30 Jan 2020 - 28 June 2020)	(29 June 2020 – 30 April 2021)	(1 May 2021 – 24 February 2021)	
<i>Aspergillus fumigatus</i> isolates (azole resistant)	46 (4)	120 (2)	137(12)	<b>303(18)</b>
Probable/Proven cases of CAPA*	15	38	44	<b>97</b>
<i>Candida</i> spp.: Candidemia	63	133	17	<b>213</b>
<i>Bordetella pertussis</i>	0	0	0	<b>0</b>
<i>Haemophilus influenzae</i>	3	2	0	<b>5</b>
<i>Neisseria meningitidis</i>	2	0	0	<b>2</b>
<i>Streptococcus pneumoniae</i>	40	45	14	<b>99</b>

x

\*\*

## \*COVID-19-associated pulmonary aspergillosis

Please note fungal data refers to secondary infections only. Mycology data contains results from Mycology reference laboratory data, Candidaemia is representative of deep infection. One case of osteomyelitis, one case of ventriculitis and one case of endocarditis was documented in wave two. Fungal data are also included in the overall numbers in slides 6-8 but have been stratified here with additional details. *Bordetella pertussis* co-infection is defined as +/- 28 d Culture/PCR (based on pertussis sample date), +/- 28 Serology/Oral fluid (anti-pertussis toxin Ig) (based on pertussis symptom onset date, excluding cases without onset date). *Haemophilus influenzae*, *Neisseria meningitidis* and *Streptococcus pneumoniae* co-infection is defined as +/- 2d. *Legionella*, *Mycoplasma* and gastrointestinal infection data not included. Please note, testing in W1 was not open to the community and therefore W1 cases are predominantly hospitalised patients vs. W2 and W3.

# Appendix 1: Pre-/co-/secondary infection definitions with COVID-19

The day pertains to the date of the sample collection that yielded a positive result. These definitions do not apply to persistent COVID-19 patients. Patients with persistent COVID-19 require independent clinical assessment.

Organism	Definition co-infection with SARS-CoV-2 †	Definition of infection pre-SARS-CoV-2 infection (other pathogen is primary infection) or Definition of post SARS-CoV-2 secondary infection (SARS-CoV-2 is primary infection)
Influenza A	+/- 1d	2-28d <sup>^</sup>
Influenza B	+/- 1d	2-28d <sup>^</sup>
RSV	+/- 1d	2-28d
Adenovirus	+/- 1d	2-28d
Enterovirus	+/- 1d	2-28d
Human metapneumovirus	+/- 1d	2-28d
Parainfluenza (any subtype)	+/- 1d	2-28d
Seasonal coronavirus	+/- 1d *	2-28d
Rhinovirus	+/- 1d	2-28d
Co-infections in ECMO patient (patients with most severe clinical respiratory signs)		
ECMO patients	Individual case review	Individual case review
Blood stream and respiratory infections (bacterial and fungal)		
<i>Achromobacter xylosoxidans</i>	+/- 1d	2-28d
<i>Acinetobacter</i> spp.,	+/- 1d	2-28d
<i>Aspergillus</i>	+/- 1d	2-28d (pre) 2-60d (post, continually hospitalised patients only)
<i>Bordetella pertussis</i>	+/- 28 d Culture/PCR (based on pertussis sample date) +/- 28 Serology/Oral fluid (anti-pertussis toxin Ig) (based on pertussis symptom onset date, excluding cases without onset date)	N/A (Pertussis presentation is often delayed)
<i>Burkholderia cepacia</i>	+/- 1d	2-28d
<i>Candida</i> spp	+/- 1d	2-28d (pre) 2-60d (post, continually hospitalised patients only)
<i>Chlamydia pneumoniae</i>	0-7d PCR	PCR within 14-28 d (8-13d PCR*)
<i>Enterobacter</i> spp.,	+/- 1d	2-28d
<i>Enterococcus</i> spp.	+/- 1d	2-28d
<i>E. coli</i>	+/- 1d	2-28d
<i>Haemophilus influenzae</i>	+/- 2d	3-28d

Continued overleaf

# Appendix 1 continued: Pre-/co-/secondary infection definitions with COVID-19

The day pertains to the date of the sample collection that yielded a positive result. These definitions do not apply to persistent COVID-19 patients. Patients with persistent COVID-19 require independent clinical assessment.

Organism	Definition co-infection with SARS-CoV-2 †	Definition of infection pre-SARS-CoV-2 infection (other pathogen is primary infection) or Definition of post SARS-CoV-2 secondary infection (SARS-CoV-2 is primary infection)
Blood stream and respiratory infections (bacterial and fungal)		
<i>Klebsiella</i> spp.	+/- 1d	2-28d
<i>Legionella pneumophila/species</i>	Individual case review	Individual case review
<i>Mycoplasma pneumoniae</i>	0-7d PCR, IgM serology 0-21d <16y	PCR within 14-28 d (8-13d PCR*)
<i>Neisseria meningitidis</i>	+/- 2d	3-28d
<i>Pseudomonas</i> spp.,	+/- 1d	2-28d
<i>Serratia</i> spp.,	+/- 1d	2-28d
<i>Staphylococcus aureus</i>	+/- 1d	2-28d
Coag-neg <i>Staphylococcus</i> ( <i>S. haemolyticus</i> )	+/- 1d	2-28d
<i>Stenotrophomonas</i> spp., ( <i>S. maltophilia</i> )	+/- 1d	2-28d
<i>Streptococcus</i> spp. ‡	+/- 1d	2-28d
<i>Streptococcus pneumoniae</i>	+/- 2d	3-28d
Tuberculosis		
<i>Mycobacterium tuberculosis</i>	Individual case review	Individual case review
Pathogens of the immunocompromised (eg HIV)		
HIV	Individual case review	Individual case review
Gastrointestinal infections		
Listeria	0-5d *	Individual case review
Campylobacter	0-5d *	Individual case review
Shiga toxin-producing <i>E. coli</i> (STEC)	0-5d *	Individual case review
Norovirus	0-5d *	Individual case review
Salmonella	0-5d *	Individual case review
Shigella	0-5d *	Individual case review
Anaerobes		
<i>C. difficile</i>	+/- 1d	2-28d
<i>Bacteroides</i> sp. ( <i>B. fragilis</i> and non- <i>fragilis</i> <i>Bacteroides</i> )	+/- 1d	2-28d

See next slides for notes



# Appendix 1 continued: Pre-/co-/secondary infection definitions with COVID-19

## Notes

† From SARS-CoV-2 first detection date. Not including multiple episodes of SARS-CoV-2 per patient.

\* Additional data check required. (Resistance is not detailed, data for MERS is not currently available).

^ Definition post- SARS-CoV-2 secondary infection (SARS-CoV-2 is primary infection). This has been extended from prior 14d secondary infection definition for influenza used by PHE to account for disparities in testing throughout the 28d period after SARS-CoV-2 detection.

‡ Streptococcus species includes the following groups and species:

Group	Species/other names
Anginosus Group	<i>Streptococcus anginosus</i> ; <i>Streptococcus constellatus</i> ( <i>Streptococcus constellatus</i> subspecies <i>constellatus</i> <i>Streptococcus constellatus</i> subspecies <i>pharynges</i> ); <i>Streptococcus</i> Group F; <i>Streptococcus intermedius</i> ; <i>Streptococcus milleri</i> group; <i>Streptococcus sinensis</i>
Bovis Group	<i>Streptococcus alactolyticus</i> ; <i>Streptococcus bovis</i> untyped; <i>Streptococcus equinus</i> ; <i>Streptococcus gallolyticus</i> subspecies <i>gallolyticus</i> ( <i>Streptococcus bovis</i> biotype I); <i>Streptococcus infantarius</i> ( <i>Streptococcus infantarius</i> sp <i>infantarius</i> ; <i>Streptococcus bovis</i> biotype II); <i>Streptococcus lutetiensis</i> ; <i>Streptococcus infantarius</i> subspecies <i>coli</i> ( <i>Streptococcus bovis</i> biotype II); <i>Streptococcus pasteurianus</i> ( <i>Streptococcus bovis</i> biotype II)
Closely Related Genera	<i>Abiotrophia</i> spp.; <i>Aerococcus</i> spp.; <i>Faklamia</i> spp.; <i>Gemella</i> spp.; <i>Globicatella sanguinis</i> ; <i>Granulicatella</i> spp.; <i>Leuconostoc</i> spp.; <i>Pedicoccus</i> spp.; <i>Peptostreptococcus</i> spp.
Mitis Group	<i>Streptococcus cristatus</i> ; <i>Streptococcus mitior</i> ; <i>Streptococcus mitis</i> ; <i>Streptococcus oralis</i> ; <i>Streptococcus pseudopneumoniae</i> ; <i>Streptococcus infantis</i> ; <i>Streptococcus peroris</i>
Mutans Group	<i>Streptococcus mutans</i> ; <i>Streptococcus sobrinus</i>
Other streptococci (including but not limited to)	Anaerobic streptococcus; <i>Streptococcus acidominimus</i> ; <i>Streptococcus</i> spp., other named/not fully identified; <i>Streptococcus suis</i> ; <i>Streptococcus uberis</i>
Salivarius Group	<i>Streptococcus vestibularis</i> ; <i>Streptococcus thermophilus</i>
Sanguinis Group	<i>Streptococcus gordonii</i> ; <i>Streptococcus massiliensis</i> ; <i>Streptococcus parasanguinis</i> ; <i>Streptococcus sanguinis</i>
<i>Streptococcus</i> Group A	Group A; <i>Streptococcus pyogenes</i> ; <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i>
<i>Streptococcus</i> Group B	Group B; <i>Streptococcus agalactiae</i>
<i>Streptococcus</i> Group C	Group C; <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i> ; <i>Streptococcus equi</i> subspecies <i>zooepidemicus</i>
<i>Streptococcus</i> Group G	Group G; <i>Streptococcus canis</i> ; <i>Streptococcus dysgalactiae</i> subspecies <i>equisimilis</i>