## Α

	Medium-grade fever		Low-grade fever			Risk Ratio	Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl	
Du 2020	11	26	14	26	12.8%	0.79 [0.44, 1.39]		
Hu 2020b	34	87	43	87	19.9%	0.79 [0.56, 1.11]		
Wan 2020	35	84	43	84	20.3%	0.81 [0.59, 1.13]		
Liu 2020f	9	23	8	23	9.1%	1.13 [0.53, 2.40]		
Zhang 2020f	68	145	51	145	21.9%	1.33 [1.01, 1.77]		
Zhang 2020l	25	56	17	56	15.0%	1.47 [0.90, 2.41]		
Chen 2020a	7	10	0	10	1.0%	15.00 [0.97, 231.84]		
Total (95% CI)		431		431	100.0%	1.04 [0.79, 1.37]	-	
Total events	189		176					
Heterogeneity: Tau <sup>2</sup> =	0.07; Chi² = 14.4	46, df = 6	$(P = 0.02); I^2$	= 59%				
Test for overall effect: $Z = 0.27$ (P = 0.78)							0.5 0.7 1 1.5 2	

Test for overall effect: Z = 0.27 (P = 0.78)

0.5 0.7 1 1.5 2 Low-grade fever Medium-grade fever

## В

	Low-grade	fever	High-grade	fever		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% C	M-H, Random, 95% Cl
Chen 2020a	0	10	3	10	4.1%	0.14 [0.01, 2.45]	
Zhang 2020l	17	56	14	56	18.4%	1.21 [0.66, 2.22]	- <b></b>
Liu 2020f	8	23	6	23	15.5%	1.33 [0.55, 3.24]	
Zhang 2020f	51	145	26	145	20.2%	1.96 [1.30, 2.96]	
Hu 2020b	43	87	10	87	18.2%	4.30 [2.31, 8.00]	
Wan 2020	43	84	6	84	16.4%	7.17 [3.22, 15.93]	
Du 2020	14	26	1	26	7.2%	14.00 [1.98, 98.85]	
Total (95% CI)		431		431	100.0%	2.50 [1.32, 4.73]	<b>•</b>
Total events	176		66				
Heterogeneity: Tau <sup>2</sup> =	0.48; Chi <sup>2</sup> = 3	25.95, df	= 6 (P = 0.00	002); l <sup>2</sup> =	77%		
Test for overall effect:	est for overall effect: Z = 2.81 (P = 0.005)						0.01 0.1 1 10 100 High-grade fever Low-grade fever

## С

	Medium-grade	e fever	High-grade	fever		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% Cl
Liu 2020f	9	23	6	23	12.0%	1.50 [0.64, 3.53]	
Zhang 2020l	25	56	14	56	20.1%	1.79 [1.04, 3.06]	
Chen 2020a	7	10	3	10	9.2%	2.33 [0.83, 6.54]	+
Zhang 2020f	68	145	26	145	25.6%	2.62 [1.77, 3.86]	
Hu 2020b	34	87	10	87	17.0%	3.40 [1.79, 6.44]	
Wan 2020	35	84	6	84	12.9%	5.83 [2.59, 13.13]	
Du 2020	11	26	1	26	3.1%	11.00 [1.53, 79.16]	
Total (95% CI)		431		431	100.0%	2.72 [1.89, 3.90]	•
Total events	189		66				
Heterogeneity: Tau <sup>2</sup> =	0.09; Chi <sup>2</sup> = 10.5	51, df = 6	$(P = 0.10); I^2$	= 43%			
Test for overall effect:	Z = 5.41 (P < 0.0	00001)					0.05 0.2 1 5 20 High-grade fever Medium-grade fever

S11 Fig. Risks of (A) low-grade fever  $(37\cdot3-38\cdot0^{\circ}C)$  vs medium-grade fever  $(38\cdot1-39\cdot0^{\circ}C)$ , (B) high-grade fever  $(>39\cdot0^{\circ}C)$  vs low-grade fever  $(37\cdot3-38\cdot0^{\circ}C)$ , and (C) high-grade fever  $(>39\cdot0^{\circ}C)$  vs medium-grade fever  $(38\cdot1-39\cdot0^{\circ}C)$  in adult non-severe COVID-19 patients.