## Α

	Medium-grade	fever	Low-grade	fever		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	I M-H, Random, 95% CI
Fan 2020	42	91	33	91	55.4%	1.27 [0.90, 1.81]	+=-
Li 2020a	34	59	17	59	44.6%	2.00 [1.27, 3.16]	<del></del>
Total (95% CI)		150		150	100.0%	1.56 [1.00, 2.42]	•
Total events	76		50				
Heterogeneity: Tau <sup>2</sup> =	0.06; Chi <sup>2</sup> = 2.37	0.1 0.2 0.5 1 2 5 10					
Test for overall effect: Z = 1.97 (P = 0.05)							Low-grade fever Medium-grade fever

## В

	Low-grade	fever	High-grade	fever		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
Fan 2020	33	91	16	91	67.9%	2.06 [1.22, 3.47]	_ <b></b>
Li 2020a	17	59	8	59	32.1%	2.13 [0.99, 4.54]	-
Total (95% CI)		150		150	100.0%	2.08 [1.35, 3.20]	•
Total events	50		24				
Heterogeneity: Tau <sup>2</sup> =	0.00; Chi <sup>2</sup> = 0	0.2 0.5 1 2 5					
Test for overall effect:	Z = 3.34 (P =		High-grade fever Low-grade fever				

## С

	Medium-grade	efever	High-grade	fever		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI	M-H, Random, 95% CI
Fan 2020	42	91	16	91	62.0%	2.63 [1.60, 4.32]	
Li 2020a	34	59	8	59	38.0%	4.25 [2.15, 8.39]	
Total (95% CI)		150		150	100.0%	3.15 [1.99, 4.99]	•
Total events	76		24				
Heterogeneity: Tau <sup>2</sup> =	0.02; Chi <sup>2</sup> = 1.27	0.1 0.2 0.5 1 2 5 10					
Test for overall effect: $Z = 4.89 (P < 0.00001)$							High-grade fever Medium-grade fever

S13 Fig. Risks of (A) low-grade fever (37·3-38·0°C) vs medium-grade fever (38·1-39·0°C), (B) high-grade fever (>39·0°C) vs low-grade fever (37·3-38·0°C), and (C) high-grade fever (>39·0°C) vs medium-grade fever (38·1-39·0°C) in non-survived adult COVID-19 patients.