

Deployment of Resilient Foods Can Greatly Reduce Famine in an Abrupt Sunlight Reduction Scenario

Morgan Rivers (✉ morgan@allfed.info)

ALLFED

Michael Hinge

ALLFED

Juan García Martínez

Alliance to Feed the Earth in Disasters <https://orcid.org/0000-0002-8761-7470>

Ross Tieman

Alliance to Feed the Earth in Disasters (ALLFED)

Victor Jaeck

ETH Zürich

Talib Butt

Northumbria University

David Denkenberger

University of Alaska Fairbanks <https://orcid.org/0000-0002-6773-6405>

Article

Keywords:




Posted Date: April 1st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1446444/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Deployment of Resilient Foods Can Greatly Reduce Famine in an Abrupt Sunlight Reduction Scenario

Morgan Rivers ^a, Michael Hinge ^a, Juan B. García Martínez ^a , Ross J. Tieman ^a , Victor Jaeck ^b, Talib E. Butt ^{a,c}, David C. Denkenberger ^{a,d} 

Abstract

In a nuclear war, volcanic eruption, asteroid or comet impact that causes an abrupt sunlight reduction scenario (ASRS), agricultural yields would plummet. Global society is currently unprepared for such an event, implying an urgent need for evaluation and prioritization of solutions. We show effective deployment of resilient food solutions appears sufficient to fulfill global energy and macronutrient food requirements, potentially saving billions from famine. A Monte Carlo analysis of resilient food outcomes, using a linear optimization model, shows a 95% probability of global food availability between 2,100 and 3,500 Kcals per capita per day in a nuclear winter scenario involving 150 Tg of soot in the stratosphere. Our analysis indicates nutritionally sufficient diets from resilient foods would be widely affordable, costing US\$1.73 daily, though subsidization could be needed across Southern Asia and Sub-Saharan Africa. Post-disaster conflict or insufficient international cooperation could increase costs and reduce output, hampering effective resilient food deployment.

1 Main

The global population is unprepared for several food-related global catastrophic risks (GCRs). We classified a subset of extreme food-related GCRs under the label *abrupt sunlight reduction scenarios* (ASRS), in which a catastrophic event causes a widespread, rapid reduction in sunlight reaching the surface of the Earth. At least three mechanisms for ASRS have been identified in the literature: extreme volcanic eruption, large bolide (asteroid/comet) impact, and nuclear war^{1,2}. In these scenarios, an enormous sudden ejection of aerosol material such as sulfates or soot (black carbon) takes place, causing multi-year reductions in global temperature, solar irradiation, and precipitation, leading to a global catastrophic food shock (GCFS). Nuclear war has a likelihood of ~1% per annum³, while large bolide impact has a likelihood of ~0.0001% and supervolcano eruption has a likelihood of ~0.001%⁴. Global conflicts such as the 2022 Russo-Ukrainian crisis may increase the annual probability of nuclear war above these levels.

Prevention of an ASRS is unambiguously the best outcome. However, this may be impossible, and according to the “three layers of defense” model of existential risk, a comprehensive strategy ought to include prevention, response and resilience⁵. The current work investigates the feasibility of response and resilience approaches to an

extreme ASRS, in line with the recent United Nations calls for “Defining, identifying, assessing and managing existential risks”^{6,7}.

Xia et al.⁸ have estimated a global fatality rate of 75% due to starvation in an ASRS catalyzed by a nuclear winter involving 150 Tg of soot in the stratosphere (“150 Tg scenario”), but did not evaluate scenarios involving effective global measures taken to bridge the food production shortfall. In this work, we investigate a subset of foods and food production methods known as *resilient foods* or *resilient food solutions*, which could allow for significant food production in the face of an ASRS⁹. Resilient food solutions must be scalable and amenable to rapid production ramp-up. Solutions must also be low cost, as affordability is a key factor for adequate access to food during an ASRS, just as it is today¹⁰.

In this study we investigate the reduction in global famine in scenarios involving a globally coordinated response and approximately \$30 million to \$300 million in preparation prior to a 150 Tg scenario. The primary difference between a 150 Tg nuclear winter scenario and a comparable volcanic eruption is the higher-altitude lofting in the stratosphere of soot emanating from firestorms induced from the nuclear blast, prolonging the duration of a nuclear ASRS¹¹.

We begin our analysis by analyzing two 150 Tg scenarios: one involving no response, and another involving plausible outcomes of the deployment of a suite of resilient food solutions. Based upon these estimates we find that resilient foods could feed everyone throughout the 150 Tg scenario. A Monte Carlo simulation of 10,000 different scenarios confirms our results. The relative caloric contribution of each resilient food is compared in a separate set of 1,000 scenarios. Finally, we compare resilient food costs and overall nutritive contributions to common low-cost foods in 2020. Expectations of the performance of some resilient foods are still preliminary. However, as new research refines expected resilient food outcomes, and as the food system changes over time, updated or new data can be included in the open source model for scenario analysis and easily run online using the publicly available Jupyter notebook Colab interface.

2 Results

2.1 Feasible Global Macronutrient Production

150 Tg Scenario, Without Resilient Foods

In the face of an ASRS such as a 150 Tg scenario or an equivalent volcanic eruption, human edible food from agriculture would undergo near-complete collapse, with a minimum production of only 11% of current levels after three years, given current behaviors⁴. Here we evaluate the degree of food availability in a scenario assuming continued global trade, not including disruptions due to the ongoing or additional conflicts, and assuming a relatively rapid reduction in human edible foods fed to livestock.

In the event of continued coordination, several aspects of the food system would allow for more food availability than implied from an 11% baseline. As a result of the readily apparent severity of the disaster, even with no planning we would expect that the majority of biofuel production would halt quickly, due to the rapid expected rise in food prices relative to fuels. Because continuation of livestock at present-day levels would imply fewer animals would starve than humans, and economic incentives would increase the cost of feed, we estimate the majority of human edible feed fed to animals would be redirected. As the disaster progresses, livestock would also become a key source of macronutrients, as most livestock would be consumed rather than maintained with expensive human edible food. Finally, in part due to soaring food prices relative to incomes, we estimate that waste (which includes "overconsumption" beyond minimum healthy levels) would be reduced sharply. Based on an estimated tripling of food prices, and a consumer response to this based upon individual incomes, we estimate post-harvest waste would fall to approximately 13% of production¹². See Supplementary Information section I for the food system baseline before catastrophe.

Assuming some continued food waste, a two-month delay before shutoff of biofuel production, and a three-month delay before human edible animal feed is redirected to humans, only 1,600 Kcals per capita per day are available throughout (**Figure 1**). The simulation indicates that a full 6.5 years (78 months) is required to increase food availability significantly above this level. The models indicate that calories are the primary nutrient limitation throughout, and that sufficient dietary fat and protein would be available to meet the needs of those who could obtain enough calories with a mixed diet. A rough estimation of the survival rate obtained by dividing the proportion of available calories by the minimum recommended caloric consumption (of 2,100 Kcals¹³) indicates approximately 24% of the global population would starve even if food were allocated optimally over time.




Figure 1 | Food availability after accounting for waste and a delayed halt of non-human consumption, without resilient foods. a, Expected per capita caloric availability by food source in a 150 Tg scenario with no resilient food deployment. 13% waste is assumed, with a two month delayed biofuel shutoff and a three month delay before human-edible animal feed is redirected to humans. **b,** Fraction of macronutrients on average available during the 150 Tg scenario with no resilient foods. Recovery of available food past 1,600 Kcal only occurs after 6.5 years under a wide range of assumptions, and in all cases quickly recovers past minimum present global needs. Waste was estimated using the probable reduction in waste if there were a 3:1 ratio of food price to 2020, as described in more detail in the methods section. The change in percentage waste was not delayed in the model. See Supplementary Information for detailed information on each food source.

150 Tg Scenario, With Resilient Foods

The successful implementation of resilient foods would make some coordination assumptions in the previous section more plausible. With food production combined with waste well above minimum human requirements throughout, stored food would not be in as extreme demand in the worst years of the 150 Tg scenario. Preparedness plans for resilient foods would imply a situation in which the industrial producers and decision makers are aware of the critical urgency of rapidly culling and storing livestock and reducing human edible food fed to animals and used for biofuel production. It is therefore assumed to take one month to halt biofuel production and two months to redirect

substantial fractions of animal feed to humans. International action to coordinate deployment of these foods, combined with strong international food subsidies for the world's poor, could allow sufficient nutrition in all areas able to trade and receive food. In this scenario, conflicts from food riots and resource competition would thus be greatly reduced.

The resilient food solutions with maximum life-saving potential in a 150 Tg scenario were identified as a combination of the following:

- The optimal release and equitable distribution of all forms of stored food, both nationally and internationally.
- Outdoor crops with high levels of fertilizer applied to croplands in the tropics and with rotations altered to accommodate the colder climate.
- Continued dairy output near present-day levels via prioritization of grasses, fodder crops and residues for dairy cattle.
- Continued meat production where output does not conflict with human consumption, while prioritizing grazing of ruminants on any available pasture beyond those needed for dairy.
- A seaweed industry that is rapidly scaled up to approximately 70x present-day farm area and production quantities.
- Rapid expansion of low-tech greenhouses in the tropics¹⁴.
- The rapid repurposing of paper and pulp factories into cellulosic sugar factories, (an "industrial food").
- The rapid establishment and deployment of single cell protein (SCP) factories^{15,16}, an ("industrial food").
- Continued commercial marine fishing with reduced catch yields as the ASRS progresses⁴.

For this scenario, the food production potential of resilient food solutions implemented successfully plus traditional agricultural production is estimated at approximately 2,700 Kcal per capita throughout the 150 Tg scenario after incorporating waste, feed, and biofuel production (**Figure 2**).

The primary contribution to maintain fat and protein requirements involved a selection of rotations involving increased planted area for wheat as a protein source, rapeseed for fat, and sugar beets or potatoes for caloric supply. In this scenario, caloric demand is the most important limiting factor. Seaweed met significant protein demand, due to the use of *Porphyra amplissima* with its high protein to calorie ratio. Protein was also significant from Methane SCP.




Figure 2 | Food availability after accounting for waste and a delayed halt of non-human consumption assuming resilient food scaling up in a 150 Tg scenario. a,b: Availability of food by food source, incorporating approximately 18% food waste, including overconsumption and excluding on-farm waste, and with current biofuel production

continued for one month and current animal feed usage continued for two months. **c,d:** An approximate average ASRS-resilient diet composition determined by feeding excess production from outdoor crops to livestock. In this scenario, dairy could be fully maintained, and other animal products could be maintained at significant volumes by prioritizing the most efficient feed systems. The change in percentage waste was not delayed in the model. See Supplementary Information for detailed information on each food source.

By month 48 after the ASRS onset, resilient foods would be fully scaled up, and traditional crop yields would pass their minimum point and begin to recover to normal levels over the coming years. This would allow diets to slowly revert to more traditional foods. Finally, the average human diet shown could include less seaweed and industrial foods if these foods were fed to animals rather than humans, although this has not been investigated.

Monte Carlo Simulations

We performed a set of Monte Carlo simulations to estimate the variability in resilient food outcomes. A random sampling of 10,000 scenarios over the parameter space was selected. The 2.5th percentile outcome scenario produced 2,100 Kcals per capita per day after waste, and the 97.5th percentile produced 3,500 Kcals per capita per day after waste. Calories remained the limiting factor in the majority of Monte Carlo scenarios. See the Supplementary Information, section II for details.

We also ran a separate set of 1,000 scenarios with either only one resilient food scaled up or only one resilient food removed. These scenarios reveal that if fat and protein minimum requirements are enforced, each individual resilient food is less effective than when scaled up in combination with other resilient foods. Relocation contributed to meeting macronutritional needs most when scaled in combination with other foods.




Figure 3 | Food available after accounting for delayed shut off and waste. **a** Estimated caloric contribution per person for each resilient food over 1,000 scenarios. Scenarios where the resilient food is removed from the full suite of resilient foods are shown in red. Scenarios where only one resilient food is scaled up is indicated in green. Only calories used as part of a balanced macronutrient profile are shown. Reduced variability is observed when adding foods, as evaluating the variability of scenarios in the non-resilient food case is outside the scope of this paper. Red bars are relative to scenarios similar to **Figure 2a** while green bars are relative to scenarios similar to **Figure 1a**. **b** Estimated caloric availability in 10,000 scenarios after incorporating a constant level of reduced waste of 18% due to increased food price, delayed biofuel and feed, and differences in the production and scaling up of resilient foods. While caloric availability is shown, fat and protein limitations are often influential in determining the change in number of useful calories for a scenario.

While many challenges remain before they are ready for rapid deployment at scale, solutions such as industrial foods and seaweeds could possibly scale to 2x or more of current global human minimum caloric requirements within the first few years after a 150 Tg scenario onset, given sufficiently large portions of global industrial and economic resources¹⁵⁻¹⁹.

2.2 Affordability of Macronutrient Consumption

Economic as well as physical access to foods is required in order to meet the criteria of food security, as defined by the Food and Agriculture Organization of the United Nations (FAO) in the 1996 World Food Summit²⁰. While a detailed analysis of price equilibria is beyond the scope of this paper, here we present an analysis of the cost of producing resilient foods post-disaster under a median production resilient food production scenario. This analysis allows us to determine the minimum cost of diets, as prices must exceed costs to be sustainable over the duration of the ASRS, especially with rapid capacity expansions. See Supplementary Information, section III for details.

Figure 4 shows the estimated costs three years subsequent to the disaster when outdoor crop production is at its highest cost, including an allowance for a return on capital and depreciation. Costs have been normalized onto a US\$/day, or 2,100 Kcal, basis. Our results indicate significant variation in the cost of foods in the ASRS. This ranges from around US\$0.30/2,100Kcal for cellulosic sugars, up to US\$0.60-0.90 for outdoor grown grains (wheat and maize), US\$0.95 for rapeseed oil, around US\$2.70 for seaweed and potatoes, and around US\$6.00 for milk vs US\$7.50-15.00 for chicken/pork and ruminant meat respectively.




Figure 4: Cost of production on a retail basis for a sample of foods, before and after disaster. Costs are listed as US\$ per 2,100 Kcal, estimated three years after 150 Tg scenario onset. These costs represent a significant rise when compared to current retail prices, with increases in bulk grain prices such as wheat and maize by around 250% and vegetable oils by 300%. This suggests a significant decline in food affordability post-disaster, pushing billions into food insecurity unless additional support is provided. Sources of foods that generate more calories per unit area, such as potatoes and greenhouse crops, are less affordable, ranging from US\$1.5-3 per 2,100 Kcal, versus around

US\$0.2-0.3 for the cheapest grains today. The retail premium refers to the estimated cost of moving foods from farm/factory gate to a shop available for sale to consumers, as well as a retail margin. Error bars indicate the range of high/low production costs based upon the variance in minimum and maximum yields over the period of calibration (see Supplementary Information, section III).

In order to provide an estimate of affordability post-disaster, we constructed four representative diets out of the proposed set of foods subject to meeting the constraints listed below. While diets typically do not match well to richer consumers, they are much closer to the choices made by consumers facing significant budgetary constraints²¹, and are therefore useful when considering affordability and diets for those in or close to global poverty. We compared the cost of these diets to global poverty thresholds by region²², based upon market exchange rates and spending no more than 63% of total income on food²³ (see Supplementary Information, section III).

Potential diets have been presented on four different levels of affordability:

- A "minimum cost caloric" diet costing US\$0.51 per day, based upon sourcing 30% of calories from cellulosic sugars and the remaining 70% from the next lowest-cost source, maize. However, this would be well below other macronutrient and micronutrient intake recommendations.
- A "minimum cost macronutrient" diet for protein and fat as well as carbohydrates costing US\$0.80 per day, achieved by substituting some grains and sugars for single cell proteins, seaweed and vegetable oils in the diet above.
- A micronutrient-complete diet as laid out in Pham et al.⁹, which presents diets constructed out of foods projected to be available following a ASRS. These satisfy minimum and maximum micronutrient (vitamin/mineral) bounds as well as macronutrients. There are a number of diets in the study, and we have chosen the period III diet (once resilient foods are scaled), excluding greenhouses, to get the lowest-cost diet satisfying full nutrient requirements in year three. In total, this diet is projected to cost \$1.73 daily. However, while the paper examined low cost diets post-disaster it did not minimize costs, and as such it is possible micronutrient requirements could be met at a lower cost than this estimate.
- An average "basket of foods" diet costing US\$2.56 per day, weighted evenly on a caloric basis by its share of the total, including higher-cost foods such as animal products and greenhouse cultivated crops. This diet meets minimum calories, protein and fat requirements. It however has not been assessed for micronutrients.

The affordability of each of these diets is presented in Table 1. While most could afford the diets on their pre-disaster income, at least 1.1 billion would be unable to afford a basic caloric diet at cost post-disaster, 1.35 billion would be unable to afford adequate macronutrients, just over 3 billion might struggle to consume enough vitamins and minerals without supplementation/fortification and over 3.7 billion would be unable to consume the average basket of foods produced.

Table 1: Affordability of dietary benchmarks post disaster - Year 3 - US\$ international exchange rates

Cheapest calories	Cheapest macronutrients	All nutrients met	Average diet
-------------------	-------------------------	-------------------	--------------

	met									
Daily cost per capita	\$0.51			\$0.80			\$1.73			\$2.56
Necessary income per day at 63% of budget	\$0.81			\$1.28			\$2.75			\$4.07
Region	Unable to afford calories		Unable to afford macronutrients		Unable to afford micronutrients		Unable to afford average diet			
	<i>million people</i>	<i>%</i>	<i>million people</i>	<i>%</i>	<i>million people</i>	<i>%</i>	<i>million people</i>	<i>%</i>		
East Asia and Pacific	17	0.7%	31	1.3%	422	17.9%	790	33.5%		
Europe and Central Asia	19	2.1%	22	2.3%	49	5.3%	83	9.0%		
Latin America and the Caribbean	17	2.6%	21	3.3%	84	12.8%	140	21.5%		
Middle East and North Africa	6	1.2%	10	2.1%	122	26.3%	195	41.9%		
Other high Income	4	1.0%	4	1.0%	6	1.6%	8	2.3%		
South Asia	644	34.7%	822	44.3%	1,557	83.9%	1,634	88.0%		
Sub-Saharan Africa	372	32.7%	440	38.7%	813	71.5%	892	78.5%		
World Total	1,078	13.9%	1,348	17.4%	3,053	39.3%	3,742	48.2%		

This affordability issue suggests that extensive support involving income transfers, food coupons or price subsidies would be needed post-disaster in order to ensure sufficient nutrition for a section of the global poor. International support post-disaster would be necessary across Sub-Saharan Africa and South Asia, particularly for the urban poor.

Affordability on a cost basis is an important first step in determining economic access to food, as it sets the minimum prices paid by consumers. Prices could however be significantly higher than this level, as described in the Discussion section. Meanwhile, in the case of a complete or partial nationalization of the food system, costs are still a relevant consideration for the capital, labor and inputs needed to produce each form of nutrition.

The same method to calculate affordability as is used above applied to current prices suggests that around 169 million people would not be able to afford sufficient macronutrients from 2015 to 2019. This compares to an average of 642 million people reported to be in severe food insecurity over the same period by the FAO¹⁰, with the difference due in large part to factors beyond direct macronutrient affordability on a cost basis, including conflict-induced food insecurity and failures in infrastructure. As such, these additional factors have significant implications for food security post-disaster, and more analysis is needed to understand how they would apply.

3 Discussion

Maintaining sufficient production of nutritious and affordable foods is a commonly identified element of food security²⁴, but some have challenged the centrality of production to modern food systems thinking²⁵. Global food shock analyses typically involve only a few percentage points lower production than expected trends in global output^{26,27,28}. In many ASRS outcomes, even with optimal allocation and distribution, sufficient access for the world's population would be impossible. Thus, we focus on food production and affordability as our primary form of analysis in this paper, while still acknowledging the vital importance of broader factors such as equitable food allocation and distribution in both ASRS and other food shocks²⁵.

Key future work to prevent global famine from lack of production in an ASRS includes but is not limited to: 1) research on food production methods, production ramp-up and technology deployment, as well as nutritional qualities of the foods, 2) further development of technologies and techniques conducive to a faster response such as fast construction and rapid repurposing, and 3) policy outcomes such as the creation and distribution of effective disaster response plans. Arguably, these areas of work can also inform preparedness and resilience against less-extreme global catastrophic food risks, such as a multiple bread-basket failure due to other forms of climate, or weather shocks, or crop diseases and pests.

While convenient for modeling purposes, the assumptions of continued global trade and other forms of international cooperation may not hold in a world where billions more people are at risk of starvation. Export bans, food riots, resource competition, and ensuing global conflict are potential outcomes, which would significantly increase famine compared to our analysis above. In a 150 Tg scenario, denial of the severity of the ASRS would likely be limited after several months, as the sky would appear dark globally in the months following the global catastrophic event^{29,30}, and outdoor crop production is estimated to fall to roughly half of present-day levels after only 12 months without relocation. This could lead to widespread panic without a credible response, significantly complicating the prospect of efficient global rationing and allocation of food stocks. By contrast, demonstrating that widespread starvation post-disaster is not a guaranteed outcome could prevent panic and counterproductive negative-sum actions.

It may also be that rapid shutoff of non-human sources of demand such as biofuels and large scale animal feed systems would not occur without sufficient regulatory legislation in advance of the ASRS. We therefore recommend including contingencies in biofuel legislation to quickly phase out food-based biofuel production in case of a GCFS.

The sociotechnical systems which constitute the global food system are complex in nature and will exhibit nonlinear dynamics as system variables change^{31,32}. Complex systems often exhibit tipping points – thresholds which once surpassed result in a conformational change of the system to another state through positive feedback loops^{33–35}. Aggregate food availability below the global population's nutritional needs may represent such a tipping point.

The Monte Carlo analysis implies that with a coordinated resilient food production response there is a low likelihood for output to dip below this hypothetical tipping point over a wide range of resilient food outcomes. Strengthening food resilience and response capabilities prior to an ASRS could thus be critical to prevent positive feedback loops. The results of our analysis demonstrate that it is possible that the current global population could be fed even in a severe ASRS, although at a higher cost compared to current resources devoted to food production. As a result the world could continue global trade in order to reduce the severity of the crisis.

Global GDP dedicated to agriculture, forestry and fishing was in excess of 9% as recently as the 1970s³⁶, and while accurate estimates of GDP further back are hard to make, employment in agriculture was dominant for much of human history³⁷. Meanwhile, agriculture, forestry and fishing is only 3.5-4% of GDP in the present day by World Bank World Development Indicator estimates³⁸. We estimate an expansion of resilient foods would require an additional 3.7% of global GDP based on their capital and operating costs (see the supplementary information for details), raising the total of GDP dedicated to agriculture, forestry and fishing to around 7-7.5%. As a result, while GDP dedicated to food production would need to increase sharply during an ASRS, such an increase may still be feasible and could still be within the upper bounds of living memory.

Despite this potential to meet nutritional needs, it is far from certain that cooperation would actually be secured or maintained, and other social and economic factors could also impact the post disaster response. Producing surplus nutrition is a necessary but not sufficient step in securing global cooperation and meeting individual needs. Expectations for the outcome of a severe ASRS incorporating these considerations are the topic of future work. As a result it is possible that as understanding of the relevant resilient food solutions and systems improve, the conclusions listed may change.

Additional modeling challenges are involved when considering social instability and calculating price equilibriums where output of vital staples are disrupted and future production is highly uncertain. Accordingly, full modeling of food markets and prices under such a scenario are left to future work. However, standard financial systems could experience severe disruption and could potentially collapse under such a scenario, even with continued trade. In addition, there are a number of factors that could alter our cost results, including import and consumption taxes as well as breakdowns in infrastructure. In addition, conflict and continued demand from biofuels or animal feed have the potential to raise prices further post-disaster, especially in the early months.

The ability to coordinate between trading blocs is uncertain following a GCFS and will significantly impact global capacity to respond. In certain types of ASRS, infrastructure could be affected such that countries may no longer be able to communicate and trade internally or externally after the GCFS³⁹. In a nuclear exchange, international trust may have eroded as well. In the context of outdoor crop production, this would reduce the ability to send agricultural inputs located in the Global North to tropical regions which could best use them, and for excess yields to be distributed to deficit zones.

Finally, transitioning to inherently more resilient food production systems prior to shocks (e.g., closed-environment systems and modular designs) would also be conducive to warding off the impacts of ASRS^{15,16} as well as other food system⁴⁰ risks. In a similar vein, expanding use of seaweed as a food and feed⁴¹ source could directly draw down CO₂, and reduce cattle methane emissions, improve food security around the world today by its use

as a low-cost food source⁴², and reduce the food security impacts of trade restrictions⁴⁰, pests⁴⁰, ASRS⁴³, and a loss of electrical and/or industrial function^{44,45}.

While pathways for prevention of massive volcanic eruption⁴⁶ or bolide impact⁴⁷ require more research, there is an urgent and well-established need for large-scale global nuclear arms reduction to small fractions of the current global arsenal⁴⁸, in part because nuclear war continues to pose a risk for large-scale societal collapse or human extinction. We encourage work along these lines, in parallel with ongoing research and preparations for a GCFS should the worst occur.

We recommend business continuity managers and decision makers working in disaster risk management to promote policies for the creation of preparedness and response plans against GCFS, as has been done for other high-impact low-probability risks, both natural (e.g., tsunamis)⁴⁹ and anthropogenic (e.g., nuclear plant accidents)⁵⁰, even at the international level⁵¹. These regional preparedness arrangements could complement systems such as the international "emergency platform" proposed by the UN Secretary-General for responding to GCRs⁷ such as those characterized in this work.

4 Methods

The model presented was designed to determine the feasibility of feeding everyone through deployment of resilient foods. Details of the linear optimization model can be found in the Supplementary Information, section IV. All resilient food production methods evaluated here have been deployed at a large scale at the time of writing. As low-cost foods are included preferentially, economic considerations are then applied to estimate how many people could likely afford the food produced. Resilient foods were selected according to anticipated resource constraints (e.g., industrial capacity, availability of production inputs) and the value of the food output – roughly macronutrient content (i.e., calories, fat, protein) per unit cost.

Previously estimated growth rate models are used for estimating the growth rate of industrial foods^{8,15}. We account for the management of food reserves and livestock carried in from before the onset of a 150 Tg scenario, as well as redirection of crops away from biofuel production and animal feed. Food waste and estimated reductions of waste at each stage (production, distribution, retail, and consumption above minimum requirements) are also accounted for.

Key assumptions used in the model include:

- Continued global trade - Trade is maintained and production activities are coordinated to maximize total output.
- No hoarding or looting of food at the household, retail or wholesale level.
- Financial stability is assumed
 - Second order analyses and economic effects of the likely financial system disarray are not considered.

- Income is assumed to be in line with current levels and distributions.
- Infrastructure intact - No direct destruction of infrastructure/industry is considered.
- Population maintained - Reductions in population due to direct or indirect effects are not considered in the results.

Key additional assumptions for the resilient foods section include:

- Sufficient preparation and knowledge in appropriate sectors on how to implement resilient foods.
- Key relations between organizations being put in place early after onset and remaining functional throughout the food shortage.
- Appropriate supporting industries are spun up starting early after the onset.

Future work in scenario analysis will study the effect of varying these assumptions.

Population Energy and Macronutrient Requirements

Energy and macronutrient requirements represent the minimum recommended daily intake for an average weight adult (62 kg): Energy - 2,100 Kcal/day¹³, Protein - 51 g/day and Fat - 47 g/day⁵². This is in line with previous assessments of GCFS^{9,15-18}. Quality of protein (i.e., the full complement of amino acids in sufficient proportions), and quality of fat (i.e., sufficient unsaturated fat, and omega-3 and omega6 fatty acids) are beyond the scope of this paper, although baskets of the same resilient foods as discussed in this paper have been calculated to provide a protein-complete diet⁹. Resilient food diets appear to be able to largely meet key macronutrient and micronutrient requirements at a population scale with adequate nutritional planning⁹. Therefore, meeting specified daily quantities of macronutrients is considered nutritionally sufficient within the model. Future work will combine food availability and nutrition analyses.

Waste

Waste information was determined using the FAOSTAT Supply Utilization Accounts database between 2014 and 2018⁵³. Over one third of current agricultural production on a caloric basis goes to waste before coming to any use. Agricultural waste consists of harvest losses, distribution losses, and retail/household waste. Harvest losses are already accounted for in the estimates of current-day agricultural production and were not adjusted in the ASRS, which is pessimistic as there would be effort to reduce harvest losses given the higher prices. Distribution losses refer to losses in processing, transit and storage, post-farm but before they are delivered to the retail level. Distribution losses are largely a function of existing quality of storage and transport infrastructure⁵⁴, and are assumed to be maintained. Distribution losses vary widely by crop/food variety, and so the percentage loss appropriate for each agricultural category in FAOSTAT is assumed⁵³. Retail/household waste refers to food damaged or not consumed at the retail level onwards, such as shops rejecting or failing to sell products or households discarding food once purchased.

In order to estimate waste in the “no resilient foods” case, a tripling of food prices is assumed. In the “no resilient foods” case, applying empirical linear extrapolation between food price and waste¹² and aggregating over all countries, retail and household waste were estimated to be reduced from 24% to 14% globally post-catastrophe in the resilient foods scenario, and from 24% to 10% in the scenario without resilient foods.

Stored Food

Available stored food at the time of the disaster has been estimated based upon reported crop year end stocks from the United States Department of Agriculture’s Production, Supply and Distribution (USDA PSD) Database⁵⁵. These have been adjusted onto a calendar year basis (as crop years and calendar years do not typically align), and takes the month of May as its starting point: the month of the abrupt sunlight reduction in the climate model (see Supplementary Information, section V for details of this process).

This gives total stored food of around 1.65 billion tonnes, dry caloric value. This considers storage of all bulk grains, sugar, soybeans and other oil crops, as well as any vegetable oils. Data on storage for fruits, vegetables and tubers are not available in the database and so have not been included; however, these are likely to be small by comparison due to their higher perishability, and this presents a more conservative view of stored foods. In order to match production and consumption in the present day, stored food was assigned the same dietary fat and protein percentages as outdoor crop production.

Allocation of Output

In the cases of validation of the run model and 2020 data, once nutrient flows have been optimized, they are allocated to an end use, firstly direct human consumption, followed by any surplus outdoor crop production going to animal feed. Human consumption is assumed to equal the total population, multiplied by the dietary needs per person and taking into account distribution and retail/household waste discussed above.

Any surplus above this threshold is assigned to animal feed systems (see Supplementary Information, section VI). In our allocation, we have assumed that long-term use of foods for other uses such as biofuels is negligible. Prices of crops would be high post-disaster, likely making the use of biofuels highly uneconomical, and it would be expected that biofuels policy would also adjust to reflect the new reality of food scarcity.

4.1 Resilient Food Modules and Interactions

Relocated Outdoor Growing

The severity of a 150 Tg scenario with no resilient foods was estimated by scaling down year 2020 production in yields using the analysis of Xia et al⁴. Because it is expected that the majority of crop production in a 150 Tg scenario would come from currently tropical areas, the seasonality of production in the tropics (defined as 24° north to 24° south) was used as a proxy for monthly variation in global production throughout the time period.

For the relocated case simulation, sugar beets and potatoes produced the most calories per hectare, rapeseed produced the most fat per hectare, and wheat produced the most

protein per hectare. An 80% reduction in planting area of crops outside of these staples was assumed, and the remaining area would be planted with a combination of potatoes, rapeseed, wheat, and sugar beet. Under the assumption that global fertilizer production would be largely unchanged and allocated efficiently to tropical crop growing regions without the need for increased fertilizer production, crop models were run with a uniform 100 kg/ha of elemental nitrogen equivalent applied to all cropland, implying low nitrogen stress growing conditions. While these crops would not be the only crops able to be grown outdoors post-disaster, they represent a reasonable high yield, cold-tolerant rotation for macronutrients, which would be likely combined alongside some ongoing cultivation of maize, rice, pulses and vegetables where possible. As a result, the one standard deviation lower bound is the current global distribution applied to the tropics existing cropland, while the one standard deviation upper bound assumes full adoption in existing food cropland. Expanding arable land would in principle be feasible over the course of 150 Tg scenario, but it was conservatively not included in the analysis.

The varieties and types of relocated crops were instrumental in meeting fat and protein requirements. Rapeseed was found to be an efficient source of fat for a 150 Tg scenario. It is assumed that the protein from rapeseed meal would not be used to feed humans, as rapeseed meal is not deemed to be safe for human consumption⁹ and it is not guaranteed a technological solution would be achieved in a reasonable timeframe. Wheat was the largest contributor of protein per hectare in the relocated crop model, closely followed by soybean. However, soybean produced many fewer calories per hectare in the relocated case and its area was reduced compared to the present day to allow for other crops to be grown.

The improvement in yields was used to linearly scale the reduction in crop yields over time, such that the biggest improvement in rotation was three years after the GCFS, but the fully recovered yields would remain unchanged from the 2020 crop yields projection. Outdoor growing yields remain exogenous to the optimization model, although excess production from any month is stored and redistributed in months after the excess production occurs. More details on outdoor growing can be found in Supplementary Information, section VII.

Greenhouses

While greenhouses today feed a negligible number of people, low-tech polymer-based greenhouses have been estimated to provide a large fraction of calories as a resilient food source in a 150 Tg scenario at an affordable price in past work¹⁴. While these low-tech greenhouses reduce the CO₂ levels, air circulation, and incoming photosynthetically active radiation (PAR) for crops, they increase the average temperature, humidity, and thus increase available *growing degree days*. Greenhouses would also allow for growth in some regions which would otherwise be unable to grow significant quantities of food. To avoid overly optimistic outcomes, we have assumed that newly constructed greenhouses would be built only on the available outdoor growing cropland area, and subtracted this fraction of area from available outdoor cropland.

Greenhouses are estimated to produce a 50% yield increase versus outdoor growing post-disaster in a 150 Tg scenario, based on a *growing degree day* evaluation of maize and potatoes (see Supplementary Information, section VIII).

Seaweed


Low-tech marine seaweed farm designs hold potential to be a low-cost, rapidly scaleable, and nutritious food source. Seaweed tolerates low sunlight conditions and cold temperatures⁵⁶, and can quickly scale to high productivity⁴². For these reasons, seaweed is an especially promising food source in an ASRS. We selected *Porphyra amplissima* as a representative example for its cold tolerance and high protein content⁵⁷. Species of *Porphyra* are also widely consumed in Southeastern countries today⁵⁸. However, other varieties would also be suitable, such as those in the *Gracilaria* family. Due to nutritional and palatability concerns, seaweed has been capped at a maximum of 10% of daily calories. The linear optimization model used to estimate seaweed growth and deployment of new farms is listed in Supplementary Information section IX.

Dairy and Meat Output

While crop yields would be severely reduced in an ASRS, efficient allocation of agricultural residues could be used to maintain a significant amount of dairy production. Prioritizing maintenance of dairy is justified by the favorable feed and protein conversion efficiency of dairy as compared to beef⁵⁹, with around 400 kcal and 21 g of protein per kg of inedible feed for milk and 22 kcal and 2 g of protein per kg of inedible feed for beef (see Supplementary Information, section VI for a detailed breakdown).

For technical feasibility calculations, it was estimated that livestock was reduced to levels that could be maintained by a combination of grasslands, agricultural residues, fodder crops and excess human edible food for the 150 Tg scenario. An iterative market equilibrium was used to estimate that feed would be maintained such that approximately 50% of current-day meat production on a caloric equivalent (mostly chicken and pork) could be maintained from Year 2 given the central assumptions of the resilient food model (see Supplementary Information, Section VI). This system is summarized below, in **Figure 5**.

In both resilient and non-resilient 150 Tg scenarios, the meat obtained by culling livestock populations was allocated uniformly over the first 60 months of the scenarios. Livestock slaughter capacity scaling up might be needed but was not considered.







Figure 5: a-c Examples of animal feed system modeling, before and after the disaster. We assume feed is applied to the most efficient use in terms of calories/protein, up to the limit set by output in the present day. This allows for significant output of animal products despite a sharp decline in feed output **c** Year 3 with surplus human edible foods, based upon the resilient food scenario under the median scenario. All values are in dry caloric tonnes (normalized to 4,000,000 kcal/metric tonne) to allow comparison between different foods.

Marine Fish

The catching of marine fish could continue at a diminished rate during the 150 Tg scenario⁴. This accounts for roughly half of current aquatic caloric production. Meanwhile, aquaculture systems typically use human edible food and were assumed to cease post-disaster. Marine fish was treated as an exogenous variable.

Industrial Foods

The ramp-up of industrial resilient food production in the 150 Tg scenario was modeled as a combination of the two options with highest technology readiness at the current time of writing. The first one is sugars from lignocellulosic biomass assuming rapid repurposing of pulp and paper mills to sugar biorefineries¹⁸. The second one is single cell protein (SCP) production from natural gas based on fast construction of large-scale fermentation facilities¹⁵. The rate of growth of industrial foods in the current model was hard-coded from the results of the cited papers, and is thus treated as an exogenous variable. First a

large wave of paper factory repurposing is assumed (~two thirds of the current global pulp and paper capacity), from then onwards the available industrial resources are assumed to be invested in methane SCP production. SCP can serve as a useful food product due to its high-quality protein content and micronutrient profile, despite the higher resource intensity and unit costs compared to lignocellulosic sugar. Fast repurposing for sugar production can help bridge the sudden gap in food production, but after the most promising facilities have been repurposed, a switch to more nutritionally rich products appears reasonable.

In the 150 Tg scenario, multiple industrial responses would likely develop in parallel at different paces and with varying degrees of success, as happened during the COVID-19 pandemic^{60,61}, potentially including others not considered in this model such as foods from CO₂ (other SCPs¹⁶, carbohydrates⁶², electrosynthesized foods¹⁷, etc.) or synthetic fats¹⁹. However, this is arguably a realistic representation of the potential of leveraging industrial resources for increasing resilient food production from inedible raw materials.

The ramp-up values from the literature used in this work were estimated based on the capital expenditure budget of chemical and related industrial sectors. There is some chance that qualified labor, material resources, equipment construction capacity and supply chains of other industries could also be quickly and efficiently diverted to food production, resulting in values ~10 times higher than the ones used in this work for industrial foods, but this is speculative and might require preparedness plans to be in place prior to an ASRS. Further research should be conducted to properly characterize the potential of agriculture-independent industrial food production for GCFS response.

5 Data Availability

Supplementary Information includes more detail on methodology, while the supplementary spreadsheet at https://github.com/allfed/allfed-integrated-model/blob/main/Supplemental_Data.xlsx contains quantitative assumptions used by the model.

6 Code Availability

The Python code used and associated Jupyter notebook is available at <https://github.com/allfed/allfed-integrated-model>.

References

1. Denkenberger, D. C. & Pearce, J. M. Feeding everyone: Solving the food crisis in event of global catastrophes that kill crops or obscure the sun. *Futures* **72**, 57–68 (2015).

2. Bostrom, N. & Cirkovic, M. M. *Global Catastrophic Risks*. (OUP Oxford, 2008).
3. Baum, S., De Neufville, R. & Barrett, A. A model for the probability of nuclear war. *Baum Seth Neufville Robert Barrett Anthony Model Probab. Nucl. War* March 8 2018 *Glob. Catastrophic Risk Inst. Work. Pap. 18-1* **18**, (2018).
4. Denkenberger, D. & Pearce, J. *Feeding everyone no matter what: Managing food security after global catastrophe*. (Academic Press, 2014).
5. Cotton-Barratt, O., Daniel, M. & Sandberg, A. Defence in Depth Against Human Extinction: Prevention, Response, Resilience, and Why They All Matter. *Glob. Policy* **11**, 271–282 (2020).
6. Handmer, J. et al. *A Framework for Global Science in support of Risk Informed Sustainable Development and Planetary Health*.
<https://www.undrr.org/publication/framework-global-science-support-risk-informed-sustainable-development-and-planetary> (2021).
7. UN Secretary-General et al. *OUR COMMON AGENDA: Report of the Secretary-General*.
<https://www.un.org/en/content/common-agenda-report/> (2021).
8. Xia, L. et al. *Global Famine after Nuclear War*. *Submitted to Nature Food* (2021)
doi:10.21203/rs.3.rs-830419/v1.
9. Pham, A. et al. Nutrition in Abrupt Sunlight Reduction Scenarios: Envisioning Feasible Balanced Diets on Resilient Foods. *Nutrients* **14**, 492 (2022).
10. FAO, I. *The State of Food Security and Nutrition in the World 2020: Transforming food systems for affordable healthy diets*. (FAO, IFAD, UNICEF, WFP and WHO, 2020).
doi:10.4060/ca9692en.
11. Robock, A., Oman, L. & Stenchikov, G. Nuclear winter revisited with a modern climate model and current nuclear arsenals: Still catastrophic consequences. *J. Geophys. Res.* **112**, (2007).
12. Verma, M. van den B., Vreede, L. de, Achterbosch, T. & Rutten, M. M. Consumers discard

- a lot more food than widely believed: Estimates of global food waste using an energy gap approach and affluence elasticity of food waste. *PLOS ONE* **15**, e0228369 (2020).
13. WHO. *The Management of nutrition in major emergencies*. (World Health Organization, 2000).
 14. Alvarado, K. A., Mill, A., Pearce, J. M., Vocaet, A. & Denkenberger, D. Scaling of greenhouse crop production in low sunlight scenarios. *Sci. Total Environ.* **707**, 136012 (2020).
 15. García Martínez, J. B., Pearce, J. M., Throup, J., Cates, J. & Denkenberger, D. C. Methane Single Cell Protein: securing protein supply during global food catastrophes. (2020) doi:10.31219/osf.io/94mkg.
 16. García Martínez, J. B. *et al.* Potential of microbial protein from hydrogen for preventing mass starvation in catastrophic scenarios. *Sustain. Prod. Consum.* **25**, 234–247 (2021).
 17. García Martínez, J. B., Brown, M. M., Christodoulou, X., Alvarado, K. A. & Denkenberger, D. C. Potential of microbial electrosynthesis for contributing to food production using CO₂ during global agriculture-inhibiting disasters. *Clean. Eng. Technol.* **4**, (2021).
 18. Throup, J. *et al.* Rapid repurposing of pulp and paper mills, biorefineries, and breweries for lignocellulosic sugar production in global food catastrophes. *Food Bioprod. Process.* **131**, 22–39 (2022).
 19. García Martínez, J. B., Alvarado, K. A. & Denkenberger, D. C. Synthetic fat from petroleum as a resilient food for global catastrophes: Preliminary techno-economic assessment and technology roadmap. *Chem. Eng. Res. Des.* **177**, 255–272 (2022).
 20. World Food Summit - Final Report - Part 1.
<https://www.fao.org/3/w3548e/w3548e00.htm>.
 21. Allen, R. C. Absolute Poverty: When Necessity Displaces Desire. *Am. Econ. Rev.* **107**, 3690–3721 (2017).
 22. Qinghua Zhao. *PovcalNet*. (World Bank, 2021).

23. Herforth, A. et al. *Cost and affordability of healthy diets across and within countries: Background paper for The State of Food Security and Nutrition in the World 2020*. FAO Agricultural Development Economics Technical Study No. 9. (Food & Agriculture Org., 2020).
24. Godfray, H. C. J. et al. Food Security: The Challenge of Feeding 9 Billion People. *Science* **327**, 812–818 (2010).
25. Lang, T. & Barling, D. Food security and food sustainability: reformulating the debate. *Geogr. J.* **178**, 313–326 (2012).
26. Tigchelaar, M., Battisti, D. S., Naylor, R. L. & Ray, D. K. Future warming increases probability of globally synchronized maize production shocks. *Proc. Natl. Acad. Sci.* **115**, 6644–6649 (2018).
27. Jones, A. W. & Phillips, A. Historic Food Production Shocks: Quantifying the Extremes. *Sustainability* **8**, 427 (2016).
28. Cottrell, R. S. et al. Food production shocks across land and sea. *Nat. Sustain.* **2**, 130–137 (2019).
29. Brönnimann, S. & Krämer, D. Tambora and the "Year Without a Summer" of 1816. A Perspective on Earth and Human Systems Science. *Geogr. Bernensia* **90**, (2016).
30. Zerefos, C. S., Gerogiannis, V. T., Balis, D., Zerefos, S. C. & Kazantzidis, A. Atmospheric effects of volcanic eruptions as seen by famous artists and depicted in their paintings. *Atmospheric Chem. Phys.* **7**, 4027–4042 (2007).
31. Puma, M. J., Bose, S., Chon, S. Y. & Cook, B. I. Assessing the evolving fragility of the global food system. *Environ. Res. Lett.* **10**, 024007 (2015).
32. Helbing, D. Globally networked risks and how to respond. *Nature* **497**, 51–59 (2013).
33. Mouritsen, O. G. *Introduction to Phase Transitions and Critical Phenomena*. By H. Eugene Stanley, Oxford University Press, Oxford, 1987. *Int. J. Quantum Chem.* **35**, 583–584 (1989).

34. Dobson, I., Carreras, B. A., Lynch, V. E. & Newman, D. E. Complex systems analysis of series of blackouts: Cascading failure, critical points, and self-organization. *Chaos Interdiscip. J. Nonlinear Sci.* **17**, 026103 (2007).
35. Buldyrev, S. V., Parshani, R., Paul, G., Stanley, H. E. & Havlin, S. Catastrophic cascade of failures in interdependent networks. *Nature* **464**, 1025–1028 (2010).
36. Wik, M., Pingali, P. & Brocai, S. *Global Agricultural Performance : Past Trends and Future Prospects*. <https://openknowledge.worldbank.org/handle/10986/9122> (2008).
37. Herrendorf, B., Rogerson, R. & Valentinyi, Á. Chapter 6 - Growth and Structural Transformation. in *Handbook of Economic Growth* (eds. Aghion, P. & Durlauf, S. N.) vol. 2 855–941 (Elsevier, 2014).
38. World Development Indicators | DataBank.
<https://databank.worldbank.org/reports.aspx?source=world-development-indicators>.
39. Baum, S. & Barrett, A. *A Model for the Impacts of Nuclear War*.
<https://papers.ssrn.com/abstract=3155983> (2018) doi:10.2139/ssrn.3155983.
40. Tzachor, A., Richards, C. E. & Holt, L. Future foods for risk-resilient diets. *Nat. Food* **2**, 326–329 (2021).
41. Roque, B. M. *et al.* Red seaweed (*Asparagopsis taxiformis*) supplementation reduces enteric methane by over 80 percent in beef steers. *PLOS ONE* **16**, e0247820 (2021).
42. Yarish, C. *et al.* *Seaweed Aquaculture for Food Security, Income Generation and Environmental Health in Tropical Developing Countries*. <https://doi.org/10.1596/24919> (2016).
43. Mill, A. *et al.* Preventing global famine in case of sun-blocking scenarios: Seaweed as an alternative food source. in (2019).
doi:https://www.researchgate.net/publication/337199859_Preventing_global_famine_in_case_of_sun-blocking_scenarios_Seaweed_as_an_alternative_food_source_Key_findings.

44. Denkenberger, D. C. *et al.* Feeding everyone if the sun is obscured and industry is disabled. *Int. J. Disaster Risk Reduct.* **21**, 284–290 (2017).
45. Denkenberger, D., Sandberg, A., Tieman, R. J. & Pearce, J. M. Long-term cost-effectiveness of interventions for loss of electricity/industry compared to artificial general intelligence safety. *Eur. J. Futur. Res.* **9**, 11 (2021).
46. Denkenberger, D. C. & Blair, R. W. Interventions that may prevent or mollify supervolcanic eruptions. *Futures* **102**, 51–62 (2018).
47. Cheng, A. F. *et al.* DART mission determination of momentum transfer: Model of ejecta plume observations. *Icarus* **352**, 113989 (2020).
48. Denkenberger, D. C. & Pearce, J. M. A National Pragmatic Safety Limit for Nuclear Weapon Quantities. *Safety* **4**, 25 (2018).
49. Instituto Geográfico Nacional (IGN) & Civil Protection. *Spanish State Civil Protection Plan for tsunami risk.*
<https://www.ign.es/web/resources/acercaDe/libDigPub/Plan-Estatal-Maremotos.pdf>
 (2021).
50. U.S. FEMA. *Radiological Emergency Preparedness.*
<https://www.fema.gov/emergency-managers/practitioners/hazardous-response-capabilities/radiological> (2019).
51. von Hillebrandt-Andrade, C., Hincapié-Cárdenas, C., Aliaga, B. & Brome, A. UNESCO IOC CARIBE EWS Tsunami Ready Pilot Program. NH41C-1004
<https://ui.adsabs.harvard.edu/abs/2018AGUFMNH41C1004V> (2018).
52. USDA & USDHHS. *2015-2020 Dietary Guidelines for Americans.* 144
<http://health.gov/dietaryguidelines/2015/guidelines/> (2015).
53. Food and Agriculture Organization of the United Nations. FAOSTAT. *FAOSTAT Statistical Database* <http://www.fao.org/faostat/en/#home>.
54. FAO. *Global food losses and food waste – Extent, causes and prevention.* ISBN

- 978-92-5-107205-9 (2011).
55. United States Department of Agriculture. *Production, Supply and Distribution*. (Foreign Agricultural Service, 2021).
56. Bird, N. L., Chen, L. C.-M. & McLachlan, J. Effects of Temperature, Light and Salinity on Growth in Culture of *Chondrus crispus*, *Furcellaria lumbricalis*, *Gracilaria tikvahiae* (Gigartinales, Rhodophyta), and *Fucus serratus* (Fucales, Phaeophyta). *Bot. Mar.* **22**, (1979).
57. Peñalver, R. *et al.* Seaweeds as a Functional Ingredient for a Healthy Diet. *Mar. Drugs* **18**, 301 (2020).
58. Venkatraman, K. L. & Mehta, A. Health Benefits and Pharmacological Effects of *Porphyra* Species. *Plant Foods Hum. Nutr.* **74**, 10–17 (2019).
59. Mottet, A. *et al.* Livestock: On our plates or eating at our table? A new analysis of the feed/food debate. *Glob. Food Secur.* **14**, 1–8 (2017).
60. Okorie, O. *et al.* Manufacturing in the Time of COVID-19: An Assessment of Barriers and Enablers. *IEEE Eng. Manag. Rev.* **48**, 167–175 (2020).
61. Poduval, A. *et al.* Barriers in repurposing an existing manufacturing plant: a total interpretive structural modeling (TISM) approach. *Oper. Manag. Res.* (2021) doi:10.1007/s12063-021-00209-9.
62. García Martínez, J. B., Alvarado, K. A., Christodoulou, X. & Denkenberger, D. C. Chemical synthesis of food from CO₂ for space missions and food resilience. *J. CO₂ Util.* **53**, (2021).

Acknowledgements

This research was supported by the Alliance to Feed the Earth in Disasters (ALLFED).

Thanks to Aron Mill, Joshua Pearce, Kyle Alvarado, Adin Richards, and Gal Hochman for their insights.

Author Information

Affiliations

^a *Alliance to Feed the Earth in Disasters (ALLFED), Fairbanks, AK, USA.*

^b *ETH Zürich, Department of Mathematics, 8092 Zurich, Switzerland*

^c *Northumbria University, Faculty of Engineering and Environment, City Campus, Newcastle-upon-Tyne, NE1 8ST, United Kingdom*

^d *University of Alaska Fairbanks, Mechanical Engineering and Alaska Center for Energy and Power, Fairbanks, AK, 99775 USA*

*Corresponding author: morgan@allfed.info

Ethics Declarations

Competing interests

The authors declare no competing interests.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInformation.pdf](#)