

Detection of alpha and betacoronaviruses in multiple Iberian bat species

Ana Falcón · Sonia Vázquez-Morón · Inmaculada Casas · Carolina Aznar ·
Guillermo Ruiz · Francisco Pozo · Pilar Perez-Breña · Javier Juste ·
Carlos Ibáñez · Inazio Garin · Joxerra Aihartza · Juan E. Echevarría

Received: 2 February 2011 / Accepted: 20 June 2011 / Published online: 16 July 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract Bat coronaviruses (CoV) are putative precursors of the severe acute respiratory syndrome (SARS) CoV and other CoV that crossed the species barrier from zoonotic reservoirs into the human population. To determine the presence and distribution of CoV in Iberian bats, 576 individuals of 26 different bat species were captured in 13 locations in Spain. We report for the first time the presence of 14 coronaviruses in 9 Iberian bat species. Phylogenetic analysis of a conserved CoV genome region (RdRp gene) shows a wide diversity and distribution of alpha and betacoronavirus in Spain. Interestingly, although some of these viruses are related to other European BatCoV, or to Asian CoV, some of the viruses found in Spain cluster in new groups of α and β CoV.

GenBank accession numbers: Partial sequences of RdRp gene of bat coronaviruses obtained in this study were deposited in GenBank under the accession numbers HQ184049 to HQ184062.

A. Falcón · S. Vázquez-Morón · I. Casas · C. Aznar · G. Ruiz ·
F. Pozo · P. Perez-Breña · J. E. Echevarría
Centro Nacional de Microbiología, Instituto de Salud Carlos III,
Madrid, Spain

A. Falcón (✉)
Centro Nacional de Biotecnología, CSIC, Darwin 3,
Campus Cantoblanco, 28049 Madrid, Spain
e-mail: afalcon@cnb.csic.es

J. Juste · C. Ibáñez
Estación Biológica de Doñana, CSIC, Seville, Andalusia, Spain

I. Garin · J. Aihartza
Department of Zoology and Animal Cell Biology,
University of the Basque Country (UPV/EHU),
Leioa, The Basque Country, Spain

The emergence of infectious diseases is a major threat to global public health in this century (WHO, World Health Report <http://www.who.int/whr/previous/en/index.html>) and many of these new infectious human diseases are caused by viruses emerging from wildlife. In the last 50 years, more than 30 new infectious human diseases have been identified (WHO, World Health Report <http://www.who.int/whr/previous/en/index.html>) including the Severe Acute Respiratory Syndrome (SARS). The aetiological agent of this disease was identified as a previously unknown coronavirus (SARS-CoV) [1] and BatCoV are putative precursors of SARS-CoV [2]. The outbreak of SARS-CoV and subsequent identification of two additional human coronaviruses (HCoV-NL63 [3] and HCoV-HKU1 [4]) has drawn human and animal health attention to *Coronavirinae* subfamily, that includes three genera, *Alphacoronavirus* (α CoV), *Betacoronavirus* (β CoV) and *Gammacoronavirus* (γ CoV), replacing the classical groups 1, 2 and 3 [5] (<http://talk.ictvonline.org/media/g/vertebrate-2008/default.aspx>).

The relevance and possible re-emergence of the pandemic SARS-CoV and other emerging viruses of zoonotic origin have activated surveillance systems of hazard agents in wild animals, including bats. As a result of these studies, bats have been described as putative reservoirs for some emerging viruses affecting humans [6]. BatCoV are putative precursors of CoV affecting humans and mammals [7, 8], including SARS-CoV [2] and other CoV that crossed the species barrier from zoonotic reservoirs into the human population [9]. In fact, association of some of these CoV to certain bat species has been suggested [10, 11], reinforcing the notion that there may be a relationship between some BatCoV and their hosts. Nowadays the presence of CoV has been shown in bats in China [12, 13], North and South America [14–17], Africa [18] and a number of regions in

Europe [11, 19–21] but not the Iberian Peninsula (Spain and Portugal), which is a bridge for European and African bat populations [22]. Thus, surveillance of wildlife reservoirs of putative zoonotic CoV is necessary, not only to unveil the ecology of these viruses, but also to permit early detection of viruses that might pose a threat to human health.

To determine the presence and distribution of putative zoonotic CoV in Iberian bats, 576 individuals from 26 bat species were captured and sampled in 13 different locations throughout Spain during 2004–2007 (Fig. 1). These samples were collected in the context of bat rhabdoviruses and lyssaviruses Surveillance Program in Spain. Most of the sampled bat species are also distributed across Europe, but *Eptesicus isabellinus* is a meridional serotine bat restricted to North Africa and the Iberian Peninsula [23], and *Myotis escalerai* is endemic in the Iberian Peninsula. Bats were caught with mist nets mainly as they left diurnal roost and by hand with polyethylene butterfly nets within roosts. Oro-pharyngeal swabs (n=390) between 2004 and 2007 as well as faecal samples from individual bats (n=216) in 2007 were taken before bats were released (Table 1).

Oro-pharyngeal swabs collected between 2004 and 2006 were preserved in 1 ml of lysis buffer (4 M GuSCN (Sigma), OS% N-lauroyl Sarcosine (Sigma), 1 mM dithiothreitol (DTT, Sigma), 25 mM Sodium Citrate and 20 µg/tube Glycogen (Boehringer Mannheim). Oro-pharyngeal swabs and faeces collected in 2007 were preserved in both 1 ml of lysis buffer and 1 ml of viral transport medium (VTM) (Eagle's minimal essential medium (EMEM) supplemented with 10 UI/ml of penicillin, 10 µg/ml of streptomycin, 160 µg/ml of gentamicin, 50 UI/ml of mycostatin and 1% of



Fig. 1 Geographical location of bat capture sites in Spain. 1: A Coruña, 2: Lugo, 3: La Rioja, 4: Gerona, 5: Castellón, 6: Valencia, 7: Alicante, 8: Málaga, 9: Cádiz, 10: Sevilla, 11: Huelva, 12: Cáceres, 13: Menorca. The stars show the locations where positive samples were found

bovine serum albumin). All samples were frozen at -80°C before sending them to the Rabies Reference Laboratory at the Centro Nacional de Microbiología, ISCIII in Madrid. Faecal samples were clarified by centrifugation. Total nucleic acid was extracted from a 200 µl aliquot of each specimen for PCR assays and the rest were stored to -80°C in two different aliquots. Final pellets were always resuspended to 55 µl of water.

A pan-coronavirus nested PCR was designed in the RdRp gene. A total of 5 µl of extracted RNA was added to 45 µl of reaction mixture of OneStep RT-PCR kit (QIAGEN, Valencia, CA, USA) containing 200 µM dNTPs and 60 pmol of generic CoV-specific degenerated primers (forward 5'-CARATGAATYTTAARTAYGC-3' and reverse 5'-TGYTGWGARCAAAAAYTCRTG-3') and following manufacturer indications. Amplifications were carried out into thin-walled reaction tubes (Sorenson, BioScience, UT) in a PTC-200 (Peltier Thermal Cycler, MJ Research, Watertown, MA). Nested PCR amplifications were performed using 2 µl of first amplification product and 48 µl of reaction mixture containing 60 mM Tris-HCl (pH 8.5), 15 mM $(\text{NH}_4)_2\text{SO}_4$, 200 µM dNTPs (Amersham Pharmacia Biotech, Piscataway, NJ), 3 mM MgCl_2 , 35 pmol of generic CoV-specific degenerated primers (forward 5'-ATGGGWTGGGAYTAYCCIAARTG-3' and reverse 5'-ACRTRRTTYTGRWARTA-3') and 1.25 U AmpliTaq DNA Polymerase (Perkin-Elmer Cetus, Norwalk). Amplification product size of 512 nt was visualized by agarose gel electrophoresis and sequenced directly in both directions using an automated ABI PRISM 377 model sequencer. For phylogeny reconstruction, consensus sequences were aligned together with others obtained from public genomic databases using the program CLUSTAL X (version 1.83) (Table 2). A Bayesian phylogenetic inference was obtained using Mr Bayes version 3.1 [24] with random starting trees without constraints. For the analyses GTR substitution model, gamma estimation and two simultaneous runs of 10^7 generations were done, each with four Markov chains, and the trees were sampled every 100 generations. Amino acid identity was calculated with MEGA 4 using the pairwise deletion option. The alignment comprised the same 396 bp of the RdRp gene used for the phylogenetic reconstruction.

A total of 26 out of the 30 known bat species known for Iberian Peninsula were screened for CoV and 14 samples taken from 9 bat species, all included in the family Vespertilionidae, were positive for CoV RNA (Tables 1 and 2). Twelve of them were found within faecal samples (5.5%) of 7 different bat species in 6 locations and two were obtained in oral samples (0.5%) of 2 other different species in the same location (Table 1). In view of these results, it is not surprising that the presence of CoV RNA is significantly more frequent in faeces than in oral cavity

Table 1 Results of detection of CoV RNA in faecal or oral samples of bats collected in Spain

Bat Species	Faecal samples positive/no. tested	Oral samples positive/no. tested	Location	Genus
<i>Barbastella barbastellus</i>	0/4	0/2	3, 4	
<i>Eptesicus isabellinus</i>	1/8	NA	10 ^a	β
<i>Eptesicus serotinus</i>	0/7	NA	1, 12	
<i>Hypsugo savii</i>	2/26	0/10	2, 4 ^a , 12 ^{a,b}	α, β
<i>Miniopterus schreibersii</i>	0/2	1/71	2, 5, 6 ^a , 7, 8, 9, 12, 13	α
<i>Myotis alcathoe</i>	0/1	NA	3	
<i>Myotis bechsteinii</i>	0/2	0/3	4, 8	
<i>Myotis blythii</i>	NA	1/11	6 ^a , 7, 12	α
<i>Myotis capaccinii</i>	NA	0/14	5, 6, 13	
<i>Myotis daubentonii</i>	1/39	0/52	2, 3, 8 ^a , 11, 12	α
<i>Myotis emarginatus</i>	NA	0/2	8, 12	
<i>Myotis escalerae</i>	NA	0/15	11, 7	
<i>Myotis myotis</i>	1/1	0/17	6, 7, 8 ^a , 12	α
<i>Myotis mystacinus</i>	0/5	NA	2, 3	
<i>Myotis nattereri</i>	0/3	0/3	4	
<i>Nyctalus lasiopterus</i>	5/37	0/137	3, 8 ^{a,b} , 9, 10, 11	α
<i>Nyctalus leisleri</i>	0/23	0/11	2, 3, 4, 8	
<i>Pipistrellus kuhlii</i>	1/4	0/6	8, 12 ^a	α
<i>Pipistrellus pipistrellus</i>	0/3	0/1	3, 12	
<i>Pipistrellus pygmaeus</i>	NA	0/1	12	
<i>Pipistrellus</i> sp.	1/29	0/5	1, 2, 4, 8, 12 ^{a,b}	α
<i>Plecotus auritus</i>	0/7	NA	1, 3, 4	
<i>Plecotus austriacus</i>	0/7	0/10	4, 11, 12	
<i>Rhinolophus euryale</i>	NA	0/13	6, 7, 12	
<i>Rhinolophus ferrumequinum</i>	0/3	0/5	4, 8, 12	
<i>Rhinolophus hipposideros</i>	0/4	NA	4, 12	
<i>Rhinolophus mehelyi</i>	NA	0/1	12	
Total	12/216	2/390	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13	α, β

NA no samples available

^a Locations where positives samples were found

^b These samples were collected in different localities than other positive samples with the same number location

($p < 0.0001$, FISHER EXACT TEST). It is of interest that none of the viruses has been found in oropharyngeal and faecal samples of the same individual, when both samples were available. This fact may indicate either that the infection was at different stage in the different individuals at the time of sampling or that replication of virus may take place independently in the intestinal and respiratory tracts [25]. Most of the CoV RNA sequences found in faecal samples (83%) correspond to α CoV, the remaining two belonging to β CoV. All CoV RNAs from oral samples (100%) were found to contain viral RNA sequences corresponding to α CoV. In agreement with all previous studies [7], none of the coronavirus detected in Spanish bats belong to group γ .

The phylogenetic analysis of Spanish BatCoV was performed using 396nt out of the 512 nt RT-PCR amplified

fragments. 116nt fragment information was lost to allowed us include more sequences from other European countries and other continents deposited in GenBank to perform a meaningful analysis. This small part of the RdRp gene has been previously used, and sufficiently represents the full gene information, for phylogenetic analysis of BatCoV [11, 14, 20]

The phylogenetic reconstruction showed 6 different lineages of Spanish BatCoV (Fig. 2). BatCoV A and B were closely related to other α BatCoV found in China [12], although they appeared to display certain genetic differentiation (Fig. 2). *Myotis daubentonii*-associated CoV H, and *Pipistrellus*-associated CoV K, clustered respectively with lineages 4 and 3 of α CoV previously described in Germany and are hosted by the same bat species or genera [11] (Fig. 2). BatCoV L was closely

Table 2 Data of interest related to the 91 coronavirus sequences used for the generation of the phylogenetic tree

Access no	Host species	Country	Genus	Cluster
DQ249221	Bat	China	β	HKU5
DQ249219	Bat	China	β	HKU5
DQ249218	<i>Pipistrellus</i> sp.	China	β	HKU5
DQ648809	Bat	China	β	
DQ648807	Bat	China	β	
DQ249217	<i>Pipistrellus</i> sp.	China	β	HKU5
DQ648819	Bat	China	β	
DQ249215	<i>Tylonycteris</i>	China	β	HKU4
DQ249214	<i>Tylonycteris</i>	China	β	HKU4
DQ074652	<i>Tylonycteris</i>	China	β	HKU4
DQ249216	<i>Tylonycteris</i>	China	β	HKU4
DQ648803	Bat	China	β	
HQ184059	<i>Hypsugo savii</i>	Spain	β	
HQ184062	<i>Eptesicus isabellinus</i>	Spain	β	
GQ404795	<i>Rhinolophus hipposideros</i>	Slovenia	β	
GQ404796	<i>Rhinolophus hipposideros</i>	Slovenia	β	
GQ404797	<i>Rhinolophus hipposideros</i>	Slovenia	β	
DQ022305	<i>Rhinolophus sinicus</i>	China	β	
NC_009696	<i>Rhinolophus macrotis</i>		β	
NC_004718	Human		β	
NC_009021	<i>Rousettus leschenaulti</i>	China	β	HKU9
NC_006577	Human		β	
NC_006852	Mouse		β	
NC_007732	Pig		β	
NC_005147	Human		β	
EF544563	<i>Myotis occultus</i>	USA	α	
EF544565	<i>Myotis occultus</i>	USA	α	
HQ184049	<i>Miniopterus schreibersii</i>	Spain	α	
HQ184050	<i>Myotis blythii</i>	Spain	α	
DQ648838	Bat	China	α	
DQ648855	<i>Rhinolophus ferrumequinum</i>	China	α	
DQ648854	<i>Rhinolophus ferrumequinum</i>	China	α	
NC_003436	Pig		α	
EU375862	<i>Myotis dasycneme</i>	Germany	α	
EU375859	<i>Myotis dasycneme</i>	Germany	α	
EU375858	<i>Myotis dasycneme</i>	Germany	α	
EU375855	<i>Myotis dasycneme</i>	Germany	α	
EU375863	<i>Myotis dasycneme</i>	Germany	α	
EU375861	<i>Myotis dasycneme</i>	Germany	α	
EU375856	<i>Myotis dasycneme</i>	Germany	α	
EU375854	<i>Myotis dasycneme</i>	Germany	α	
EU375857	<i>Myotis dasycneme</i>	Germany	α	
EU375865	<i>Myotis bechsteinii</i>	Germany	α	
EU375853	<i>Myotis bechsteinii</i>	Germany	α	
EU375860	<i>Myotis bechsteinii</i>	Germany	α	
EU375869	<i>Pipistrellus nathusii</i>	Germany	α	
EU375864	<i>Pipistrellus nathusii</i>	Germany	α	
EU375870	<i>Pipistrellus pygmaeus</i>	Germany	α	

Table 2 continued

Access no	Host species	Country	Genus	Cluster
EU375868	<i>Pipistrellus pygmaeus</i>	Germany	α	
EU375867	<i>Pipistrellus pygmaeus</i>	Germany	α	
HQ184060	<i>Pipistrellus</i> sp.	Spain	α	
DQ648822	Bat	China	α	
DQ648821	Bat	China	α	
DQ648824	Bat	China	α	
DQ648823	Bat	China	α	
EU375875	<i>Myotis daubentonii</i>	Germany	α	
EU375873	<i>Myotis daubentonii</i>	Germany	α	
EU375874	<i>Myotis daubentonii</i>	Germany	α	
EU375872	<i>Myotis daubentonii</i>	Germany	α	
EU375866	<i>Myotis daubentonii</i>	Germany	α	
HQ184056	<i>Myotis daubentonii</i>	Spain	α	
EU375871	<i>Myotis daubentonii</i>	Germany	α	
DQ648833	<i>Myotis ricketti</i>	China	α	
DQ249224	<i>Myotis ricketti</i>	China	α	HKU6
DQ648837	<i>Myotis ricketti</i>	China	α	
DQ249235	<i>Rhinolophus sinicus</i>	China	α	HKU2
DQ249213	<i>Rhinolophus sinicus</i>	China	α	HKU2
DQ648840	Bat	China	α	
DQ249228	<i>Miniopterus</i> sp.	China	α	HKU8
EU834954	<i>Miniopterus australis</i>	Australia	α	
EU834952	<i>Miniopterus australis</i>	Australia	α	
EU834955	<i>Miniopterus schreibersii</i>	Australia	α	
EU834953	<i>Rhinolophus megaphyllus</i>	Australia	α	
DQ648835	<i>Miniopterus schreibersii</i>	China	α	
DQ648796	Bat	China	α	
DQ648797	Bat	China	α	
DQ249226	<i>Miniopterus magnater</i>	China	α	HKU7
HQ184061	<i>Hypsugo savii</i>	Spain	α	
HQ184051	<i>Nyctalus lasiopterus</i>	Spain	α	
HQ184054	<i>Nyctalus lasiopterus</i>	Spain	α	
HQ184053	<i>Nyctalus lasiopterus</i>	Spain	α	
HQ184052	<i>Nyctalus lasiopterus</i>	Spain	α	
HQ184055	<i>Nyctalus lasiopterus</i>	Spain	α	
HQ184057	<i>Myotis myotis</i>	Spain	α	
HQ184058	<i>Pipistrellus kuhlii</i>	Spain	α	
EU834951	<i>Myotis macropus</i>	Australia	α	
NC_002645	Human		α	
NC_005831	Human		α	
AY994055	Cat		α	
NC_002306	Pig		α	
NC_001451	Chicken		γ	

related to cluster HKU7 described in China [10]. BatCoV C, D, E, F, G and I belonged to the same lineage and showed an intriguing new independent cluster (significant posterior probability = 0.95) including BatCoV I' (Fig. 2). In addition, BatCoV J and M were genetically related to β CoV, although they did not really associate with any of their previously described lineages. Sequences corresponding to two different groups of CoV were found in the same bat species (*Hypsugo savii*) as it was also found for other bat species elsewhere [12]. Spanish BatCoV described here were also classified calculating amino acid distances of these viruses from phylogenetic neighbours and related reference species [21]. Amino acid distance criteria recently described for separating RdRp grouping units (RGU) were adopted [21]. Interestingly, the amino acid distance criteria (>4.8% for alphacoronavirus and >6.3% for betacoronavirus) reinforced the presence of the new group of alphacoronaviruses mentioned above including Spanish BatCoV C, D, E, F, G and I; and additionally showed four new tentative groups (Fig. 2). BatCoV A and B were included in a new alphacoronavirus group and BatCoV I' constituted another independent alphacoronavirus group. BatCoV J, and M represented two new betacoronavirus groups and BatCoV H, K and L remained as part of several established species (Fig. 2).

It has been previously suggested that some CoV associate to certain bat species [10, 11]. However, we found that different bat species from the same colony or location harbour CoV of the same genetic lineage (BatCoV A and B; G and I), indicating a greater diversity and higher complexity than previously described for the ecology of BatCoV. Similar exceptions were found in China and South America [12, 16, 26] and could also be observed with Australian BatCoV (Fig. 2 and Table 2).

In conclusion, previous studies showed the presence of BatCoV in Europe. However, to our knowledge, this is the first report describing the presence of CoV RNA in Iberian bat species. Phylogenetic data indicate high diversity, wide distribution and complex ecology of CoV in bats captured in diverse Spanish locations. The sequences reported herein provide new insights into the diversity of coronaviruses and describe new phylogenetic lineages that appear to diverge from all genotypes previously detected in other European locations. Future studies should clarify whether such apparently high diversity reflects the bio-geographical peculiarities of the Iberian Peninsula or not. This study contributes with a new dataset to the global surveillance of emerging BatCoV with pathogenic potential in humans. Our data reinforce the notion that the ecology and transmission of CoV in bat reservoirs is far from being completely understood and that more studies will be necessary to evaluate the magnitude of the potential threat that these viruses pose to human health.

Acknowledgments We gratefully thank S. Perlman and P. Woo for providing MHV RNA and HCoV-HKU1 RT-PCR products respectively, as controls for pancoronavirus PCR development. We also thank L. Enjuanes for scientific discussions and P. Gastaminza for critical comments on the manuscript. We are grateful to David García, Ana Popa, Grupo Drosera, Museu de Granollers and everyone who participated in the collection of field samples and the Genomic Unit of the National Center of Microbiology for analyzing the genomic sequences. This work was supported by a Spanish Ministry of Science and Innovation grant (Projects SAF2006-12784-C02-01 and SAF2006-12784-C02-02) and a grant from the Basque Government (ref. IT301-10).

Conflict of interest The authors declare no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Drosten C, Günther S, Preiser W, van der Werf S et al (2003) Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *N Engl J Med* 348:1967–1976
2. Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, Wang H et al (2005) Bats are natural reservoirs of SARS-like coronaviruses. *Science* 310:676–679
3. van der Hoek L, Pyrc K, Jebbink MF, Vermeulen-Oost W, Berkhout RJ, Wolthers KC, Wertheim-van Dillen PM, Kaandorp J, Spaargaren J, Berkhout B (2004) Identification of a new human coronavirus. *Nat Med* 10:368–373
4. Woo PC, Lau SK, Chu CM, Chan KH, Tsoi HW, Huang Y, Wong BH, Poon RW, Cai JJ, Luk WK, Poon LL, Wong SS, Guan Y, Peiris JS, Yuen KY (2005) Characterization and complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia. *J Virol* 79:884–895
5. de Groot RJ, Ziebur J, Poon LL, Woo PC, Talbot P, Rottier PJM, Holmes KV, Baric R, Perlman S, Enjuanes L, Gorbalenya AE (2010) Taxonomic structure of the Coronaviridae. In: Fauquet CM, Mayo MA, Maniloff J, Desselberg U, King A (eds) *Virus taxonomy*. International committee on taxonomy of viruses. Academic Press, London
6. Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T (2006) Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev* 19:531–545
7. Vijaykrishna D, Smith GJ, Zhang JX, Peiris JS, Chen H, Guan Y (2007) Evolutionary insights into the ecology of coronaviruses. *J Virol* 81:4012–4020
8. Woo PCY, Lau SKP, Huang Y, Yuen KY (2009) Coronavirus diversity, phylogeny and interspecies jumping. *Exp Biol Med* 234:1117–1127
9. Pfeifferle S, Oppong S, Drexler JF, Gloza-Rausch F, Ipsen A, Seebens A, Müller MA, Annan A, Vallo P, Adu-Sarkodie Y, Kruppa TF, Drosten C (2009) Distant relatives of severe acute respiratory coronavirus and close relatives of human coronavirus 229E in bats, Ghana. *Emerg Infect Dis* 15:1377–1384
10. Woo PC, Lau SK, Li KS, Poon RW, Wong BH, Tosí HW, Yip BC, Huang Y, Chan KH, Yuen KY (2006) Molecular diversity of coronaviruses in bats. *Virology* 20:180–187
11. Gloza-Rausch F, Ipsen A, Seebens A, Götsche M, Panning M, Felix Drexler J, Petersen N, Annan A, Grywna K, Müller M, Pfeifferle S, Drosten C (2008) Detection and prevalence patterns

- of group I coronaviruses in bats, northern Germany. *Emerg Infect Dis* 14:626–631
12. Tang XC, Zhang JX, Zhang SY, Wang P, Fan XH, Li LF, Li G, Dong BQ, Liu W, Cheung CL, Xu KM, Song WJ, Vijaykrishna D, Poon LLM, Peiris JSM, Smith GJD, Chen H, Guan Y (2006) Prevalence and genetic diversity of coronavirus in bats from China. *J Virol* 80:7481–7490
 13. Chu DK, Poon LL, Chan KH, Chen H, Guan Y, Yuen KY, Peiris JS (2006) Coronaviruses in bent-winged bats (*Miniopterus* spp.). *J Gen Virol* 87:2461–2466
 14. Dominguez SR, O’Shea TJ, Oko LM, Holmes KV (2007) Detection of group I coronaviruses in bats in North America. *Emerg Infect Dis* 13:1295–1300
 15. Misra V, Dumonceaux T, Dubois J, Willis C, Nadin-Davis S, Severini A, Wandeler A, Lindsay R, Artsob H (2009) Detection of polyoma and corona viruses in bats of Canada. *J Gen Virol* 90:2015–2022
 16. Carrington CV, Foster JE, Zhu HC, Zhang JX, Smith GJ, Thompson N, Auguste AJ, Ramkissoon V, Adesiyun AA, Guan Y (2008) Detection and phylogenetic analysis of group I coronaviruses in South American bats. *Emerg Infect Dis* 14:1890–1893
 17. Brandao PE, Scheffer K, Villarreal LY, Achkar S, Oliveira Rde N, Fahl Wde O, Castilho JG, Kotait I, Richtzenhain LJ (2008) A coronavirus detected in the vampire bat *Desmodus rotundus*. *Braz J Infect Dis* 12:466–468
 18. Tong S, Conrardy C, Ruone S, Kuzmin IV, Guo X, Tao Y, Niezgoda M, Haynes L, Agwanda B, Breiman RF, Anderson LJ, Rupprecht CE (2009) Detection of novel SARS-like and other coronaviruses in bats from Kenya. *Emerg Infect Dis* 15:482–485
 19. Balboni A, Palladini A, Bogliani G, Battilani M (2010) Detection of a virus related to betacoronaviruses in Italian greater horseshoe bats. *Epidemiol Infect* 18:1–4
 20. Rihtaric D, Hostnik P, Steyer A, Grom J, Toplak I (2010) Identification of SARS-like coronaviruses in horseshoe bats (*Rhinolophus hipposideros*) in Slovenia. *Arch Virol* 155:507–514
 21. Drexler JF, Gloza-Rausch F, Glende J, Corman VM, Muth D, Goettsche M, Seebens A, Niedrig M, Pfefferle S, Yordanov S, Zhelyazkov L, Hermanns U, Vallo P, Lukashev A, Müller MA, Deng H, Herrler G, Drosten C (2010) Genomic characterization of severe acute respiratory syndrome-related coronavirus in European bats and classification of coronaviruses based on partial RNA-dependent RNA polymerase gene sequences. *J Virol* 84:11336–11349
 22. García-Mudarra JL, Ibáñez C, Juste J (2009) The Straits of Gibraltar: barrier or bridge to Ibero-Moroccan bat diversity? *Biol J Linn Soc* 96:434–450
 23. Ibáñez C, García-Mudarra JL, Ruedi M, Stadelmann B, Juste J (2006) The Iberian contribution to cryptic diversity in European bats. *Acta Chiropter* 8:277–297
 24. Huelsenbeck JP, Ronquist F (2001) MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755
 25. Liu W, Tang F, Fontanet A, Zhan L, Zhao QM, Zhang PH, Wu XM, Zuo SQ, Baril L, Vabret A, Xin ZT, Shao YM, Yang H, Cao WC (2004) Long-term SARS coronavirus excretion from patient cohort, China. *Emerg Infect Dis* 10:1841–1843
 26. Wang LF, Shi Z, Zhang S, Field H, Daszak P, Eaton BT (2006) Review of bats and SARS. *Emerg Infect Dis* 12:1834–1840