



LUND UNIVERSITY

Land-atmosphere exchange of methane from soil thawing to soil freezing in a high-Arctic wet tundra ecosystem

Tagesson, Torbern; Mölder, Meelis; Mastepanov, Mikhail; Sigsgaard, Charlotte; Tamstorf, Mikkel P.; Lund, Magnus; Falk, Julie Maria; Lindroth, Anders; Christensen, Torben; Ström, Lena

Published in:
Global Change Biology

DOI:
[10.1111/j.1365-2486.2012.02647.x](https://doi.org/10.1111/j.1365-2486.2012.02647.x)

2012

Document Version:
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):

Tagesson, T., Mölder, M., Mastepanov, M., Sigsgaard, C., Tamstorf, M. P., Lund, M., Falk, J. M., Lindroth, A., Christensen, T., & Ström, L. (2012). Land-atmosphere exchange of methane from soil thawing to soil freezing in a high-Arctic wet tundra ecosystem. *Global Change Biology*, 18(6), 1928-1940. <https://doi.org/10.1111/j.1365-2486.2012.02647.x>

Total number of authors:
10

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Land-atmosphere exchange of methane from soil thawing to soil freezing in a high-Arctic wet tundra ecosystem

TORBERN TAGESSON*, MEELIS MÖLDER*, MIKHAIL MASTEPANOV*, CHARLOTTE SIGSGAARD†, MIKKEL P. TAMSTORF‡, MAGNUS LUND‡, JULIE M. FALK*, ANDERS LINDROTH*, TORBEN R. CHRISTENSEN* and LENA STRÖM*

*Department of Physical Geography and Ecosystems Analysis, Lund University, Sölvegatan 12, 223 62, Lund, Sweden, †Institute of Geography and Geology, University of Copenhagen, Øster Voldgade 10, DK-1350, Copenhagen, Denmark, ‡Department for Bioscience - Arctic Environment, Århus University, Frederiksborgvej 399, Roskilde, DK-4000, Denmark

Abstract

The land-atmosphere exchange of methane (CH₄) and carbon dioxide (CO₂) in a high-Arctic wet tundra ecosystem (Rylekærene) in Zackenberg, north-eastern Greenland, was studied over the full growing season and until early winter in 2008 and from before snow melt until early winter in 2009. The eddy covariance technique was used to estimate CO₂ fluxes and a combination of the gradient and eddy covariance methods was used to estimate CH₄ fluxes. Small CH₄ bursts were observed during spring thawing 2009, but these existed during short periods and would not have any significant effect on the annual budget. Growing season CH₄ fluxes were well correlated with soil temperature, gross primary production, and active layer thickness. The CH₄ fluxes remained low during the entire autumn, and until early winter. No increase in CH₄ fluxes were seen as the soil started to freeze. However, in autumn 2008 there were two CH₄ burst events that were highly correlated with atmospheric turbulence. They were likely associated with the release of stored CH₄ from soil and vegetation cavities. Over the measurement period, 7.6 and 6.5 g C m⁻² was emitted as CH₄ in 2008 and in 2009, respectively. Rylekærene acted as a C source during the warmer and wetter measurement period 2008, whereas it was a C sink for the colder and drier period of 2009. Wet tundra ecosystems, such as Rylekærene may thus play a more significant role for the climate in the future, as temperature and precipitation are predicted to increase in the high-Arctic.

Keywords: Micrometeorology, eddy covariance, gradient method, methane, carbon balance, tundra, climate change, land-atmosphere interactions

Received 13 November 2011 and accepted 17 December 2011

Introduction

Approximately 16% of the global soil area is underlain by permafrost (Tarnocai *et al.*, 2009). Ecosystems in these areas are vulnerable to the effects of predicted global climate change, and a range of possible feedback mechanisms to climate could arise from these ecosystems (Chapin *et al.*, 2000). Global warming is not evenly distributed over the world, and especially northern high-latitude areas are projected to see a larger increase in temperature, precipitation, and growing season length than the rest of the world (ACIA, 2005; IPCC, 2007).

Northern wet tundra ecosystems are characterized by waterlogged, anaerobic, and cold conditions, which effectively reduce decomposition rates and favor the formation of peat. The net carbon (C) emission from these ecosystems is the balance between the carbon dioxide (CO₂) absorbed by gross primary production

(GPP), and the C decomposed and released as CO₂ or methane (CH₄). During the Holocene, these areas acted as sinks for C, and approximately 50% of the global soil organic C pool is today stored in the northern permafrost regions (Smith *et al.*, 2004; McGuire *et al.*, 2009; Tarnocai *et al.*, 2009). As the permafrost areas are thawing (Serreze *et al.*, 2000), previously frozen C becomes available for decomposition, and could have a major impact on the global C cycle (Tarnocai, 2006).

The waterlogged and anaerobic conditions also make northern wetlands one of the largest natural sources of atmospheric CH₄ (Whalen, 2005). A change in this major CH₄ source would have significant effects on the greenhouse gas budgets of these areas. As concerns regarding the future climate change grow, there is an increasing need to quantify the greenhouse gas budget of global ecosystems. There are still large uncertainties regarding the CH₄ source strength of wet tundra ecosystems, and its relationship to environmental variables. For example, Mastepanov *et al.* (2008) showed unexpectedly high CH₄ fluxes during the refreezing of the active layer. Whalen & Reeburgh (1988), showed

Correspondence: Torbern Tagesson, tel. + 46 0704 993 936, fax + 46 046 222 0321, e-mail: torbern.tagesson@nateko.lu.se

similar autumn CH₄ fluxes for areas covered with mosses in Alaska where 40% of the annual fluxes were released during the freeze-in period.

Most CH₄ flux studies in high-latitude ecosystems have been based on the closed-chamber technique (e.g., Whalen & Reeburgh, 1988; Joabsson & Christensen, 2001; Corradi *et al.*, 2005; Mastepanov *et al.*, 2008). However, chamber-based measurements decouple the surface from the atmosphere, and the influence of, for example, atmospheric turbulence on the CH₄ fluxes can hereby not be studied. Net exchange of greenhouse gases between the ground surface and the atmosphere can be measured at a scale of hectares to km² with micrometeorological methods like the eddy covariance technique (EC) or in lack of fast response instruments, by using gradient methods. Only a few studies have reported CH₄ flux measurements with these methods from high-Arctic wet tundra ecosystems (Friborg *et al.*, 2000; Sachs *et al.*, 2008; Wille *et al.*, 2008).

The high-Arctic region constitutes approximately 34% of the non-glaciated Arctic area (Walker *et al.*, 2005) and is defined as areas where the average air temperature of the warmest month is below 7 °C (Bliss, 1997; Walker *et al.*, 2005). The spatial coverage of wetlands north of 45° is 3.72 × 10⁶ km², whereof 62% is underlain by permafrost (Avis *et al.*, 2011). The primary aim of this study was to use micrometeorological methods to investigate ecosystem-scale CH₄ and CO₂ fluxes in a high-Arctic wet tundra ecosystem from spring thaw, throughout the growing season, and into the freeze-in and early winter period during 2008 and 2009. As study site, representative of the non-glaciated high-Arctic areas underlain by permafrost, we selected Rylekærene a wet area situated in the Zackenberg valley, Northeast Greenland. A long time series of CH₄ fluxes in combination with long term data of other ecosystem properties, that is, CO₂ fluxes and environmental conditions, will allow us to assess the seasonal behavior of CH₄ fluxes, and the influence of ecosystem properties. Subsequently, a second aim was to identify how well these ecosystem properties determine CH₄ fluxes at an ecosystem level. In addition, our measurements of both CH₄ and CO₂ fluxes allowed us to estimate C budgets for the land-atmosphere exchange of the measured area, and a final aim was to investigate inter-annual variation in these budgets.

Materials and methods

Site description

The field measurements were carried out in the fen area, Rylekærene, in the Zackenberg valley (74°28'N 20°34'W), situated in the Northeast Greenland National Park. The average air

temperature of the warmest month in the valley is 5.8 °C, and the mean annual temperature is −9 °C (Hansen *et al.*, 2008). The snow melt usually begins in late May and ends in mid-June. Rylekærene is a typical high-Arctic heterogeneous wetland area. It is a patterned fen characterized by alternating high, dry heath areas, and low, wet fen areas. During a field inventory in 2007, the dominant plant communities were registered for every 15 m² within a 1.4 km² area covering Rylekærene (Fig. 1) (Tagesson, 2011). The dominant plant communities were fen, grassland (dominated by *Arctagrostis latifolia*, *Eriophorum triste*, and *Alopecurus alpinus*), *Cassiope tetragona* heath, *Dryas octopetala* heath, *Vaccinium uliginosum* heath and *Salix arctica* snowbed (Fig. 1) (Tagesson, 2011). The fen areas were further divided into continuous fen (flat areas dominated by *Eriophorum scheuchzeri*, *Carex stans* and *Dupontia psilosantha*) and hummocky fen (hummocks dominated by *E. triste*, *S. arctica* and *A. latifolia*) (Bay, 1998). The moss species at the site was dominated by true mosses, for example, *Tomenthypnum*, *Scorpidium*, *Aulacomnium* and *Drepanocladus*. The maximum active layer thickness in the valley ranges between 0.40 m at the fen to about 0.80 m at *Dryas* sites (Elberling *et al.*, 2008). The peat layer in the fen is 0.20–0.30 m thick (Christensen *et al.*, 2000). There were mainly mineral soils below the peat and in the remaining plant communities. The pH of the fen pore water at 10 cm peat depth was 6.9 ± 0.2 (*n* = 50; Ström, unpublished results). There have been extensive ecological, biogeographic, climatic, and hydrological research and monitoring carried out in the Zackenberg research area since 1995 (Meltofte *et al.*, 2008).

Turbulence and CO₂ flux measurements

A 3 m tower was placed in the center of Rylekærene so that fen areas were located in the main upwind direction (Fig. 1). The study took place between 25 June and 28 October 2008, and between 16 May and 24 October 2009. An EC system consisting of a 3-axis sonic anemometer (Metek, GmbH, Elmshorn, Germany), and an open-path CO₂/H₂O infrared gas analyzer (LI-7500: LI-COR Inc, Lincoln, NE, USA) was installed at 3.3 m above the surface. The gas analyzer was tilted 32° from vertical next to the sonic anemometer. This system provided both the direct CO₂ fluxes, and the turbulence data (friction velocity and stability) necessary for the further CH₄ flux calculations. The CO₂ flux calculations followed standard EC methods (Kaimal & Finnigan, 1994).

The sonic anemometer and gas analyzer data were sampled at 20 Hz, but the statistics were calculated for 30 min periods. The raw data were screened for spikes four times with a running mean of 100 values (Hojstrup, 1993). All velocity and temperature values that deviated from the running mean more than 4.5, 5.0, 5.5 and 6.0 times the running standard deviation were removed, for each run respectively. These factors were determined empirically to give an optimum despiking. Higher factors (8.0, 8.5, 9.0 and 9.5) were needed for the CO₂ concentration data. Linear detrending was applied to all data, and the 2D rotation to wind velocity data (Kaimal & Finnigan, 1994). Data were filtered for fully developed turbulent conditions according to Foken (2008) using the integral




Fig. 1 The Zackenberg valley and the investigation area surrounding the Zackenberg research station (ZERO). A field-inventory map of the dominant plant communities observed in Rylekærerne in July 2007 (Tagesson, 2011) is put on top of an IKONOS satellite image from 23 July 2000. The red mark on the Greenland map is the location of Zackenberg.

turbulence characteristics. CO₂ concentration data were filtered for steady-state conditions following Vickers & Mahrt (1997), with an empirically determined threshold value of 0.02. In addition, all half-hourly values with a spike removal of more than 1% were rejected (Foken, 2008). The CO₂ data were additionally filtered according to a diagnostic value from the LI-7500, indicating problems caused by precipitation, icing, fog, and dirt on the sensor. In total, 43% of the combined CO₂ flux and turbulence data were filtered away. Schotanus corrections were applied on the sensible heat fluxes (Schot-

anus *et al.*, 1983). Consequently, the CO₂ fluxes were Webb corrected (Webb *et al.*, 1980), which added between -5 and 10 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (on average 0.64 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$). The CO₂ fluxes were also corrected for heating of the open-path sensor itself according to Burba *et al.* (2008) (on average 1.14 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$). The correction for heating of the open-path sensor was originally designed for vertically mounted sensors. However, no correction is available for sensors mounted at an angle. The frequency losses were estimated by comparing the average of 450 half-hour co-spectra

of the vertical wind and the concentration of CO₂ against the average co-spectra of the vertical wind and the temperature. No transfer functions were applied for compensating for frequency losses (see below). Totally, all applied corrections added on average 1.77 μmol CO₂ m⁻² s⁻¹.

Methane flux measurements

We lacked a fast response instrument necessary for the EC technique to be applicable, and the CH₄ fluxes were thus estimated by combining gradient and EC methods. The CH₄ concentrations were measured at two levels (0.70 and 2.75 m) on the tower at 1 Hz rate. The system consisted of a laser off-axis integrated cavity output spectroscopy analyzer (LGR; DLT200, Fast Methane Analyzer, repeatability 1 ppb at 0.1 Hz, Los Gatos Research, Mountain View, CA, USA). Air was pumped at a rate of 4 L min⁻¹ with two membrane pumps (THOMAS EMS3025-63ri, Sheboygan, WI, USA) from the two heights through two 16 m long high-density polyethylene tubes (inner diameter 4 mm). The air from the two different heights was first passed through two 5 L mixing chambers to minimize fluctuations of CH₄ concentration. Subsamples were taken after the mixing chambers and entered the LGR at a rate of 0.4 L min⁻¹. The measurements were switched between the two levels every 5 min. The first ½ minute was discarded and remaining 4.5 min of 1 Hz data were averaged. For the flux calculations, average and standard deviation for the concentrations over the 30 min periods were calculated. Data were filtered for steady-state conditions with an empirically determined standard deviation threshold value of 20 ppb. In total, 49% of the combined CH₄ concentrations and turbulence data were filtered away.

According to Monin–Obukhov similarity theory vertical turbulent fluxes (F_{CH_4}) can be calculated as

$$F_{\text{CH}_4} = -\rho \frac{\kappa u_* (C_{\text{CH}_4, z_2} - C_{\text{CH}_4, z_1})}{\ln(z_2/z_1) - \psi_H(\zeta)_{z_2} + \psi_H(\zeta)_{z_1}} \quad (1)$$

where κ (0.4) is the von Kármán constant, u_* is the friction velocity, C_{CH_4, z_1} is the CH₄ concentration at 0.7 m and C_{CH_4, z_2}

is the CH₄ concentration at 2.75 m, z_1 is the 0.7 m measurement height, z_2 is the 2.75 m measurement height and $\psi(\zeta)$ is the integrated form of the universal function (Foken, 2008). There are few studies with universal functions for trace gases, and the functions for sensible heat (H) is widely used (Högström, 1988). The CH₄ concentrations at z_1 and z_2 were measured and u_* and H were obtained by the sonic anemometer measurements.

The footprint of the CH₄ flux measurements was estimated by a model that combines a Lagrangian stochastic dispersion model and similarity theory (Hsieh *et al.*, 2000). The model relates footprint to atmospheric stability, measurement height, and surface roughness length. The measurement height was set to the arithmetic mean for stable conditions, and the geometric mean for unstable conditions (Horst, 1999). Surface roughness length was set to 0.01 m (Wieringa, 1993). The Obukhov length was given by the sonic anemometer measurements. The 80% cumulative flux distance was on average 260 m, and the average point of maximum contribution was 22 m from the tower. The average 80% cumulative flux distance is shown in Fig. 1. The fractions of different plant communities for different wind directions are included in Table 1.

Environmental variables

Soil temperature at 0.05, 0.10 and 0.15 m depth were measured (Tinytag Plus, Gemini Data Loggers, Chichester, UK) every 5 min at a site 265 m south of the tower (Fig. 1). In addition, soil temperature at a depth of 0.02 m ($T_{0.02}$) was measured every 10 min at the tower site using Copper-Constantan thermocouples. Photosynthetic active radiation (PAR) was measured with a JYP-1000 sensor (SDEC, Tauxigny, France). Soil gas pressure profiles relative to the atmospheric pressure (24PCEFA6G, Honeywell S&C, Morristown, NJ, USA) were measured at four plots in the surroundings of the tower at 0.10, 0.20, 0.30 and 0.40 m depth from the 20 July until the soil temperature dropped down to 0 °C in 2008. No pressure sensors were used during 2009. Some of the sensors broke during the field season, and only data from two sensors at 0.10 m, one sensor at 0.20 m, four sensors at 0.30 m, and three sensors

Table 1 Fractions of the plant communities in the CH₄ flux footprint area of the tower. Quadrant 1 is between 315° and 45°, quadrant 2 is between 45° and 135°, quadrant 3 is between 135° and 225° and quadrant 4 is between 225° and 315°. The fractions for the dominant wind direction ± 20° for the growing season and the autumn and early winter periods are also included

Plant community	Entire footprint	Quadrant 1	Quadrant 2	Quadrant 3	Quadrant 4	Dominant wind direction growing season	Dominant wind direction autumn/early winter
Continuous fen	0.25	0.40	0.23	0.28	0.14	0.38	0.49
Hummocky fen	0.11	0.23	0.13	0.07	0.09	0.05	0.26
Grassland	0.25	0.23	0.18	0.21	0.37	0.27	0.22
<i>Cassiope tetragona</i>	0.07	0.03	0.15	0.07	0.03	0.06	0.02
<i>Vaccinium uliginosum</i>	0.02	0.02	0.01	0.03	0.00	0.02	0.01
<i>Dryas octopetala</i>	0.02	0.00	0.00	0.05	0.01	0.04	0.00
<i>Salix arctica</i>	0.16	0.07	0.30	0.14	0.13	0.11	0.00
Water	0.04	0.00	0.00	0.05	0.09	0.01	0.00
Gravel	0.08	0.01	0.00	0.10	0.14	0.06	0.00

at 0.40 m depth were used in the final analysis. The soil gas pressures were corrected for temperature effects on the sensors.

All sensors were connected to a CR-1000 data logger in combination with a multiplexer (Campbell scientific Inc., North Logan, UT, USA). Data were sampled every 30 s, and stored as 10 min averages.

Active layer thickness (AL) and water table depth (WtD) were measured manually during 15 occasions at 15 plots around the tower in 2008. Active layer thickness was measured at five plots and WtD at one plot during 33 occasions in 2009. Precipitation and snow depth were measured at the Climate station (C1) in Fig. 1 (ClimateBasis, 2010).

Data handling

Partitioning of CO₂ fluxes into GPP and ecosystem respiration. The measured net ecosystem exchange (NEE) of CO₂ from the growing season was partitioned into GPP, and ecosystem respiration (ER). The growing season was defined to start the day snow depth exceeded 0.10 m and end the first of two consecutive days with T_{0.02} below 0 °C (Tamstorf *et al.*, 2007). The 2008 growing season data were divided into 4 day data sets, except for the last period, which got 7 days. For 2009, the data were divided into 4 day data sets, except for the beginning (1–12 June 2009) and end of the growing season (after the 19 August), which were divided into 12 and 9.5 day intervals, respectively. The reason for the longer periods in the beginning and the end of the growing seasons was that GPP constitute a small fraction of the measured data and it was hereby harder to separate it from ER. A light response curve was fitted to each data set:

$$NEE = -(F_{\text{csat}} + R_d)(1 - e^{\frac{-\alpha(\text{PAR})}{F_{\text{csat}} + R_d}}) + R_d \quad (2)$$

where F_{csat} is CO₂ flux at light saturation, R_d is the ecosystem dark respiration, and α is initial slope of the light response curve (Falge *et al.*, 2001). By subtracting R_d , the curve was forced through zero, and GPP could hereby be estimated:

$$GPP = -(F_{\text{csat}} + R_d)(1 - e^{\frac{-\alpha(\text{PAR})}{F_{\text{csat}} + R_d}}) \quad (3)$$

Ecosystem respiration was calculated by subtracting the modeled GPP from the measured NEE values. The light response curves were well fitted for all periods, except the 1–12 June 2009. For partitioning of this period, an exponential regression was fitted with ER against air temperature, with data measured 7 days before, and after this period. Ecosystem respiration for the 1 June to 12 June 2009 was estimated using the regression on the air temperature data set, and GPP was estimated by subtracting modeled ER from measured NEE.

The budgets of the land-atmosphere exchange of carbon. For calculation of the CH₄ and CO₂ budgets, the time series needed to be gap-filled. Gaps were filled from 24 June to 31 October 2008, and 16 May to 31 October 2009. Gaps in data originated from, (1) the filtering described above, (2) a late start-up during the growing season 2008, (3) the measurements were stopped before the 31 October 2008 and 2009, (4)

power failures, (5) technical issues, and (6) a polar bear attack. The gaps in the CH₄ fluxes were 57%, the gaps in GPP were 7%, and the gaps in ER were 49% of the full time series. Gaps that were shorter than or equal to 2 h were gap-filled by linear interpolation between the two neighboring values. Remaining gaps in the CH₄ fluxes that were shorter than 24 h were set to daily averages. Gaps in daily average CH₄ fluxes during the growing season were filled with a linear regression model against T_{0.02}, whereas outside the growing seasons, they were filled by linear interpolation between daily averages the days before, and after the gap. For the remaining gaps in the GPP and ER time series, a mean diurnal value of a 7 day gliding window was used (Falge *et al.*, 2001).

The errors in the CO₂ flux budgets were estimated following Aurela *et al.* (2002). These estimates do not cover all error sources, but gives an estimate of the key uncertainties in the flux measurements. Aurela *et al.* (2002) used closed-path EC measurements. The errors associated with frequency losses were thus estimated by comparing the average half-hour co-spectra of the vertical wind and the concentration of CO₂ against the average co-spectra of the vertical wind and the temperature for five different groups with 90 half-hour values each.

The random errors in the CH₄ flux measurements were calculated by setting a measurement error of u_* (E_{u^*}) to 0.05 m s⁻¹, an error of 10 W for H (E_H) (Foken, 2008) and an error of 1 ppb for the CH₄ concentration difference between 0.7 and 2.75 m ($E_{\Delta\text{CH}_4}$) (Los Gatos Research). The errors in CH₄ flux budgets associated with the gap filling (E_{gf}) and the filtering of the measured data (E_f) were estimated following Aurela *et al.* (2002). Finally all errors (E_t) were added together using the error accumulation principle:

$$E_t = \sqrt{E_{u^*}^2 + E_H^2 + E_{\Delta\text{CH}_4}^2 + E_{\text{gf}}^2 + E_f^2} \quad (4)$$

Seasonal dynamics of CH₄ fluxes. We divided the measurement periods into three different seasons, based on the stages of physical and biological conditions; (i) spring, (ii) growing season, and (iii) autumn and early winter. As stated above the growing season was defined as ranging from the day the snow depth exceeded 0.10 m until the first of two consecutive days with soil temperature at 0.02 m depth below 0 °C (Tamstorf *et al.*, 2007). Autumn and early winter was defined to start the day after the end of the growing season until the 31 October. Spring 2009 was set to start on 16 May, that is, date of the first CH₄ flux measurements and end when the growing season started. Dates for the different seasons are given in Table 2.

Table 2 The three different seasons in which the measurement periods were divided into for 2008 and 2009

Season	2008	2009
Spring	–	16 May–31 May
Growing season	24 June–16 September	1 June–6 September
Autumn and early winter	17 September–31 October	7 September–31 October

The relationships between daily average CH₄ fluxes and daily average soil temperatures at the different depths, GPP, WtD, AL, and friction velocity were tested by fitting linear and exponential regressions. Data from the gaps in the daily average CH₄ fluxes were not used in the analysis against environmental variables, as they were not true measured data. An exponential regression between half-hourly CH₄ fluxes and friction velocity measured after soil was completely frozen 2008 was also fitted.

Diurnal dynamics of CH₄ fluxes. We analyzed the different seasons for diurnal variation. The CH₄ fluxes were normalized by dividing each half-hourly flux value with the mean daily flux. Only days with more than 70% data coverage were used in the analysis (in total 29 and 27 days for 2008 and 2009, respectively).

To avoid effects of the seasonal variation, 1 week data sets at the peak of the growing seasons were used to study variation in half-hourly CH₄ fluxes. The data sets were from 26 July to 1 August 2008 and from 6 July to 12 July 2009. The variation in half-hourly CH₄ fluxes during these periods was analyzed by fitting exponential and linear regressions against half-hour values of soil temperature at the different depths, GPP, fraction of emissions from the different plant communities in the footprint and friction velocity.

Results

Environmental conditions

Average air temperature during June–August was 6.9 °C in 2008 and 5.0 °C in 2009. Both years were above the long-term average of 4.6 °C since the start of the measurements in 1996 (ClimateBasis, 2010). The AL reached its maximum (~0.50 m) in the beginning of August both years (Figs 2c and 3c). Maximum snow depths were 1.30 and 0.17 m in the winters preceding the growing seasons 2008 and 2009, respectively. The area became snow free on 24 June 2008 and on 1 June 2009. It rained 161 and 117 mm in 2008 and 2009, respectively. The difference in temperatures and precipitation lead to different hydrological conditions, and the water table was closer to the soil surface in 2008 (deepest WtD: –0.03 m) than in 2009 (deepest WtD: –0.10 m) (Figs 2d and 3d).

There were no changes in the soil gas pressures until the soil froze at any of the depths (Fig. 2f). After the soil started to freeze the 23 September 2008, there was a strong increase in soil gas pressure at all depths, gradually increasing as the frozen ground advanced downwards.

Diurnal variation in CH₄ fluxes

There were large variations in half-hourly CH₄ fluxes, but no systematic diurnal variation could be seen for

any of the seasons 2008 or 2009. Normalized half-hourly data from the growing seasons 2008 and 2009 are shown in Fig. 4. Shorter periods within the different seasons were also studied, but no diurnal variation was seen in these periods either.

None of the environmental factors gave any good degree of explanation to the variation in half-hourly CH₄ fluxes during the peak of the growing season. The fraction of emission that derived from continuous fen was the factor that best explained the variation in half-hourly CH₄ fluxes (linear correlations: 2008: *P*-value = 0.0002, *R*²=0.17, *n* = 80; 2009: *P*-value < 0.0001, *R*²=0.09, *n* = 181).

Seasonal dynamics in CH₄ fluxes

Spring 2009. The CH₄ flux was on average 0.18 mg CH₄ m⁻² h⁻¹ during spring 2009. The highest CH₄ emissions with values up to 1.75 mg CH₄ m⁻² h⁻¹ were observed on 21, 23 and 28 May 2009. These emissions do however not have any significant impacts on a seasonal or annual budget since they existed only for short periods of time (Fig. 3a). There were no correlations between variation in CH₄ fluxes and any of the measured variables. The reason for the lack of correlations was the low steady CH₄ fluxes without any large variations.

Growing seasons 2008 and 2009. The average growing season CH₄ fluxes were larger in 2008 than in 2009, that is, 4.60 mg and 2.94 mg CH₄ m⁻² h⁻¹, respectively. Peak daily average CH₄ fluxes were 10.62 mg CH₄ m⁻² h⁻¹ the 5 August in 2008, and 6.64 mg CH₄ m⁻² h⁻¹ the 9 July in 2009 (Figs 2a and 3a).

All input variables were co-varying over the season. The temporal autocorrelation meant that the data could not be regarded as statistically independent, and the underlying assumptions for testing of significance in a regression analysis were hereby not fulfilled. It is consequently hard to tell which exact factors control the CH₄ fluxes. Despite this we still used regression analysis as an attempt to investigate how well the factors determine the CH₄ fluxes. We only present the coefficient of determination (*R*² values), whereas the significance level (*P*-values) is not presented in the results. The lack of diurnal cycles allowed us to average fluxes daily, since missing data only increased uncertainty, but did not add any systematic errors.

Soil temperature at 0.02 m was the depth that best explained the variation in daily average CH₄ fluxes (linear correlation: 2008: *R*² = 0.79, *n* = 73; 2009: *R*² = 0.55, *n* = 98) (Fig. 5a). There was also a strong linear correlation between daily average CH₄ fluxes




Fig. 2 Time series of CH₄ fluxes, CO₂ fluxes, and environmental variables during 2008. (a) CH₄ fluxes, (b) gross primary production (GPP), and ecosystem respiration (ER), (c) soil temperature (T) at 0.05, 0.10 and 0.15 m depth, and active layer thickness (AL), (d) precipitation and water table depth (WtD), (e) average daily wind speed and (f) soil gas pressure (P) normalized by the annual mean together with soil temperature at 0.02 m depth. The dotted lines show the boundary between the different seasons.

and GPP (2008: $R^2 = 0.72$, $n = 73$; 2009: $R^2 = 0.66$, $n = 98$) (Fig. 5b).

No correlation between daily average CH₄ fluxes and WtD existed for the growing season in 2009 ($R^2 = 0.03$, $n = 26$) and a negative linear correlation ($R^2 = 0.22$, $n = 14$) was seen in 2008. There was a strong linear correlation between daily average CH₄ fluxes and AL, if only data up until the peak of the growing seasons were used (2008: $R^2 = 0.58$, $n = 8$; 2009: $R^2 = 0.90$, $n = 10$). No correlation was seen between CH₄ fluxes and friction velocity for either 2008 or 2009.

Autumn and early winter 2008 and 2009. Average autumn and early winter CH₄ fluxes were 0.74 and 1.22 mg CH₄ m⁻² h⁻¹ for 2008 and 2009, respectively. There were no increases in CH₄ fluxes after the soil started to freeze at 0.02 m depth on 23 September 2008, and on 3 October 2009. There were no correlations between daily average CH₄ fluxes and any of the environmental variables. Large CH₄ bursts were seen, at the 21 and 24–25 October 2008, and these occurred during very windy conditions. For the data measured after soil was completely frozen (18 October 2008) there was an exponential relationship ($R^2 = 0.63$,




Fig. 3 Time series of CH₄ fluxes, CO₂ fluxes, and environmental variables during 2009. (a) CH₄ fluxes, (b) gross primary production (GPP), and ecosystem respiration (ER), (c) soil temperature (T) at 0.05, 0.10 and 0.15 m depth and active layer thickness (AL), (d) precipitation, and water table depth (WtD) and (e) average daily wind speed in m s^{-1} . No soil gas pressure estimates were done during 2009. The dotted lines show the boundary between the different seasons.

$n = 322$) between half-hourly CH₄ fluxes and friction velocity (Fig. 6).

The budgets of the land-atmosphere exchange of carbon

If all C fluxes were added together (24 June to 31 October 2008 and 16 May to 31 October 2009), Rylekærene acted as a C source for the measurement period 2008 ($+3.8 \text{ g C m}^{-2}$), whereas it acted as a C sink for the measurement period 2009 (-47.6 g C m^{-2}). Over the measurement period, 7.6 and 6.5 g C m^{-2} were emitted as CH₄ in 2008 and in 2009, respectively. During the measurement period 2008, Rylekærene switched from being

a sink of -3.8 g C m^{-2} to a source of $+3.8 \text{ g C m}^{-2}$ because of the CH₄ emissions. During the measurement period 2009, 14% of the C balance was emitted as CH₄. The C budgets for the different seasons are shown in Table 3.

The error analysis for the CO₂ flux measurements and the gap filling procedure resulted in total accumulated errors of ± 5.68 and $\pm 4.45 \text{ g C}$ for 2008 and 2009, respectively. The error analysis of the CH₄ flux measurements and gap filling of the CH₄ fluxes resulted in total accumulated errors of ± 0.63 and $\pm 0.77 \text{ g C}$ for 2008 and 2009, respectively. According to the co-spectrum analysis, 99% of the frequencies were measured by the




Fig. 4 Half-hourly CH₄ fluxes normalized by the daily mean flux. Only data from the growing seasons 2008 and 2009 with daily data coverage larger than 70% were used.

system. No transfer functions were thus applied for compensating for losses of high and low frequencies. The separate components of the error analysis are given in Table 4.

Discussion

Seasonal dynamics in CH₄ flux

Spring 2009. The increased spring-time emissions were observed simultaneously to the start of snow melt, possibly indicating a release of CH₄ from the snow cover, as it settled during snow thawing. These were minor releases during short periods of time, and no extra CH₄ bursts occurred as the ice covering the fen was melting. The increased CH₄ emissions were very small in comparison to previous studies showing bursts as the soil thaws. Hargreaves *et al.* (2001) showed that 11% of total annual CH₄ fluxes were released during spring thaw using the EC technique, whereas both Wille *et al.* (2008) (using the EC technique) and Tokida *et al.* (2007) (using closed chambers) showed spring thaw bursts equal in size to episodic ebullition releases during high summer.

Growing season. The start of the growing season greenhouse gas exchange at high-Arctic areas is strongly controlled by timing of the snow melt (Groendahl *et al.*, 2007; Mastepanov, 2010). This was confirmed in the present study as both years had similar CH₄ fluxes 2 days after snow melt. The increase over the season was much faster in 2008 than in 2009. The main reason was most likely the more rapid increase in soil temperature in 2008 than in 2009. The relationship between CH₄ emissions and soil temperature has previously been shown to have both strong exponential relation-




Fig. 5 (a) Daily average CH₄ fluxes against soil temperature at 0.02 m depth ($T_{0.02}$) and (b) against gross primary production (GPP) for the growing seasons 2008 and 2009. The solid lines are regressions for the 2008 data series, and the dotted lines are regressions for the 2009 data series (2008: CH₄ flux = $0.60 \times T_{0.02} + 0.57$, $R^2 = 0.79$, $n = 73$; 2009: CH₄ flux = $0.37 \times T_{0.02} + 0.01$, $R^2 = 0.55$, $n = 98$); (2008: CH₄ flux = $-0.007 \times \text{GPP} + 1.90$, $R^2 = 0.72$, $n = 73$; 2009: CH₄ flux = $-0.004 \times \text{GPP} + 1.08$, $R^2 = 0.66$, $n = 98$).

ships (Hargreaves *et al.*, 2001; Rinne *et al.*, 2007; Wille *et al.*, 2008) and relationships that were weak or absent (Wickland *et al.*, 2006; Sachs *et al.*, 2008). When the relationship between the net CH₄ fluxes and soil temperature deviates from an exponential one, this implies that other factors than CH₄ production exert a strong control on the net emissions. Examples of such factors could be metabolic activity of the methanotrophs, sizes of the zones for CH₄ production and oxidation, substrate availability, and storage or resistances along the transport pathways from the site of production to the atmosphere.

The WtD regulates the sizes of the zones for CH₄ production and oxidation (Bubier & Moore, 1994). There was no consistent relationship between seasonal




Fig. 6 Half-hourly CH_4 fluxes against friction velocity (u_*), after the soil was completely frozen the 18 October 2008. The solid line is an exponential regression with CH_4 flux against u_* ($\text{CH}_4\text{flux} = 0.09 \times e^{6.39 \times u_*} - 0.18$, $R^2 = 0.63$, $n = 322$).

variation in CH_4 fluxes and WtD, that is, a lack of correlation 2009 and a negative correlation 2008. An explanation could be that the WtD is usually at its peak right after the snow melt, where after it decreases over the summer as the water runs off or is evapotranspired. Low CH_4 fluxes were consequently measured in the beginning of the growing seasons, when the WtD was high, and high fluxes at the peak of the growing season when the WtD was low. The water table was closer to the soil surface in 2008 than in 2009. It may be suggested that WtD explains the inter-annual variation in CH_4 flux whereas there are other factors with a stronger control on the annual CH_4 fluxes. In agreement with this reasoning, many previous studies showing a positive relationship between CH_4 flux and WtD have mainly showed spatial and long-term dependences (Hargreaves & Fowler, 1998; Huttunen *et al.*, 2003; Bu-

bier *et al.*, 2005). Another factor setting the zone for CH_4 production is the AL. The main part of the CH_4 fluxes originates from the upper part of the soil (Whalen, 2005), and CH_4 fluxes were therefore not affected by AL when it approached its maximum.

The quantity and quality of the substrate to be decomposed are additional important environmental variables regulating CH_4 fluxes (Whiting & Chanton, 1993; Joabsson & Christensen, 2001). The most important substrates for methanogens are fresh organic matter from root exudates, and recently produced litter (Chanton *et al.*, 1995; Ström & Christensen, 2007). The availability of fresh organic matter is closely linked to GPP (Joabsson & Christensen, 2001), hence the observed strong relationship between the CH_4 fluxes and GPP. This has also been shown in previous studies (Whiting & Chanton, 1993; Friberg *et al.*, 2000; Joabsson & Christensen, 2001).

No correlation was seen between CH_4 fluxes and friction velocity during the growing seasons. It has been shown that there are high correlations between atmospheric turbulence and diffusive transfers between water–air interfaces (MacIntyre *et al.*, 1995) and most studies reporting an effect of friction velocity have had a high surface coverage of water bodies (MacIntyre *et al.*, 1995; Hargreaves *et al.*, 2001; Sachs *et al.*, 2008; Wille *et al.*, 2008).

Autumn. Our study did not show any autumn CH_4 emission pulse during the onset of soil freezing, as has been suggested by land-atmosphere CH_4 flux measurements, and atmospheric concentration measurements at high northern latitudes (Khalil & Rasmussen, 1983; Dlugokencky *et al.*, 1994; Mastepanov *et al.*, 2008). There are few studies addressing the issue of wintertime CH_4 fluxes in permafrost areas (Whalen &

Table 3 The carbon budgets for the different seasons. The periods for the different seasons are given in Table 2. All daily average flux values are in g C m^{-2} per 24 h period. The Flux budgets are total accumulated flux in g C m^{-2} . No measurements were done during spring 2008

Flux	Spring	Growing season		Autumn and early winter		Entire measurement period	
	2009	2008	2009	2008	2009	2008	2009
Daily CH_4 fluxes	0.003	0.084	0.051	0.011	0.027	0.059	0.039
Daily GPP	0.00	-2.57	-3.11	0.00	0.00	-1.68	-1.80
Daily ER	0.37	1.89	2.16	1.19	0.60	1.65	1.49
Daily net C exchange	0.37	-0.59	-0.90	1.20	0.63	0.03	-0.28
CH_4 flux budgets	0.1	7.1	5.0	0.5	1.5	7.6	6.5
GPP budgets	0.0	-218.1	-304.9	0.0	0.0	-218.1	-304.9
ER budgets	5.9	160.9	211.7	53.5	33.2	214.3	250.8
Net C exchange budgets	6.0	-50.1	-88.2	53.9	34.7	3.8	-47.6

GPP, gross primary production; ER, ecosystem respiration.

Table 4 The separate error components from the error analysis of CO₂ and CH₄ flux budgets. The errors are in g C for the entire measurement period. There were no errors associated with high and low frequency loss for the CH₄ flux measurements as it was not based on fast gas concentration measurements

	2008	2009
Error CO ₂ flux measurements (g C)		
Random	±0.26	±0.32
Gap filling	±5.36	±4.24
Frequency losses	±0.04	±0.54
Filtering for steady state conditions	±0.59	±0.60
Filtering for turbulent conditions	±1.77	±1.05
Total	±5.68	±4.45
Error CH ₄ flux measurements (g C)		
Friction velocity measurements	±0.41	±0.55
Sensible heat flux measurements	±0.18	±0.23
CH ₄ gradient measurements	±0.22	±0.29
Gap filling	±0.24	±0.08
Filtering for steady state conditions	±0.28	±0.39
Filtering for turbulent conditions	±0.10	±0.09
Total	±0.63	±0.77

Reeburgh, 1988; Mastepanov *et al.*, 2008; Wille *et al.*, 2008). Wille *et al.* (2008) did not see any increase in the wintertime CH₄ fluxes as the soil started to freeze in the Lena delta, north-eastern Siberia. Whalen & Reeburgh (1988) showed that 40% of the annual fluxes were winter fluxes from moss covered surfaces in Alaska, whereas no increases were seen for other vegetation covers. Mastepanov *et al.* (2008), showed that an equal amount of CH₄ was emitted during the freeze-in period as was emitted during the entire growing season in 2007 at the edge of Rylekærene.

Both Mastepanov *et al.* (2008) and Whalen & Reeburgh (1988) explained the increase in CH₄ fluxes as the soil started to freeze with physical processes. The soil gas pressure increased as the soil started to freeze down toward the permafrost bottom. Accumulated CH₄ is prevented from diffusing downwards due to the impermeable permafrost, and it is hereby pressed up through the soil matrix. Pathways through the soil could both be voids in the frozen moss matrix (Whalen & Reeburgh, 1988), and through the aerenchyma of senescent vascular plants (Hargreaves *et al.*, 2001; Mastepanov, 2010). In accordance with this hypothesis, there was indeed a strong increase in soil gas pressure as the soil started to freeze the 23 September 2008 (Fig. 2f). Despite the increase in soil pressure, there was no increase in the CH₄ fluxes.

A plausible explanation for the lack of autumn bursts in 2008 and 2009 is that other environmental conditions need to be fulfilled in order for the burst to occur. It was not enough with an increase in soil gas pressure. It

could be that it was also necessary with an accumulation of CH₄ produced during the growing season. Whalen & Reeburgh (1988) only observed increased autumn emissions in moss covered areas, whereas measurements at plots with vascular plants did not show any increase in CH₄ fluxes during soil freezing. Shannon *et al.* (1996), have shown that CH₄ transporting plants deplete both the dissolved CH₄ pool in the pore water, and the concentration of CH₄ in trapped gas bubbles in the saturated rhizosphere. In case there are no vascular plants, a larger fraction of the produced CH₄ is stored in the soil. This CH₄ can then be released due to the increased soil pressure during soil freezing. There was no apparent difference in vascular plant coverage between the site where the Mastepanov *et al.* (2008) chambers were positioned and the main parts of Rylekærene (where the current study was conducted). The Mastepanov *et al.* (2008) site, however, has a slightly different vascular plant composition with a higher coverage of the grass *A. latifolia*, indicating lower plant mediated CH₄ transport. This could possibly result in a CH₄ accumulation at the Mastepanov *et al.* (2008) site.

An interesting feature was the bursts of CH₄ during the storm events in October 2008. It might be that accumulated CH₄ existed in cavities in the moss and vegetation layer, and in cracks in the frozen soil. The high wind speed increased the turbulence, which likely resulted in turbulence-induced large-scale ebullition, possibly triggered by changes in air pressure, and ventilation of these cavities. This has also been observed in other studies (Massman *et al.*, 1997; Black *et al.*, 2000; Aurela *et al.*, 2002). No similar phenomena were seen in 2009. However, there was a 0.30 m deep snow layer covering the area in 2009, and an increase in turbulence could not affect the soil due to this protective layer.

Diurnal variation in CH₄ flux

Diurnal variation in CH₄ fluxes is often site specific and has been observed at some sites (Shannon *et al.*, 1996; Hargreaves & Fowler, 1998; Kim *et al.*, 1998), whereas not at others (Kormann *et al.*, 2001; Rinne *et al.*, 2007). We did not find any indications of diurnal variations in the present study. It could be that the lack of diurnal variation was associated with the absence of true night conditions during the main part of the growing seasons. Average nighttime (22:00–6:30 hours) PAR was 103 and 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for the growing seasons 2008 and 2009, respectively. The lack of diurnal variation indicates that the CH₄ fluxes were not directly affected by the stomatal closure or had other pathways than the stomata for CH₄ transport to the atmosphere

(Morrissey *et al.*, 1993). It could also be that the vegetation had sufficient water and was not limited by transpiration.

Tagesson (2011) showed that there were CH₄ emissions from continuous and hummocky fen, whereas they were undetectable in the dry plant communities (grassland, *S. arctica* snowbed, and the heath communities). This could explain the relationship between half-hourly CH₄ fluxes and fraction of continuous fen in the footprint of the tower. The continuous fen had a water table close to the soil surface and a high coverage of vascular plants; consequently it had the largest CH₄ fluxes. Furthermore, species composition of vascular plants can affect CH₄ emissions and substrate availability for methanogens and pointed to the importance of *Eriophorum* species (*E. scheuchzeri*, *E. angustifolium* and *E. vaginatum*) in this respect (Ström *et al.*, 2005, 2012; Ström & Christensen, 2007). These species are common in the continuous fen areas.

Representativeness of the site

Walker *et al.* (2005) made a hierarchical subdivision of the Arctic vegetation along an altitudinal temperature gradient from north to south, that is, (A) cushion-forb, (B) prostrate dwarf-shrub, (C) hemiprostrate dwarf-shrub (D) erect dwarf-shrub, and (E) low shrub subzones. The main features distinguishing Subzone C from Subzone B are the presence of the hemiprostrate shrub *C. tetragona* and well-differentiated plant communities in mires, snowbeds and creek sides. The mean July temperature at the southern boundary of Subzone B is approximately 5 °C and of Subzone C about 7 °C (Walker *et al.*, 2005). The average air temperature of the warmest month in the Zackenberg valley is 5.8 °C. Based solely on temperature the Zackenberg valley would belong to subzone B. However, the presence of fen, grassland, *C. tetragona* heath, *D. octopetala* heath, *V. uliginosum* heath and *S. arctica* snowbed (Bay, 1998) would primarily put the valley in to subzone C. Combined subzone B and C covers about 32% of the tundra zone, including most of the islands in the Canadian Archipelago, most of northern Greenland, south-western Svalbard, Novaya Zemlya, most of the northern fringe of mainland Russia, and the New Siberian Islands (Walker *et al.*, 2005). Thus, the plant communities in the Zackenberg valley and in Rylekærene comprise a good example of this part of the Arctic tundra.

The budgets of the land-atmosphere exchange of carbon

The carbon budgets were quite large, but in the same range as other Arctic studies (Corradi *et al.*, 2005; Wille *et al.*, 2008). Both total growing season and daily

averages of GPP and ER were larger in 2009 than in 2008 (Table 3), whereas the total growing season CH₄ fluxes were larger in 2008. There was no difference in average incoming PAR, whereas both temperature and WtD were higher in 2008 than in 2009. The growing season constituted a larger fraction of the measurement period in 2008 than in 2009 and the sink function for 2008 should thus be larger in comparison to 2009. It could be that the vegetation was water stressed in 2008, as the WtD was almost at the soil surface throughout the entire growing season. Further, it could be that the drier conditions in 2009 increased the mineralization rates resulting in increased GPP; ER was also higher in 2009 than in 2008 in accordance with this hypothesis. An explanation to the low ER in 2008 could be the very shallow aerobic zone. This lowered the fraction of C respired as CO₂, and resulted in a larger fraction respired as CH₄. The % of the C balance that was emitted as CH₄ was similar to studies in the Siberian arctic tundra (Friborg *et al.*, 2003; Wille *et al.*, 2008). The year 2008 was warmer and wetter than 2009. Global circulation models predict increased temperatures and increased precipitation in the future for the northern latitude ecosystems. Arctic wetlands such as Rylekærene may thus play a more significant role for the climate in the future.

Acknowledgements

The work was funded by FORMAS. We are grateful to personnel at Zackenberg field station for logistic support.

References

- ACIA (2005) *Arctic Climate Impact Assessment*. Cambridge University Press, New York.
- Aurela M, Laurila T, Tuovinen JP (2002) Annual CO₂ balance of a subarctic fen in northern Europe: importance of the wintertime efflux. *Journal of Geophysical Research*, **107**, 4607.
- Avis CA, Weaver AJ, Meissner KJ (2011) Reduction in areal extent of high-latitude wetlands in response to permafrost thaw. *Nature Geoscience*, **4**, 444–448.
- Bay C (1998) *Vegetation Mapping of Zackenberg valley, Northeast Greenland*. Danish Polar Center & Botanical Museum, University of Copenhagen, Copenhagen.
- Black TA, Chen WJ, Barr AG *et al.* (2000) Increased carbon sequestration by a boreal deciduous forest in years with a warm spring. *Geophysical Research Letters*, **27**, 1271–1274.
- Bliss LC (1997) Arctic ecosystems of North America. In: *Polar and Alpine Tundra* (ed. Wielgolaski FE), pp. 551–683. Elsevier, Amsterdam, NL.
- Bubier JL, Moore TR (1994) An ecological perspective on methane emissions from northern wetlands. *Trends in Ecology & Evolution*, **9**, 460–464.
- Bubier J, Moore T, Savage K, Crill P (2005) A comparison of methane flux in a boreal landscape between a dry and a wet year. *Global Biogeochemical Cycles*, **19**, GB1023.
- Burba GG, McDermitt DK, Grelle A, Anderson DJ, Xu L (2008) Addressing the influence of instrument surface heat exchange on the measurements of CO₂ flux from open-path gas analyzers. *Global Change Biology*, **14**, 1854–1876.
- Chanton JP, Bauer JE, Glaser PA *et al.* (1995) Radiocarbon evidence for the substrates supporting methane formation within northern Minnesota peatlands. *Geochimica et Cosmochimica Acta*, **59**, 3663–3668.
- Chapin FS, McGuire AD, Randerson J *et al.* (2000) Arctic and boreal ecosystems of western North America as components of the climate system. *Global Change Biology*, **6**, 211–223.

- Christensen TR, Friberg T, Sommerkorn M *et al.* (2000) Trace gas exchange in a high-arctic valley 1. Variations in CO₂ and CH₄ flux between tundra vegetation types. *Global Biogeochemical Cycles*, **14**, 701–713.
- ClimateBasis (2010) *The Zackenberg Database*. Available at: <http://zdb.dmu.dk>. (accessed 10 February 2010)
- Corradi C, Kolle O, Walter K, Zimov SA, Schulze ED (2005) Carbon dioxide and methane exchange of a north-east Siberian tussock tundra. *Global Change Biology*, **11**, 1910–1925.
- Dlugokencky EJ, Steele LP, Lang PM, Masarie KA (1994) The growth rate and distribution of atmospheric methane. *Journal of Geophysical Research*, **99**, 17021–17043.
- Elberling B, Tamstorf MP, Michelsen A *et al.* (2008) Soil and plant community characteristics and dynamics at Zackenberg. In: *Advances in Ecological Research-High Arctic Ecosystem Dynamics in a Changing Climate* (eds Meltofte H, Christensen TR, Elberling B, Forchhammer MC, Rasch M), pp. 223–248. Academic Press, Amsterdam.
- Falge E, Baldocchi D, Olson R *et al.* (2001) Gap filling strategies for defensible annual sums of net ecosystem exchange. *Agricultural and Forest Meteorology*, **107**, 43–69.
- Foken T (2008) *Micrometeorology*. Springer-Verlag, Berlin.
- Friberg T, Christensen TR, Hansen BU, Nordstroem C, Soegaard H (2000) Trace gas exchange in a high-arctic valley 2. Landscape CH₄ fluxes measured and modeled using eddy correlation data. *Global Biogeochemical Cycles*, **14**, 715–723.
- Friberg T, Soegaard H, Christensen TR, Lloyd CR, Panikov NS (2003) Siberian wetlands: where a sink is a source. *Geophysical Research Letters*, **30**, 2129.
- Groendahl L, Friberg T, Soegaard H (2007) Temperature and snow-melt controls on interannual variability in carbon exchange in the high Arctic. *Theoretical and Applied Climatology*, **88**, 111–125.
- Hansen BU, Sigsgaard C, Rasmussen L *et al.* (2008) Present-day climate at Zackenberg. In: *Advances in Ecological Research-High-Arctic Ecosystem Dynamics in a Changing Climate* (eds Meltofte H, Christensen TR, Elberling B, Forchhammer MC, Rasch M), pp. 111–149. Academic Press, Amsterdam.
- Hargreaves KJ, Fowler D (1998) Quantifying the effects of water table and soil temperature on the emission of methane from peat wetland at the field scale. *Atmospheric Environment*, **32**, 3275–3282.
- Hargreaves KJ, Fowler D, Pitcairn CER, Aurela M (2001) Annual methane emission from Finnish mires estimated from eddy covariance campaign measurements. *Theoretical and Applied Climatology*, **70**, 203–213.
- Högström U (1988) Non-dimensional wind and temperature profiles in the atmospheric surface layer: a re-evaluation. *Boundary-Layer Meteorology*, **42**, 55–78.
- Hojstrup J (1993) A statistical data screening procedure. *Measurement Science & Technology*, **4**, 153–157.
- Horst TW (1999) The footprint for estimation of atmosphere-surface exchange fluxes by profile techniques. *Boundary-Layer Meteorology*, **90**, 171–188.
- Hsieh CI, Katul G, Chi TW (2000) An approximate analytical model for footprint estimation of scalar fluxes in thermally stratified atmospheric flows. *Advances in Water Resources*, **23**, 765–772.
- Huttunen JT, Nykänen H, Turunen J, Martikainen PJ (2003) Methane emissions from natural peatlands in the northern boreal zone in Finland, Fennoscandia. *Atmospheric Environment*, **37**, 147–151.
- IPCC (2007) *Climate change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Joabsson A, Christensen TR (2001) Methane emissions from wetlands and their relationship with vascular plants: an Arctic example. *Global Change Biology*, **7**, 919–932.
- Kaimal JC, Finnigan J (1994) *Atmospheric boundary layer flows: their structure and measurement*. Oxford University Press, Oxford.
- Khalil MAK, Rasmussen RA (1983) Sources, sinks, and seasonal cycles of atmospheric methane. *Journal of Geophysical Research*, **88**, 5131–5144.
- Kim JN, Verma SB, Billesbach DP (1998) Seasonal variation in methane emission from a temperate Phragmites-dominated marsh: effect of growth stage and plant-mediated transport. *Global Change Biology*, **5**, 433–440.
- Kormann R, Müller H, Werle P (2001) Eddy flux measurements of methane over the fen “Murnauer Moos”, 11°11'E, 47°39'N, using a fast tunable diode laser spectrometer. *Atmospheric Environment*, **35**, 2533–2544.
- MacIntyre S, Wanninkhof R, Chanton J (1995) Trace gas exchange across the air-water interface in freshwater and coastal marine environments. In: *Biogenic Trace Gases: Measuring Emissions from Soil and Water* (eds Matson P, Harriss R), pp. 52–97. Blackwell, Oxford, UK.
- Massman WJ, Sommerfeld RA, Mosier AR, Zeller KF, Hehn TJ, Rochelle SG (1997) A model investigation of turbulence-driven pressure-pumping effects on the rate of diffusion of CO₂, N₂O, and CH₄ through layered snowpacks. *Journal of Geophysical Research*, **102**, 18851–18863.
- Mastepanov M (2010) Towards a Changed View on Greenhouse Gas Exchange in the Arctic- New Findings and Improved Techniques. Unpublished PhD thesis, Lund University, Meddelanden från Lunds Universitets Geografiska Institutioner, Avhandlingar 188, Lund, 121 pp.
- Mastepanov M, Sigsgaard C, Dlugokencky EJ, Houweling S, Ström L, Tamstorf MP, Christensen TR (2008) Large tundra methane burst during onset of freezing. *Nature*, **456**, 628–630.
- McGuire AD, Anderson LG, Christensen TR *et al.* (2009) Sensitivity of the carbon cycle in the Arctic to climate change. *Ecological Monographs*, **79**, 523–555.
- Meltofte H, Christensen TR, Elberling B, Forchhammer MC, Rasch M (2008) *Advances in Ecological Research-High Arctic Ecosystem Dynamics in a Changing Climate*. Academic Press, Amsterdam.
- Morrissey LA, Zobel DB, Livingston GP (1993) Significance of stomatal control on methane release from Carex-dominated wetlands. *Chemosphere*, **26**, 339–355.
- Rinne J, Riutta T, Pihlatie M *et al.* (2007) Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique. *Tellus B*, **59**, 449–457.
- Sachs T, Wille C, Boike J, Kutzbach L (2008) Environmental controls on ecosystem-scale CH₄ emission from polygonal tundra in the Lena River Delta, Siberia. *Journal of Geophysical Research*, **113**, G00A03.
- Schotanus P, Nieuwstadt FTM, Debruin HAR (1983) Temperature-measurement with a sonic anemometer and its application to heat and moisture fluxes. *Boundary-Layer Meteorology*, **26**, 81–93.
- Serreze MC, Walsh JE, Chapin FS III *et al.* (2000) Observational evidence of recent change in the northern high-latitude environment. *Climatic Change*, **46**, 159–207.
- Shannon RD, White JR, Lawson JE, Gilmour BS (1996) Methane efflux from emergent vegetation in peatlands. *Journal of Ecology*, **84**, 239–246.
- Smith LC, Macdonald GM, Velichko AA *et al.* (2004) Siberian peatlands: a net carbon sink and global methane source since the early Holocene. *Science*, **303**, 353–356.
- Ström L, Christensen TR (2007) Below ground carbon turnover and greenhouse gas exchanges in a sub-arctic wetland. *Soil Biology and Biochemistry*, **39**, 1689–1698.
- Ström L, Mastepanov M, Christensen TR (2005) Species-specific effects of vascular plants on carbon turnover and methane emissions from wetlands. *Biogeochemistry*, **75**, 65–82.
- Ström L, Tagesson T, Mastepanov M, Christensen TR (2012) Presence of *Eriophorum scheuchzeri* enhances substrate availability and methane emission in an Arctic wetland. *Soil Biology and Biochemistry*, **45**, 61–70.
- Tagesson T (2011) Land-atmosphere exchange of carbon in a high-Arctic wet tundra ecosystem. Unpublished PhD thesis, Lund University, Meddelanden från Lunds Universitets Geografiska Institutioner, Avhandlingar 192, Lund, 33 pp.
- Tamstorf M, Illeris L, Hansen B, Wisz M (2007) Spectral measures and mixed models as valuable tools for investigating controls on land surface phenology in high arctic Greenland. *BMC Ecology*, **7**, 9.
- Tarnocai C (2006) The effect of climate change on carbon in Canadian peatlands. *Global and Planetary Change*, **53**, 222–232.
- Tarnocai C, Canadell JG, Schuur EAG, Kuhry P, Mazhitova G, Zimov S (2009) Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles*, **23**, GB2023.
- Tokida T, Mizoguchi M, Miyazaki T, Kagemoto A, Nagata O, Hatano R (2007) Episodic release of methane bubbles from peatland during spring thaw. *Chemosphere*, **70**, 165–171.
- Vickers D, Mahrt L (1997) Quality control and flux sampling problems for tower and aircraft data. *Journal of Atmospheric and Oceanic Technology*, **14**, 152–156.
- Walker DA, Reynolds MK, Daniëls FJA *et al.* (2005) The circumpolar Arctic vegetation map. *Journal of Vegetation Science*, **16**, 267–282.
- Webb EK, Pearman GI, Leuning R (1980) Correction of the flux measurements for density effects due to heat and water vapour transfer. *Quarterly Journal of The Royal Meteorological Society*, **106**, 85–100.
- Whalen SC (2005) Biogeochemistry of methane exchange between natural wetlands and the atmosphere. *Environmental Engineering Science*, **22**, 73–94.
- Whalen SC, Reeburgh WS (1988) A methane flux time series for tundra environments. *Global Biogeochemical Cycles*, **2**, 399–409.
- Whiting GJ, Chanton JP (1993) Primary production control of methane emission from wetlands. *Nature*, **364**, 794–795.
- Wickland KP, Striegl RG, Neff JC, Sachs T (2006) Effects of permafrost melting on CO₂ and CH₄ exchange of a poorly drained black spruce lowland. *Journal of Geophysical Research*, **111**, G02011.
- Wieringa J (1993) Representative roughness parameters for homogeneous terrain. *Boundary-Layer Meteorology*, **63**, 323–363.
- Wille C, Kutzbach L, Sachs T, Wagner D, Pfeiffer E-M (2008) Methane emission from Siberian arctic polygonal tundra: eddy covariance measurements and modeling. *Global Change Biology*, **14**, 1395–1408.