

# The NUBASE2016 evaluation of nuclear properties<sup>\*</sup>

G. Audi (欧乔治)<sup>1</sup> F.G. Kondev<sup>2</sup> Meng Wang (王猛)<sup>3,4;1</sup> W.J. Huang(黄文嘉)<sup>1</sup> S. Naimi<sup>5</sup>

<sup>1</sup> CSNSM, Univ Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, 91405 Orsay, France

<sup>2</sup> Argonne National Laboratory, Argonne, IL 60439, USA

<sup>3</sup> Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, People's Republic of China

<sup>4</sup> Joint Department for Nuclear Physics, Institute of Modern Physics, CAS and Lanzhou University, Lanzhou 730000, China

<sup>5</sup> RIKEN Nishina Center, Wako, Saitama 351-0198, Japan

**Abstract:** This paper presents the NUBASE2016 evaluation that contains the recommended values for nuclear and decay properties of 3437 nuclides in their ground and excited isomeric ( $T_{1/2} \geq 100$  ns) states. All nuclides for which any experimental information is known were considered. NUBASE2016 covers all data published by October 2016 in primary (journal articles) and secondary (mainly laboratory reports and conference proceedings) references, together with the corresponding bibliographical information. During the development of NUBASE2016, the data available in the “Evaluated Nuclear Structure Data File” (ENSDF) database were consulted and critically assessed for their validity and completeness. Furthermore, a large amount of new data and some older experimental results that were missing from ENSDF were compiled, evaluated and included in NUBASE2016. The atomic mass values were taken from the “Atomic Mass Evaluation” (AME2016, second and third parts of the present issue). In cases where no experimental data were available for a particular nuclide, trends in the behavior of specific properties in neighboring nuclides (TNN) were examined. This approach allowed to estimate values for a range of properties that are labeled in NUBASE2016 as “non-experimental” (flagged “#”). Evaluation procedures and policies used during the development of this database are presented, together with a detailed table of recommended values and their uncertainties.

AMDC: <http://amdc.impcas.ac.cn/>

**Keywords:** NUBASE2016 evaluation, nuclear properties, Atomic Mass Evaluation (AME), excitation energies of isomers, Evaluated Nuclear Structure Data File (ENSDF)

**PACS:** 21.10.-k, 21.10.Dr, 21.10.Hw, 21.10.Tg **DOI:** 10.1088/1674-1137/41/3/030001

## 1 Introduction

NUBASE is a database containing values of the main nuclear properties, such as masses, excitation energies of isomers, half-lives, spins and parities, and decay modes and their intensities, for all known nuclides in their ground and excited isomeric states. The information presented in NUBASE represents the fundamental building blocks of modern nuclear physics, and specifically, of nuclear structure and nuclear astrophysics research. The first version of NUBASE was published in 1997 [1] and since then it has been widely used in many fields from fundamental physics to applied nuclear sciences. The present publication includes updated information of all nuclear properties given in the previous publications of NUBASE [1–3].

One of the main applications of NUBASE2016 is the “Atomic Mass Evaluation” (AME2016, second and third parts of this issue) where it is imperative to have an unambiguous identification of all states involved in a particular de-

decay, reaction or mass-spectrometric measurement. This is the primary reason for which the two evaluations are published jointly in the present issue, for the third time since the publication of the NUBASE2003 [2].

Furthermore, with the advances of modern mass-spectrometry techniques (see for example the special issue of “one hundred years of mass spectrometry” for relevant topics [4]) and the availability of intense stable and rare-isotope beams, a large number of unstable nuclei can be produced in a single experiment in their ground and/or isomeric states, and their masses can be measured with high precision. Thus, NUBASE2016 can be particularly useful in future mass measurements, where an unambiguous identification of complex mass-spectrometric data would be required.

Applications of this database in astrophysics network calculations and in theoretical studies of nuclear properties, where complete and reliable data for all known nuclei are needed, are also envisioned.

Received 10 March 2017

\* This work has been undertaken with the endorsement of the IUPAP Commission on Symbols, Units, Nomenclature, Atomic Masses and Fundamental Constants (SUNAMCO).

1) E-mail: wangm@impcas.ac.cn

©2017 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

Last, but not least, the evaluated data presented in NUBASE2016 are also useful for specialists in applied nuclear fields, such as reactor engineering and design, fuel manufacture and transport, waste management, material analysis, medical diagnostics and radiotherapy, and others, where one needs to access basic information for a given nuclide.

The information presented in NUBASE2016 fulfills several user-demanded requirements, namely that it is: a) *complete* – includes all measured quantities and their uncertainties, b) *up-to-date* – results from the most recent publications are included, c) *credible and reliable* – identifies and resolves contradictory results that exist in the scientific literature, as well as in other nuclear physics databases, d) *properly referenced* – provides comprehensive information for the traceability of all included data.

Most of the data included in NUBASE2016 are in principle available in two other evaluated databases: the “Evaluated Nuclear Structure Data File” (ENSDF) [5] and the “Atomic Mass Evaluation” (AME2016). Therefore, the demand for NUBASE could be partially fulfilled by combining these two databases into a single, ‘horizontal’ structure, which exists in AME, but not in ENSDF. Therefore, NUBASE2016 can be considered at a first level as a critical combination of those two evaluations.

During the development of the present version of NUBASE, it was imperative to examine all available literature for several nuclides in order to revise results adopted in ENSDF, and to ensure that the recommended data are presented in a consistent way (*credibility and reliability* requirements). It was also necessary to include all the available experimental data, i.e. not only results that were published recently (*up-to-date* requirement), but also older data that were missing in ENSDF (*completeness* requirement). This implied that some extra evaluation work was necessary. The corresponding conclusions are added as remarks in the NUBASE2016 table, and in the discussions below. Complete bibliographical references are given for all added experimental data in Table I (see Section 2.8).

There is no strict literature cut-off date for the results presented in the NUBASE2016 evaluation: all data available to the authors until October 2016 have been included. Results that were not incorporated for special reasons, e.g. the need for a heavy revision of the evaluation at too late a stage of development, are added, whenever possible, in remarks to the relevant data.

During the preparation of NUBASE2016, we noticed that Ref. [6] reports important decay data for proton-rich nuclides  $^{67}\text{Kr}$ ,  $^{63}\text{Se}$  and  $^{59}\text{Ge}$ , where a two-proton emission from  $^{67}\text{Kr}$  was observed. We found that it was too heavy an effort at this stage to include these results into NUBASE2016, especially to establish the atomic mass surface in this region. They are not included in the current evaluation, but the original paper

is just mentioned here.

The contents of NUBASE2016 are described below, together with the adopted policies that were used during the development of this database. Section 3 presents the updating procedures, while the electronic distribution and interactive display of NUBASE2016 contents by means of a stand-alone PC-program are described in Section 4.

## 2 Contents of NUBASE2016

The NUBASE2016 evaluation contains recommended values for the basic nuclear ground-state properties, for 3437 nuclides, derived from all available experimental results, together with some values estimated by extrapolating neighboring ones. It also contains data for 1318 nuclides which have one or more excited isomeric states with half-lives longer than 100 ns.

Similar to the previous editions, NUBASE2016 also contains data on 186 isobaric analog states (IAS), which have their excitation energies determined either through an “internal relation” and taken from ENSDF, or through an “external relation” and then determined by the AME2016 evaluation.

For each nuclide ( $A, Z$ ), and for each state (ground or excited isomer), the following properties were compiled and, when necessary, evaluated: mass excess, excitation energy of excited isomeric states, half-life, spin and parity, decay modes and their intensities, isotopic abundance (for a stable nuclide), year of discovery and the corresponding bibliographical information for all experimental data.

References to published articles in the description sections below are given by means of the keynumber style used in the “Nuclear Science Reference” (NSR) bibliographical database [7]. However, references quoted in the NUBASE2016 tables are abbreviated with the first two digits of the year of publication being omitted from the NSR keynumbers. The complete reference list is given at the end of this issue, together with the references used in AME (see AME2016, Part II).

At the time the work on NUBASE2016 was completed, superheavy elements (SHE) up to  $Z = 118$  were officially named by The Commission on Nomenclature of Inorganic Chemistry of the International Union of Pure and Applied Chemistry (IUPAC) [8]:

113	Nihonium	(Nh),
115	Moscovium	(Mc),
117	Tennesine	(Ts),
118	Oganesson	(Og).

We were not able to include the new names in AME2016 and NUBASE2016, but instead we used the provisional symbols Ed, Ef, Eh, and Ei for elements 113, 115, 117, and 118, respectively.




Figure 1. Chart of the nuclides displaying the accuracy 'u' of masses (created by NUCLEUS-AMDC).

NUBASE2016 contains numerical and bibliographical data for all known nuclides for which at least one property is known experimentally in their ground state, excited isomeric states with  $T_{1/2} \geq 100$  ns, and/or IAS. It also includes information on yet unobserved nuclides, estimated from the observed experimental trends of neighboring nuclides (TNN). This ensures continuity in the set of considered nuclides simultaneously in  $N$ , in  $Z$ , in  $A$  and in  $N - Z$ . The chart of nuclides defined in this way has a smooth contour.

For experimentally unknown properties, values were also estimated from TNN. Similarly to AME2016, the estimated values are flagged with the symbol ‘#’ to indicate non-experimental information.

As a rule, one standard deviations ( $1\sigma$ ) are used in NUBASE2016 to represent the uncertainties associated with the quoted experimental values. Unfortunately, authors of research articles do not always define the meaning of their reported uncertainties and those values were assumed to be one standard deviations. In many cases, uncertainties are not even given at all and were estimated by us, considering the limitations of the experimental method.

Values and corresponding uncertainties for properties given in NUBASE2016 are rounded, even if unrounded values were given in the literature or in ENSDF. In cases where the two furthest left significant digits in the uncertainty were larger than a given limit (set to 30 for masses and energies to be consistent with AME2016, and set to 25 for all other quantities, as used in ENSDF), values and uncertainties were rounded accordingly (see examples in the ‘Explanation of table’). In a few cases that were deemed essential for traceability purposes (e.g. isotopic abundances) the original (unrounded) value is also provided in an associated comment.

## 2.1 Mass excess

In NUBASE2016 the mass excess values (in keV), defined as being differences between the atomic mass (in mass units) and the mass number, together with their one-standard-deviation uncertainty, are taken from the mass tables of the AME2016 evaluation.

In general, knowledge of masses can provide valuable information on decay modes, in particular for a particle-decay instability, or  $\beta$ -delayed particle-decay, for nuclei far from the line of stability. Such information is used in NUBASE2012, for example for  $^{10}\text{He}$ ,  $^{39}\text{Sc}$ ,  $^{62}\text{As}$ , or  $^{63}\text{As}$ . In some cases, the claimed observations of decay modes were rejected when it was found that they were not allowed through simple energetics.

Figure 1 displays the mass accuracy from the main table, as a function of  $N$  and  $Z$ .

## 2.2 Isomers

In the first version of NUBASE [1], a definition for excited isomers was adopted: excited states with a half-life longer than one millisecond. Within this definition, all  $\beta$ -

decaying states were included in this category, since they have a lower half-life limit of one millisecond. However, already at that time, it was noticed that such a definition had several drawbacks, particularly for neutron-deficient alpha- and proton-decaying nuclides, where much shorter-lived states were known to exist. Moreover, several cases are known where isomers with half-lives far below one millisecond survive longer than the ground state itself, e.g.  $^{216}\text{Fr}$ .

With the publication of NUBASE2003 [2], the definition of isomers was extended to half-lives longer than 100 ns, and such states are now included in NUBASE2016. The main reasons for this change were to include:

- all proton- and alpha-decaying states observed in many neutron-deficient nuclei,
- isomers that may be detected in mass-spectrometric experiments performed at accelerator facilities following the immediate detection of the produced nuclei, and
- all possible isomers that may be detected in such experiments in the future.

In NUBASE2016, isomers are tabulated in order of increasing excitation energy and identified by appending the letters ‘ $m$ ’, ‘ $n$ ’, ‘ $p$ ’, ‘ $q$ ’, or ‘ $r$ ’ to the nuclide name, e.g.  $^{90}\text{Nb}$  for the ground state,  $^{90}\text{Nb}^m$  for the first excited isomer,  $^{90}\text{Nb}^n$  for the second one, and  $^{90}\text{Nb}^p$ ,  $^{90}\text{Nb}^q$ , and  $^{90}\text{Nb}^r$  for the third, fourth and fifth ones, respectively. In the cases of  $^{179}\text{Ta}$  and  $^{214}\text{Ra}$  a sixth isomer had to be included, and they were labeled provisionally with the letter ‘ $x$ ’.

Suffix ‘ $x$ ’ also applies to mixtures of levels which are used in the atomic mass evaluation. These mixtures occur in spallation reactions or in fission and they appear in mass measurements performed using mass spectrometers. For each mixture, the excitation energy and the relative production rate of isomeric state with respect to ground state are given.

The excitation energy of a given isomer can be determined using different experimental methods, which, in general, belong to the category of either internal or external relations. A typical internal relation is via the  $\gamma$ -ray decay energy, or a combination of such  $\gamma$ -ray energies. The most accurate values for the excitation energies of isomers deduced by this approach can be found in ENSDF, where a least-squares fitting procedure is applied to all  $\gamma$  rays along the decay path of a particular isomer. However, when no such internal relations can be established, then the relation to other nuclides (external relations) can be used to deduce the mass (or energy) difference between excited and ground-state isomers. In all such cases, the most accurate values can only be derived using the AME evaluation procedure and the values are therefore taken from AME2016. The origin (the method used to establish the external relation) of each isomer data element is then indicated by a two-letter code, next to the isomer excitation energy, in the NUBASE2016 table. For internal relations, the origin field is left blank and the numerical values are taken either from ENSDF or from literature updates. In the latter case, a least-squares fit to the measured  $\gamma$ -ray decay energies from

complex level schemes was applied, in accordance with the current ENSDF policies.

It also happens that connections between excited and ground state isomers can be obtained by both internal relations and one, or more, external relations with comparable accuracies. All relations are then combined within the AME2016 data by adding an equation that relates the excitation energy obtained from ENSDF (or from literature), so that AME2016 derives the best combination of all data. For example, the AME2016 derives the mass of  $^{178}\text{Lu}^m$  at 66% from  $E_x(\text{IT})=120(3)$  keV [1993Bu02] and at 34% from  $^{176}\text{Lu}(\text{t,p})^{178}\text{Lu}^m=4482(5)$  keV [1981Gi01]. The adjusted excitation energy is thus 123.8(2.6) keV.

In some cases, excitation energies known from internal relations are essential in order to determine the mass of the ground state. Those values are labeled in the NUBASE table with ‘IT’ in the origin field. They are entered as an equation in AME2016 so that the ground state mass can be derived. For example, the mass of  $^{62}\text{Mn}$  was listed as unknown in AME2012, since it was the excited isomer that was measured in a Penning trap experiment [2012Na15]. However, the excitation energy of  $^{62}\text{Mn}^m$  was determined recently via  $\gamma$ -ray spectroscopy [2015Ga38], so the mass of the ground state is established experimentally. An interesting case is the mass and excitation energy of  $^{186}\text{Tl}^n$ , where its mass is experimentally known from a Penning trap (ISOLTRAP) measurement [2014Bo26]. The well known transition from  $^{186}\text{Tl}^n$  to  $^{186}\text{Tl}^m$  allows to determine not only the mass of the latter, but also the excitation energy of the  $\alpha$ -decaying isomers in the parent nuclides  $^{190}\text{Bi}^m$ ,  $^{194}\text{At}^m$  and  $^{198}\text{Fr}^m$ .

When the existence of an isomer is ambiguous, it is flagged with ‘EU’ (“existence uncertain”) in the origin field (e.g.  $^{73}\text{Zn}^n$ ). A comment is generally added to indicate why this existence is questioned, or where this matter is discussed in more detail. Five isomers, namely  $^{73}\text{Zn}^n$ ,  $^{138}\text{Pm}^n$ ,  $^{141}\text{Tb}^m$ ,  $^{185}\text{Bi}^n$ ,  $^{273}\text{Ds}^m$  are treated in this way in the present evaluation and the mass excess and excitation energy values are given for them all except  $^{138}\text{Pm}^n$ , for which the existence is strongly doubted.

When a particular isomer was initially reported as “discovered”, but later it was proved to be an error, it is flagged with ‘RN’ in the origin field, indicating “reported, non-existent”. Three isomers, namely  $^{117}\text{La}^m$ ,  $^{156}\text{Tm}^n$  and  $^{181}\text{Pb}^m$  are treated in this way. In these cases, no mass-excess or excitation energy values are given, and, similarly to the ‘EU’ choice above, a “non-existent” label is added.

*Note:* the use of the two flags, ‘EU’ and ‘RN’, was extended to cases where the discovery of a nuclide is questioned (e.g.  $^{260}\text{Fm}$  or  $^{289}\text{Lv}$ ). However, an estimate for the ground state mass, derived from trends in the mass surface (TMS), is always given in AME2016 and NUBASE2016.

In several instances, lower and higher limits for the excitation energy of a particular isomer are presented in ENSDF. The policy of NUBASE2016 is that a uniform distribution of

probabilities is assumed, which yields a mid-range value and a  $1\sigma$  uncertainty corresponding to 29% of the range (see Appendix B of the AME2016, Part I in this issue for a complete description of this procedure). For example, the excitation energy of the  $^{162}\text{Tm}^m$  isomer is known from ENSDF to be above the 66.90 keV level. On the other hand, there is also solid experimental evidence that it is below the 192 keV level, and so this information is presented as  $E_x = 130(40)$  keV in NUBASE2016. However, if such a value is based on theoretical considerations, or from TNN, the resulting  $E_x$  is considered as a non-experimental quantity and the value is consequently flagged with the ‘#’ symbol.

In cases where the uncertainty of the excitation energy,  $\sigma$ , is relatively large as compared to the  $E_x$  value, the assignment of the level as a ground or isomeric state is uncertain. If  $\sigma > E_x/2$ , a ‘\*’ flag is added in the NUBASE2016 table.

The ordering of several ground and excited isomeric states were reversed as compared to the recommendations in ENSDF. These cases are flagged with the ‘&’ symbol in the NUBASE2016 table. In several other instances, evidence was found for states located below the adopted ground state in ENSDF. There are also cases where the trends in neighboring nuclides, with the same parities in  $N$  and  $Z$ , strongly suggest that such a lower state should exist. Such results were added in the NUBASE tables and can be easily located, as they are flagged with the ‘&’ symbol. In a growing number of cases, new experimental information on masses led to a reversal of the ordering between previously assigned ground and excited isomeric states. Thanks to the coupling of the NUBASE2016 and AME2016 evaluations, all changes in the ordering of nuclear levels have been carefully synchronized.

Finally, there are cases where data exist on the ordering in energy of the isomers, e.g. if one of them is known to decay into the other one, or if the Gallagher-Moszkowski rule [9] points strongly to one of the two as being the ground-state. Detailed discussions can be found in Ref. [10].

### 2.2.1 Isobaric analog states (IAS)

In the previous version of NUBASE [3] we have included the  $T = 3/2$  to  $T = 3$  experimentally observed (IAS). These states are also included in NUBASE2016 and generally labelled with  $i$  or  $j$  superscripts, for members of successively higher multiplets. The experimental information about IAS has been evaluated in more detail recently in Ref. [11]. Some nuclides belong simultaneously to several categories, for example, they may be in their ground state but they may also be IAS of some other ground state nucleus, as is the general case for ground state mirror nuclei. Here, the IAS label is not present, since these nuclides are already naturally included in the database. Another exception is the set of  $N = Z$ ,  $T = 1$  odd-odd ground state nuclides which are also already part of the original dataset of ground state masses. They are:  $^{34}\text{Cl}_{17}$ ,  $^{42}\text{Sc}_{21}$ ,  $^{46}\text{V}_{23}$ ,  $^{50}\text{Mn}_{25}$ ,  $^{54}\text{Co}_{27}$ ,  $^{62}\text{Ga}_{31}$  and  $^{70}\text{Br}_{35}$ . The reader may note that the  $Z = 29$  and  $Z = 33$  nuclides are not included in this series, since their ground states are  $T = 0$ , as

expected from theory. Finally, there are eight excited isomers,  $^{16}\text{N}^m$ ,  $^{26}\text{Al}^m$ ,  $^{34}\text{Cl}^m$ ,  $^{38}\text{K}^m$ ,  $^{46}\text{V}^m$ ,  $^{50}\text{Mn}^m$ ,  $^{54}\text{Co}^m$  and  $^{72}\text{Ga}^m$ , which are also IAS. In such cases, the isomer labels ('m', 'n', ...) are used preferentially over the IAS labels. Here we note with interest that five of them have experimentally determined excitation energies, at least partly, by the JYFLTRAP-Jyväskylä Penning trap.

In NUBASE2016 there are roughly 181 unique IAS masses, of which 113 are evaluated in the AME via external relations, and 68 cases evaluated through internal relations and published in ENSDF. There are five cases where no clear experimental data is available, and although some Isobaric Multiplet Mass Equation (IMME) [12] and Coulomb Displacement Energy (CDE) [13] calculations point to a likely IAS state, their existence cannot yet be certified experimentally (for example  $^{15}\text{O}^i$ ).

The isospin multiplet assignment given in the table is the logical IAS multiplet value, and has not necessarily been deduced experimentally.

### 2.3 Half-life

Fig. 2 displays the half-lives of nuclides in NUBASE2016. In the light mass region, nuclides beyond the particle drip-lines can be studied with modern radioactive ion facilities. Most of these unbound nuclides exist for a very short time before they directly decay via particle emission. For some of them, such as  $^{19}\text{Mg}$  and  $^{26}\text{O}$ , the half-lives can be determined experimentally with novel experimental methods. For most unbound nuclei, only the total level width ( $\Gamma_{\text{cm}}$ ) can be measured and therefore the half-life ( $T_{1/2}$ ) can be deduced using the equation  $\Gamma_{\text{cm}} T_{1/2} \simeq \hbar \times \ln 2$  so that

$$T_{1/2} (\text{s}) \simeq 4.562 \cdot 10^{-22} / \Gamma_{\text{cm}} (\text{MeV}).$$

The following units are used for convenient display in NUBASE2016: seconds (s) and its sub-units, minutes (m), hours (h), days (d) and years (y) and its sub-units. Conversion between years and seconds or days could follow various definitions: Julian year, Gregorian year, tropical year 1900, epoch 2000, etc., differing only slightly from each other. A fixed value of:

$$1 \text{ y} = 31\,556\,926 \text{ s} \quad \text{or}$$

$$1 \text{ y} = 365.2422 \text{ d}$$

was adopted in NUBASE2016.

Asymmetric uncertainties for half-lives,  $T_{1/2}^{+a}_{-b}$ , are often presented in the literature. However, for these values to be used in practical applications, they need to be symmetrized. A rough symmetrization procedure was used earlier (see AME1995) where the central value was taken as the mid-value between the upper and lower  $1\sigma$ -equivalent limits,  $T_{1/2} + (a - b)/2$ , and the uncertainty was defined to be the average of the two uncertainties,  $(a + b)/2$ . A strict statistical

derivation (see Appendix A) shows that a better approximation for the central value can be obtained by using

$$T_{1/2} + 0.64 \times (a - b).$$

The exact expression for asymmetric uncertainties, adopted in NUBASE2016, is presented in Appendix A.

When two or more independent measurements were reported in the literature, the corresponding values were weighted by their reported precisions and then averaged. While doing this, the NORMALIZED CHI,  $\chi_n$  (or 'consistency factor' or 'Birge ratio'), as defined in AME2016, is considered. When  $\chi_n$  is larger than 2.5, departure from the statistical result is allowed and the external uncertainty for the average result is adopted. This follows the same policy that is discussed and adopted in AME2016. Very rarely, when  $\chi_n$  is so large that all individual uncertainties can be considered as irrelevant, the arithmetic (unweighted) average is adopted and the corresponding uncertainty is based on the dispersion of the values. In such cases, the list of values that were averaged, together with the  $\chi_n$  value (when relevant) and the reason for this choice, are given in the NUBASE2016 table. When contradictory (conflicting) results were identified in the literature, attention was focused on establishing the reason for such discrepancies, and consequently, any bad data were rejected. The justification for such decisions are given as comments in the NUBASE2016 table.

In experiments where extremely rare events are detected and where the results are very asymmetric (e.g. studies of super-heavy nuclei), the half-life values reported in different publications were not directly averaged. Instead, when the information presented in the literature was sufficient (e.g.  $^{264}\text{Hs}$ ), the decay times associated with the individual events were combined, as prescribed by Schmidt *et al.* [1984Sc13].

Some experimental results are reported in the literature as a range of values with a most probable lower and upper limit. These are treated, as in the case of isomer excitation energies (see preceding page), as a uniform distribution of probabilities.

In the NUBASE2016 table, an upper or lower limit on the half-life value is given for nuclides identified using a time-of-flight technique. The following policies were considered:

i) For *observed* nuclides, the lower limit for the half-life is given in place of the uncertainty (e.g.  $^{44}\text{Si}$ ). However, such limits should be used with caution, since they may be far below the actual half-life. In order to avoid confusion, a somewhat more realistic estimate (flagged with #), derived using TNN is also given. ii) For nuclides that were sought, but *not observed*, the upper limit is given in place of the actual half-life uncertainty. Upper limits for a dozen undetected nuclides were evaluated by F. Pougheon [1993Po.A], based on the time-of-flight of the experimental setup and the production yields expected from TNN (e.g.  $^{21}\text{Al}$ ).




Figure 2. Chart of the nuclides displaying half-lives (created by NUCLEUS-AMDC).

When ground-state half-lives for nuclides with the same parities in  $Z$  and  $N$  are found to vary smoothly, interpolation or extrapolation (TNN) is used to obtain reasonable estimates for unknown cases.

The super-allowed  $0^+ \rightarrow 0^+$  nuclear  $\beta$  decays between isospin analog states with isospin  $T=1$  and spin-parity  $J^\pi=0^+$  are of particular interest due to their pivotal role in the precise determination of  $V_{ud}$  to test the unitarity of the Cabibbo - Kobayashi - Maskawa (CKM) Matrix. The evaluation of super-allowed decays, including their half-lives, is a long-standing work carried out by J.C Hardy and I.S. Towner. In the most recent survey [14], experimental data of 20 super-allowed transitions have been compiled and carefully evaluated. Half-lives of these nuclides are compared in Fig. 3. It can be seen clearly that the values listed in NUBASE2016 agree well with the values from Ref. [14]. The only significant differences occur for  $^{18}\text{Ne}$  and  $^{42}\text{Ti}$ , for which new experimental results were published after the publication of Ref. [14].




Figure 3. Comparison of  $T_{1/2}$  for 20 super-allowed  $\beta$  emitters from NUBASE2016 (N16) and Ref. [14] (HT). The error bars at the points display the uncertainties from Ref.[14], and the shaded area displays the uncertainties in NUBASE2016.

## 2.4 Spin and parity

As for ENSDF, spin and parity values are presented with and without parentheses, based on strong and weak assignment arguments, respectively (see the introductory pages of Ref. [15]). Unfortunately, parentheses in ENSDF are also applied to estimates from theory or from TNN. In NUBASE2016, following our policy of making a clear distinction between experimental and non-experimental information, parentheses are used if the so-called “weak” argument is based on experimental observations, while the symbol ‘#’ is used for the other cases. It should also be noted that despite the well-defined evaluation policies [15], there are a number of inconsistencies in ENSDF regarding the spins

and parities for nuclear states. Often, the proposed assignments reflect the interpretation of a particular ENSDF evaluator, rather than that of firm policy rules. As a result, assignments to similar states in neighboring nuclides are put in parenthesis by one evaluator, but not by another, although similar experimental information is available.

We have tried to use a consistent approach in assigning spins and parities to nuclear states, but the survey is still far from complete and the reader may still find inconsistencies. The authors would gratefully appreciate feedback from users for such cases, to improve future versions of NUBASE.

If spins and parities are not determined experimentally, they can be estimated from TNN with the same parities in  $N$  and  $Z$ . Although, this is frequently the case for odd- $A$  nuclides, such trends are also sometimes valid for odd-odd nuclides, especially in the neighborhood of magic numbers. In all cases, the estimated values are flagged with the ‘#’ symbol.

The review of nuclear radii, moments and spins by Otten [1989Ot.A], as well as the recent compilation by MacDonald [16], were used to check and complete the spin values in NUBASE2016.

The spins and parities of odd-even, even-odd, odd-odd nuclides in their ground states are displayed in Fig. 4, Fig. 5 and Fig. 6, respectively.

## 2.5 Decay modes and their intensities

Fig. 7 displays the main decay modes of all known nuclides. The most important policy in assembling the information for the decay modes was to establish a clear distinction between a decay mode that is energetically allowed, but not yet experimentally observed (represented by a question mark alone, which refers to the decay mode itself), and a decay mode which is actually observed, but for which the intensity could not be determined (represented by ‘=?’, the question mark referring here to the quantity after the equal sign).

As in ENSDF, no corrections were made to normalize the primary intensities to 100%.

In addition to applying direct updates from the literature, partial evaluations completed by other authors were also considered and properly referenced. Those cases are mentioned below when discussing some particular decay modes.

### $\beta^+$ decay

In the NUBASE evaluations some definitions and notations for  $\beta^+$  decay were refined to provide a clearer presentation of the available information. Specifically,  $\beta^+$  denotes the decay process that includes both electron capture, labeled  $\epsilon$ , and decay by positron emission, labeled  $e^+$ . One can then symbolically write:  $\beta^+ = \epsilon + e^+$ . It is well known that for an available energy below 1022 keV, only electron capture,  $\epsilon$ , is allowed, while above that value the two processes are in competition.

*Remark:* this notation is **not** the same as the one used implicitly in ENSDF, where the combination of both modes is denoted “ $\epsilon + \beta^+$ ”.



When the two modes compete, the separated intensities are not always experimentally available and frequently they are deduced from model calculations, as is the policy in ENSDF. In continuation of one of the general NUBASE policies, in which only experimental information is used whenever possible, it was decided not to retain the separated values calculated in ENSDF (which are scarce and not always updated). Only in a few very specific cases, where the distinction is of importance, such as rare processes ( $^{91}\text{Nb}$ ,  $^{54}\text{Mn}$ ,  $^{119}\text{Te}^m$ ), separate values are given.

By the same token, both electron-capture-delayed fission ( $\epsilon\text{SF}$ ) and positron-delayed fission ( $e^+\text{SF}$ ) are given with the same symbol  $\beta^+\text{SF}$ .

### Double- $\beta$ decay

In the course of this work it was found that half-lives for double- $\beta$ -decaying nuclides were not always consistently given in ENSDF. Since the two-neutrino gs-gs transition is the dominant decay process (one exception may be  $^{98}\text{Mo}$ , for which the neutrinoless decay is predicted to be faster, see [2002Tr04]), only those half-life values or their upper-limits were presented in the NUBASE2016 table. No attempt was made to convert the upper limit results given by different authors to the same statistical confidence level (CL).

The excellent compilation of Tretyak and Zdesenko [2002Tr04] was of great help in evaluating such decays.

### $\beta$ -delayed particle decays

For delayed particle decays, intensity relations must be carefully considered. By definition, the intensity of a decay mode is the percentage of decaying parent nuclei in that mode. But traditionally, the intensities of the pure  $\beta$  decay are summed with those of the delayed particles in order to give an intensity that is assigned to the pure  $\beta$  decay. For example, if the  $(A, Z)$  nuclide has a decay described traditionally by ' $\beta^-=100$ ;  $\beta^-n=20$ ', this means that for 100 decays of the parent, 80  $(A, Z+1)$  and 20  $(A-1, Z+1)$  daughter nuclei are produced and that 100 electrons and 20 delayed neutrons are emitted. A strict notation in this case, using the definition above, would be ' $\beta^-=80$ ;  $\beta^-n=20$ '. However, in the present work, it has been decided to follow the above traditional notation.

This also holds for more complex delayed emissions. For example, a decay described by: ' $\beta^-=100$ ;  $\beta^-n=30$ ;  $\beta^-2n=20$ ;  $\beta^-\alpha=10$ ' corresponds to the emission of 100 electrons,  $(30+2\times 20=70)$  delayed-neutrons and 10 delayed- $\alpha$  particles; and in terms of residual nuclides, to 40  $(A, Z+1)$ , 30  $(A-1, Z+1)$ , 20  $(A-2, Z+1)$  and 10  $(A-4, Z-1)$ . More generally, the number of emitted neutrons per 100 decays,  $P_n$ , can be written as:

$$P_n = \sum_i i \times \beta_{in}^-;$$

and similar expressions can be written for  $\alpha$  and proton emis-

sion. The number of residual daughter nuclides  $(A, Z+1)$  populated via  $\beta^-$  decay is then:

$$\beta^- - \sum_i \beta_{in}^- - \sum_j \beta_{j\alpha}^- - \dots$$

Another special remark concerns the intensity of a particular  $\beta$ -delayed mode. In general, the primary (parent)  $\beta$  decay populates several excited states in the daughter nuclide, which can further decay by particle emission. However, in a case where the ground state of the daughter nuclide decays also by the same particle emission, some authors included its decay in the value for the corresponding  $\beta$ -delayed intensity. It has been decided to not use such an approach in NUBASE2016 for two main reasons. Firstly, the energies of delayed particles emitted from excited states are generally much higher than those emitted from the ground state, implying different subsequent processes. Secondly, the characteristic decay times from excited states are related to the parent, whereas decays from the daughter's ground state are connected to the daughter nuclide itself. For example,  $^9\text{C}$  decays via  $\beta^+$  with an intensity of 100% of which 12% and 11% populate two excited proton-emitting states in  $^9\text{B}$ , and 17% goes to an  $\alpha$ -emitting state. Thus,  $\beta^+p=23\%$  and  $\beta^+\alpha=17\%$ , from which the user of the NUBASE2016 table can derive a 60% direct feeding of the ground state of  $^9\text{B}$ . In a slightly different example,  $^8\text{B}$  decays to only two excited states in  $^8\text{Be}$ , which in turn decay by  $\alpha$ - and  $\gamma$ -ray emissions, but not to the  $^8\text{Be}$  ground state. Thus, one may write  $\beta^+=100\%$  and  $\beta^+\alpha=100\%$ , the difference of which leaves 0% for the feeding of the daughter's ground state.

Finally, the users should be aware that the percentages given in the NUBASE2016 table are related to 100 parent decaying nuclei, rather than to the primary beta-decay fraction. An illustrative example is given by the decay of  $^{228}\text{Np}$ , for which the delayed-fission probability is given in the original paper as 0.020(9)% [1994Kr13], but this value is relative to the  $\epsilon$  process, which has an intensity of 60(7)%. Thus, the renormalized delayed-fission intensity is  $0.020(9)\% \times 0.60(7) = 0.012(6)\%$  of the total decay intensity.

In compiling the data for  $\beta^+$ -delayed proton and  $\alpha$  activities, the remarkable work of Hardy and Hagberg [1989Ha.A], in which the corresponding physics was reviewed and discussed in detail, was consulted. The review of Honkanen, Äystö and Eskola [17] on delayed proton decays has also been consulted.

Similarly, the review of delayed neutron emission by Hansen and Jonson [18] was carefully examined and used in the NUBASE tables, together with the evaluation of Rudstam, Aleklett and Sihver [1993Ru01].

## 2.6 Isotopic abundances

Isotopic abundances are taken from the compilation of M. Berglund and M.E. Wieser [2011Be53] and the values are listed in the decay field with the symbol  $IS$ . These data




Figure 4. Chart of the nuclides displaying spins and parities. Only the odd-Z even-N nuclides are shown (created by NUCLEUS-AMDC).




Figure 5. Chart of the nuclides displaying spins and parities. Only the even- $Z$  odd- $N$  nuclides are shown (created by NUCLEUS-AMDC).




Figure 6. Chart of the nuclides displaying spins and parities. Only the odd-Z odd-N nuclides are shown (created by NUCLEUS-AMDC).




Figure 7. Chart of the nuclides displaying decay modes (created by NUCLEUS-AMDC).




Figure 8. Chart of the nuclides displaying the years of discovery (created by NUCLEUS-AMDC).

are given in the NUBASE tables as presented originally in [2011Be53], and so in this case the rounding policy was not applied.

## 2.7 Year of discovery

As in NUBASE2012, the present tables include information of the year of discovery for each nuclide in its ground or isomeric state. For the former, recent evaluations performed by a group at Michigan State University [19] were adopted. Similar criteria was used when assigning the year of discovery for isomeric states. The information about the year of discovery is illustrated in Fig. 8.

## 2.8 References

The year of the archival file for the nuclides evaluated in ENSDF is indicated, otherwise this entry is left blank.

References for all of the experimental updates are given by the NSR keynumber style [7], and are listed at the end of this issue. They are followed by one, two or three one-letter codes which specify the added or modified physical quantities. In cases where more than one reference is needed to describe a particular update, they are given as a remark. No reference is given for estimated values. The initials of the former and present evaluators, AHW, FGK, GAU, HWJ, JBL, MMC, WGM, XUX, are used as reference keys in cases where it may not be precisely clear that the re-interpretation of data were made by the authors.

## 3 Updating procedure

In general, NUBASE was updated via two routes: from ENSDF after each new *A*-chain evaluation is published (or from the bi-annual releases), and directly from the literature. Data available in the “eXperimental Unevaluated Nuclear Data List” (XUNDL)[20] database were also regularly consulted.

ENSDF files are retrieved from NNDC using the on-line service [5]. Computer programs, originally developed by O. Bersillon and J. Blachot [21], were used to successively:

- check that each *Z* in the *A*-chain has an ‘adopted levels’ data set; if not, a corresponding data set is generated from the ‘decay’ or ‘reaction’ data set,
- extract the ‘adopted levels’ data sets from ENSDF,
- extract the required physical quantities from these data sets, and convert them into the NUBASE format.

The processed data were used to manually update the previous version of NUBASE.

ENSDF is updated generally by *A*-chains and more recently also by individual nuclides. Its contents are extensive, since it encompasses all of the complex nuclear structure and decay properties. This is a huge effort, and it is not surprising that occasionally some older data (in particular annual reports, conference proceedings, and theses) are missing and that some recent data have not yet been included. When such

cases were revealed, they were analyzed and evaluated, as described above, and the NUBASE2016 database was updated accordingly. In principle, these new data will be included in future ENSDF evaluations and the corresponding references can then be removed from future NUBASE distributions. Unfortunately, it has been observed in the past that such a procedure was not always adhered to. In fact, in some newer ENSDF files, quotations to earlier NUBASE publications were found, which leads to an undesirable loop resulting in non-traceable information. For this reason, in such cases the original references are repeated here again.

## 4 Distribution and displays of NUBASE2016

The full contents of the present evaluation is available on-line at the Atomic Mass Data Center (AMDC) website [22], as well as at a mirror website maintained by the International Atomic Energy Agency (IAEA) [23]. An electronic ASCII file for the NUBASE2016 table is also distributed at the AMDC website. Any work that uses those files should make reference to the present publication and not to the electronic files.

The contents of NUBASE2016 can be displayed by the stand-alone PC-program called “NUCLEUS”. The charts of nuclides shown in this paper were created by using this program. The program “NUCLEUS” has been updated according to the present NUBASE2016 evaluation and can be downloaded from the AMDC website [22] and the IAEA [23].

## 5 Conclusions

The ‘horizontal’ evaluated database, NUBASE2016, which contains the recommended values for the main properties of all known nuclides in their ground and excited isomeric states, has been updated. These data originate from the intersection of two evaluated databases: ENSDF, followed by a critical assessment of the validity and completeness of those data, including new updates from the literature, and AME2016. The main requirement in developing NUBASE2016 was to cover as completely as possible all available experimental data and to provide proper references to them, especially for cases that are not already included in ENSDF. This traceability allows any user to check the recommended data and, if necessary, to undertake a re-evaluation.

As a result of this ‘horizontal’ work, better homogeneity in handling and presentation of all data was obtained for all known nuclides. Furthermore, isomeric assignments were examined more critically and the data of their excitation energies were improved.

## 6 Acknowledgments

We wish to thank many colleagues who answered our questions about their experiments and those who sent us preprints of their papers. We appreciate the help provided by

J.K. Tuli and B. Singh in solving some of the puzzles we encountered in ENSDF. Continuous interest, discussions, suggestions and encouragements from D. Lunney, Zhang Yuhu and Furong Xu were highly appreciated.

This work is supported in part by the National Key Program for S&T Research and Development (Contract No. 2016YFA0400504) and the Major State Basic Research Development Program of China (Contract No. 2013CB834401). The work at ANL was supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357. W.J. Huang acknowledges the support from the China Scholarship Council, grant No. 201404910496. S. Naimi acknowledges the support of “RIKEN Pioneering Project Funding” from the Riken project.

**Appendix A Symmetrization of asymmetric uncertainties**

Experimental data are sometimes given with asymmetric uncertainties,  $X_{-b}^{+a}$ . If these data are to be used in some practical applications, their uncertainties may need to be symmetrized. A simple method (Method 1) that was developed earlier, uses the central value to be in the middle between the upper and lower  $1\sigma$ -equivalent limits

$$X + (a - b)/2, \text{ with the uncertainty defined to be the average of the two uncertainties } (a + b)/2.$$

An alternative method (Method 2) considers the random variable  $x$  associated with the measured quantity. For this random variable, one assumes that the probability density function is an asymmetric normal distribution having a modal (most probable) value of  $x = X$ , a standard deviation  $b$  for  $x < X$ , and a standard deviation  $a$  for  $x > X$  (Fig. 9). Then the

average value of this distribution is

$$\langle x \rangle = X + \sqrt{2/\pi} (a - b),$$

with variance

$$\sigma^2 = (1 - 2/\pi)(a - b)^2 + ab. \tag{1}$$

The median value  $m$  which divides the distribution into two equal areas is given, for  $a > b$ , by

$$\text{erf}\left(\frac{m - X}{\sqrt{2}a}\right) = \frac{a - b}{2a}, \tag{2}$$

and by a similar expression for  $b > a$ .

One can then define the equivalent symmetric normal distribution that have a mean value equal to the median value  $m$  of the previous distribution with same variance  $\sigma$ .

If the shift  $m - X$  of the central value is small compared to  $a$  or  $b$ , expression (2) can be written [24]:

$$m - X \simeq \sqrt{\pi/8} (a - b)$$

$$m - X \simeq 0.6267 (a - b).$$

In order to allow for a small non-linearity that appears for higher values of  $m - X$ , the relation

$$m - X = 0.64 (a - b).$$

was adopted for Method 2. In NUBASE2016, Method 2 is used for the symmetrization of asymmetric half-lives and decay intensities. Table A illustrates the results from both methods.




Figure 9. Simulated asymmetric probability density function (heavy solid line) and the equivalent symmetric one (dashed line).



Table A. Examples of two different treatments of asymmetric half-life uncertainties.  
 Method 1 is the classical method, used previously, as in the AME1995.  
 Method 2 is the one developed in NUBASE2003, described in this Appendix.

Nuclide	Original $T_{1/2}$	Method 1	Method 2
$^{83}\text{Mo}$	6+30-3 ms	$20 \pm 17$	$23 \pm 19$
$^{100}\text{Kr}$	7+11-3 ms	$11 \pm 7$	$12 \pm 8$
$^{264}\text{Hs}$	327+448-120 $\mu\text{s}$	$490 \pm 280$	$540 \pm 300$
$^{266}\text{Mt}$	1.01+0.47-0.24 ms	$1.1 \pm 0.4$	$1.2 \pm 0.4$

## References

References in the text such as [1993Po.A] or [2015Ga38] are listed under "References used in the AME2016 and the NUBASE2016 evaluations", p. 030003-261.

- G. Audi, O. Bersillon, J. Blachot and A. H. Wapstra, Nucl. Phys. A, **624** : 1-124 (1997)
- G. Audi, O. Bersillon, J. Blachot and A. H. Wapstra, Nucl. Phys. A, **729**: 3 (2003)
- G. Audi, F.G. Kondev, M. Wang, B.Pfeiffer, X. Sun, J. Blachot and M. MacCormick, Chin. Phys. C, **36**: 12 (2012)  
<http://amdc.impcas.ac.cn/evaluation/data2012/paper/NUBASE2012.pdf>
- Edited by Yu. Litvinov and K. Blaum, Int. J. Mass Spectrom. **349 - 350**: (2013)
- T.W. Burrows, Nucl. Instrum. Meth. **286**: 595 (1990)  
<http://www.nndc.bnl.gov/ensdf/>
- T. Goigoux et al., Phys. Rev. Lett. **117**: 162501 (2016)
- B. Pritychenko, E. Běták, M.A. Kellett, B. Singh and J. Totans, Nucl. Instrum. Methods Phys. Res. A, **640**: 213 (2011)  
<http://www.nndc.bnl.gov/nsr/>
- L. Öhrström and J. Reedijk, Pure Appl. Chem. **88**: 1225 (2016)
- C.J. Gallagher, Jr. and S. A. Moszkowski, Phys. Rev. **111**: 1282 (1958)
- G. Audi, "A Lecture on the evaluation of atomic masses", in arXiv:  
<http://arxiv.org/abs/nucl-ex/0302020>
- M. MacCormick and G. Audi, Nucl. Phys. A, **925**: 61 (2014)
- E. P. Wigner, in Proceedings of the Robert A. Welch Foundation Conference on Chemical Research, edited by W. O. Milligan (Houston: Welch Foundation, 1958), Vol. 1, p. 88
- M. S. Antony, J. Britz, J. B. Bueb and A. Pape, At. Data Nucl. Data Tables , **33**: 447 (1985)  
M. S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables , **34**: 279 (1985)  
A. Pape and M. S. Antony, At. Data Nucl. Data Tables , **39**: 201 (1988)  
M. S. Antony, J. Britz and A. Pape, At. Data Nucl. Data Tables , **40**: 9 (1988)
- J. C. Hardy, I. S. Towner, Phys. Rev. C, **91**: 025501 (2015)
- General Policies, Nuclear Data Sheets, **113**: (2012)  $\nu$
- A. MacDonald, B. Karamy, K. Setoodehnia, B. Singh, Nucl. Data Sheets **114**: 397 (2013)
- J. Honkanen, J. Äystö and K. Eskola, Phys. Scr. **34**: (1986) 608
- P. G. Hansen and B. Jonson, Particle Emission from Nuclei, Vol.3, p. 157, CRC Press, Florida (1989)
- M. Thoennessen, Rep. Prog. Phys., **76**: 056301 (2013) and references therein
- <http://www.nndc.bnl.gov/xundl/>
- O. Bersillon and J. Blachot, NEANDC(E) 246/L, INDC(FR) 071/L, September 1991
- The NUBASE2016 files in the electronic distribution format and additional complementary information can be retrieved from the Atomic Mass Data Center (AMDC) at <http://amdc.impcas.ac.cn/>
- <https://www-nds.iaea.org/amdc/>
- R. D. Evans, The Atomic Nucleus (New York: McGraw-Hill, 1955) p. 766

**Table I. The NUBASE2016 table of nuclear and decay properties****EXPLANATION OF TABLE**

Data are presented in groups ordered according to increasing mass number  $A$ .

Nuclide	Nuclidic name: mass number $A = N + Z$ and element symbol (for $Z > 109$ see Section 2). Elements with upper suffix ‘ $m$ ’, ‘ $n$ ’, ‘ $p$ ’, ‘ $q$ ’, ‘ $r$ ’ or ‘ $x$ ’ indicate assignments to excited isomeric states (defined as higher states with half-lives greater than 100 ns). Suffixes ‘ $p$ ’ and ‘ $q$ ’ also indicate non-isomeric levels, but used in the AME2016. Suffix ‘ $r$ ’ also indicates a state from a proton resonance occurring in $(p, \gamma)$ reactions (e.g. $^{28}\text{Si}^r$ ). Suffix ‘ $x$ ’ also applies to mixtures of levels (with relative ratio $R$ , given in the ‘Half-life’ column), e.g. occurring in spallation reactions (indicated ‘ $\text{spmix}$ ’ in the ‘ $J^\pi$ ’ column) or fission (‘ $\text{fsmix}$ ’).																						
Mass excess	Mass excess $[M(\text{in u}) - A]$ , in keV, and its one standard deviation uncertainty as given in the ‘Atomic Mass Evaluation’ (AME2016, in the second part of this volume). Rounding-off policy: in cases where the furthest-left significant digit in the error is larger than 3, values and errors are rounded-off, but not to more than tens of keV. (Examples: $2345.67 \pm 2.78 \rightarrow 2345.7 \pm 2.8$ , $2345.67 \pm 4.68 \rightarrow 2346 \pm 5$ , but $2346.7 \pm 468.2 \rightarrow 2350 \pm 470$ ). # instead of a decimal point: value and uncertainty are not derived only from experimental data, but at least partly with estimates from TMS (see AME2016).																						
Excitation energy	For excited isomers only: energy difference, in keV, between levels adopted as higher level isomer and ground state isomer, and its one standard deviation uncertainty, as given in AME2016 when derived from the AME, otherwise as given by ENSDF. The rounding-off policy is the same as for the mass excesses (see above). # instead of a decimal point: value and uncertainty derived from trends in neighboring nuclides. The excitation energy is followed by its origin code when derived from a method other than $\gamma$ -ray spectrometry: <table> <tbody> <tr><td>MD</td><td>mass doublet</td></tr> <tr><td>RQ</td><td>reaction <math>Q</math>-value</td></tr> <tr><td>AD</td><td><math>\alpha</math> energy difference</td></tr> <tr><td>BD</td><td><math>\beta</math> energy difference</td></tr> <tr><td>p, 2p</td><td>one-, two-proton decay</td></tr> <tr><td>IT</td><td>combination of AME and <math>\gamma</math>-ray data</td></tr> <tr><td>Nm</td><td>estimated value derived using the Nilsson model</td></tr> </tbody> </table> When the existence of an isomer is questionable the following codes are used: <table> <tbody> <tr><td>EU</td><td>existence of isomer is under discussion (e.g. <math>^{73}\text{Zn}^n</math>). If existence is strongly doubted, no excitation energy and no mass are given. They are replaced by the mention “non existent” (e.g. <math>^{138}\text{Pm}^n</math>).</td></tr> <tr><td>RN</td><td>isomer has been proven not to exist (e.g. <math>^{181}\text{Pb}^m</math>). Excitation energy and mass are replaced by the mention “non existent”.</td></tr> </tbody> </table> Remark: codes EU and RN are also used when the discovery of a nuclide (e.g. $^{260}\text{Fm}$ or $^{289}\text{Lv}$ ) is questioned. In this case an estimate derived from trends in the mass surface is always given for the ground state mass. Isomeric assignment: <table> <tbody> <tr><td>*</td><td>if the uncertainty <math>\sigma</math> on the excitation energy <math>E</math> is greater than half the excitation energy (<math>\sigma &gt; E/2</math>), these quantities are followed by an asterisk (e.g. <math>^{130}\text{In}</math> and <math>^{130}\text{In}^m</math>).</td></tr> <tr><td>&amp;</td><td>when the ordering of the ground state isomer and the excited isomer are reversed as compared to ENSDF, an ampersand sign is added (e.g. <math>^{102}\text{Y}</math> and <math>^{102}\text{Y}^m</math>).</td></tr> </tbody> </table>	MD	mass doublet	RQ	reaction $Q$ -value	AD	$\alpha$ energy difference	BD	$\beta$ energy difference	p, 2p	one-, two-proton decay	IT	combination of AME and $\gamma$ -ray data	Nm	estimated value derived using the Nilsson model	EU	existence of isomer is under discussion (e.g. $^{73}\text{Zn}^n$ ). If existence is strongly doubted, no excitation energy and no mass are given. They are replaced by the mention “non existent” (e.g. $^{138}\text{Pm}^n$ ).	RN	isomer has been proven not to exist (e.g. $^{181}\text{Pb}^m$ ). Excitation energy and mass are replaced by the mention “non existent”.	*	if the uncertainty $\sigma$ on the excitation energy $E$ is greater than half the excitation energy ( $\sigma > E/2$ ), these quantities are followed by an asterisk (e.g. $^{130}\text{In}$ and $^{130}\text{In}^m$ ).	&	when the ordering of the ground state isomer and the excited isomer are reversed as compared to ENSDF, an ampersand sign is added (e.g. $^{102}\text{Y}$ and $^{102}\text{Y}^m$ ).
MD	mass doublet																						
RQ	reaction $Q$ -value																						
AD	$\alpha$ energy difference																						
BD	$\beta$ energy difference																						
p, 2p	one-, two-proton decay																						
IT	combination of AME and $\gamma$ -ray data																						
Nm	estimated value derived using the Nilsson model																						
EU	existence of isomer is under discussion (e.g. $^{73}\text{Zn}^n$ ). If existence is strongly doubted, no excitation energy and no mass are given. They are replaced by the mention “non existent” (e.g. $^{138}\text{Pm}^n$ ).																						
RN	isomer has been proven not to exist (e.g. $^{181}\text{Pb}^m$ ). Excitation energy and mass are replaced by the mention “non existent”.																						
*	if the uncertainty $\sigma$ on the excitation energy $E$ is greater than half the excitation energy ( $\sigma > E/2$ ), these quantities are followed by an asterisk (e.g. $^{130}\text{In}$ and $^{130}\text{In}^m$ ).																						
&	when the ordering of the ground state isomer and the excited isomer are reversed as compared to ENSDF, an ampersand sign is added (e.g. $^{102}\text{Y}$ and $^{102}\text{Y}^m$ ).																						

Half-life	<p>s = seconds; m = minutes; h = hours; d = days; y = years;  1 y = 31 556 926 s or 365.2422 d  adopted values for NUBASE (see text)  STABLE = stable nuclide, or nuclide for which no finite half-life value  has been found.</p> <p># value estimated from trends in neighboring nuclides with the same <math>Z</math> and <math>N</math> parities.</p> <p>subunits:</p> <table border="0"> <tr> <td>ms</td><td>:</td><td><math>10^{-3}</math></td><td>s</td><td>millisecond</td> <td>ky</td><td>:</td><td><math>10^3</math></td><td>y</td><td>kiloyear</td> </tr> <tr> <td><math>\mu</math>s</td><td>:</td><td><math>10^{-6}</math></td><td>s</td><td>microsecond</td> <td>My</td><td>:</td><td><math>10^6</math></td><td>y</td><td>megayear</td> </tr> <tr> <td>ns</td><td>:</td><td><math>10^{-9}</math></td><td>s</td><td>nanosecond</td> <td>Gy</td><td>:</td><td><math>10^9</math></td><td>y</td><td>gigayear</td> </tr> <tr> <td>ps</td><td>:</td><td><math>10^{-12}</math></td><td>s</td><td>picosecond</td> <td>Ty</td><td>:</td><td><math>10^{12}</math></td><td>y</td><td>terayear</td> </tr> <tr> <td>fs</td><td>:</td><td><math>10^{-15}</math></td><td>s</td><td>femtosecond</td> <td>Py</td><td>:</td><td><math>10^{15}</math></td><td>y</td><td>petayear</td> </tr> <tr> <td>as</td><td>:</td><td><math>10^{-18}</math></td><td>s</td><td>attosecond</td> <td>Ey</td><td>:</td><td><math>10^{18}</math></td><td>y</td><td>exayear</td> </tr> <tr> <td>zs</td><td>:</td><td><math>10^{-21}</math></td><td>s</td><td>zeptosecond</td> <td>Zy</td><td>:</td><td><math>10^{21}</math></td><td>y</td><td>zettayear</td> </tr> <tr> <td>ys</td><td>:</td><td><math>10^{-24}</math></td><td>s</td><td>yoctosecond</td> <td>Yy</td><td>:</td><td><math>10^{24}</math></td><td>y</td><td>yottayear</td> </tr> </table> <p><math>R</math>: For isomeric mixtures only, it is the production ratio of the excited isomer state to the ground state isomer.</p>	ms	:	$10^{-3}$	s	millisecond	ky	:	$10^3$	y	kiloyear	$\mu$ s	:	$10^{-6}$	s	microsecond	My	:	$10^6$	y	megayear	ns	:	$10^{-9}$	s	nanosecond	Gy	:	$10^9$	y	gigayear	ps	:	$10^{-12}$	s	picosecond	Ty	:	$10^{12}$	y	terayear	fs	:	$10^{-15}$	s	femtosecond	Py	:	$10^{15}$	y	petayear	as	:	$10^{-18}$	s	attosecond	Ey	:	$10^{18}$	y	exayear	zs	:	$10^{-21}$	s	zeptosecond	Zy	:	$10^{21}$	y	zettayear	ys	:	$10^{-24}$	s	yoctosecond	Yy	:	$10^{24}$	y	yottayear
ms	:	$10^{-3}$	s	millisecond	ky	:	$10^3$	y	kiloyear																																																																								
$\mu$ s	:	$10^{-6}$	s	microsecond	My	:	$10^6$	y	megayear																																																																								
ns	:	$10^{-9}$	s	nanosecond	Gy	:	$10^9$	y	gigayear																																																																								
ps	:	$10^{-12}$	s	picosecond	Ty	:	$10^{12}$	y	terayear																																																																								
fs	:	$10^{-15}$	s	femtosecond	Py	:	$10^{15}$	y	petayear																																																																								
as	:	$10^{-18}$	s	attosecond	Ey	:	$10^{18}$	y	exayear																																																																								
zs	:	$10^{-21}$	s	zeptosecond	Zy	:	$10^{21}$	y	zettayear																																																																								
ys	:	$10^{-24}$	s	yoctosecond	Yy	:	$10^{24}$	y	yottayear																																																																								
$J^\pi$	<p>Spin and parity:  () uncertain spin and/or parity.  # values estimated from trends in neighboring nuclides with the same <math>Z</math> and <math>N</math> parities.  high high spin.  low low spin.  am same <math>J^\pi</math> as <math>\alpha</math>-decay parent  <i>T</i> Isopin multiplet for isobaric analog states (IAS).</p> <p>For isomeric mixtures only: mix (spmix and fsmix if observed in spallation and fission, respectively).</p>																																																																																
Ens	<p>Year of the ENSDF file archive  (in order to reduce the width of the Table, the two century digits are omitted).</p>																																																																																
Reference	<p>Reference keys:  (in order to reduce the width of the Table, the two century digits are omitted. However, at the end of this volume the full reference key-number is given, ie. 2010Cr02 as opposed to 10Cr02)</p> <p>10Cr02 updates to ENSDF derived from a regular journal. These keys are taken from Nuclear Data Sheets. Where not yet available, the style 12Ma.1 is provisionally adopted.</p> <p>12Dr.A updates to ENSDF derived from an abstract, preprint, private communication, conference, thesis or annual report.</p> <p>AHW (or FGK, GAU, HWJ, JBL, MMC, WGM), re-interpretation by one of the evaluators of NUBASE.</p> <p>Mirror deduced from mirror nuclide properties.</p> <p>Imme deduced from Isobaric Multiplet Mass Equation.</p> <p>The reference key-numbers are followed by one, two or three letter codes which specifies the added or modified physical quantities:</p> <table border="0"> <tr> <td>E</td><td>for the isomer excitation energy</td> </tr> <tr> <td>T</td><td>for half-life</td> </tr> <tr> <td>J</td><td>for spin and/or parity</td> </tr> <tr> <td>D</td><td>for decay mode and/or intensity</td> </tr> <tr> <td>I</td><td>for identification</td> </tr> </table>	E	for the isomer excitation energy	T	for half-life	J	for spin and/or parity	D	for decay mode and/or intensity	I	for identification																																																																						
E	for the isomer excitation energy																																																																																
T	for half-life																																																																																
J	for spin and/or parity																																																																																
D	for decay mode and/or intensity																																																																																
I	for identification																																																																																
Year of discovery	<p>for ground states [15] and for excited isomers (see text).</p>																																																																																

Decay modes and intensities	<p>Decay modes followed by their intensities (in %), and their one standard deviation uncertainties. The special notation 1.8e-12 stands for <math>1.8 \times 10^{-12}</math>.</p> <p>The uncertainties are given - only in this field - in the ENSDF-style: <math>\alpha=25.9\ 23</math> stands for <math>\alpha=25.9 \pm 2.3\ %</math></p> <p>The ordering is according to decreasing intensities.</p> <p><math>\alpha\ ?</math> means <math>\alpha</math> decay is energetically allowed.</p> <p><math>\alpha=?</math> means <math>\alpha</math> decay has been observed but not yet quantified.</p> <table border="0"> <tr> <td><math>\alpha</math></td> <td><math>\alpha</math> emission</td> <td></td> </tr> <tr> <td>p 2p</td> <td>proton emission</td> <td>2-proton emission</td> </tr> <tr> <td>n 2n</td> <td>neutron emission</td> <td>2-neutron emission</td> </tr> <tr> <td><math>\epsilon</math></td> <td>electron capture</td> <td></td> </tr> <tr> <td><math>e^+</math></td> <td>positron emission</td> <td></td> </tr> <tr> <td><math>\beta^+</math></td> <td><math>\beta^+</math> decay</td> <td>(<math>\beta^+ = \epsilon + e^+</math>)</td> </tr> <tr> <td><math>\beta^-</math></td> <td><math>\beta^-</math> decay</td> <td></td> </tr> <tr> <td><math>2\beta^-</math></td> <td>double <math>\beta^-</math> decay</td> <td></td> </tr> <tr> <td><math>2\beta^+</math></td> <td>double <math>\beta^+</math> decay</td> <td></td> </tr> <tr> <td><math>\beta^-n</math></td> <td><math>\beta^-</math> delayed neutron emission</td> <td></td> </tr> <tr> <td><math>\beta^-2n</math></td> <td><math>\beta^-</math> delayed 2-neutron emission</td> <td></td> </tr> <tr> <td><math>\beta^+p</math></td> <td><math>\beta^+</math> delayed proton emission</td> <td></td> </tr> <tr> <td><math>\beta^+2p</math></td> <td><math>\beta^+</math> delayed 2-proton emission</td> <td></td> </tr> <tr> <td><math>\beta^-\alpha</math></td> <td><math>\beta^-</math> delayed <math>\alpha</math> emission</td> <td></td> </tr> <tr> <td><math>\beta^+\alpha</math></td> <td><math>\beta^+</math> delayed <math>\alpha</math> emission</td> <td></td> </tr> <tr> <td><math>\beta^-d</math></td> <td><math>\beta^-</math> delayed deuteron emission</td> <td></td> </tr> <tr> <td>IT</td> <td>internal transition</td> <td></td> </tr> <tr> <td>SF</td> <td>spontaneous fission</td> <td></td> </tr> <tr> <td><math>\beta^+SF</math></td> <td><math>\beta^+</math> delayed fission</td> <td></td> </tr> <tr> <td><math>\beta^-SF</math></td> <td><math>\beta^-</math> delayed fission</td> <td></td> </tr> <tr> <td><math>^{24}\text{Ne}</math></td> <td>heavy cluster emission</td> <td></td> </tr> <tr> <td>...</td> <td>list is continued in a remark, at the end of the A-group</td> <td></td> </tr> </table> <p>For long-lived nuclides:</p> <table border="0"> <tr> <td>IS</td> <td>Isotopic abundance (from [2011Be53])</td> </tr> </table>	$\alpha$	$\alpha$ emission		p 2p	proton emission	2-proton emission	n 2n	neutron emission	2-neutron emission	$\epsilon$	electron capture		$e^+$	positron emission		$\beta^+$	$\beta^+$ decay	( $\beta^+ = \epsilon + e^+$ )	$\beta^-$	$\beta^-$ decay		$2\beta^-$	double $\beta^-$ decay		$2\beta^+$	double $\beta^+$ decay		$\beta^-n$	$\beta^-$ delayed neutron emission		$\beta^-2n$	$\beta^-$ delayed 2-neutron emission		$\beta^+p$	$\beta^+$ delayed proton emission		$\beta^+2p$	$\beta^+$ delayed 2-proton emission		$\beta^-\alpha$	$\beta^-$ delayed $\alpha$ emission		$\beta^+\alpha$	$\beta^+$ delayed $\alpha$ emission		$\beta^-d$	$\beta^-$ delayed deuteron emission		IT	internal transition		SF	spontaneous fission		$\beta^+SF$	$\beta^+$ delayed fission		$\beta^-SF$	$\beta^-$ delayed fission		$^{24}\text{Ne}$	heavy cluster emission		...	list is continued in a remark, at the end of the A-group		IS	Isotopic abundance (from [2011Be53])
$\alpha$	$\alpha$ emission																																																																				
p 2p	proton emission	2-proton emission																																																																			
n 2n	neutron emission	2-neutron emission																																																																			
$\epsilon$	electron capture																																																																				
$e^+$	positron emission																																																																				
$\beta^+$	$\beta^+$ decay	( $\beta^+ = \epsilon + e^+$ )																																																																			
$\beta^-$	$\beta^-$ decay																																																																				
$2\beta^-$	double $\beta^-$ decay																																																																				
$2\beta^+$	double $\beta^+$ decay																																																																				
$\beta^-n$	$\beta^-$ delayed neutron emission																																																																				
$\beta^-2n$	$\beta^-$ delayed 2-neutron emission																																																																				
$\beta^+p$	$\beta^+$ delayed proton emission																																																																				
$\beta^+2p$	$\beta^+$ delayed 2-proton emission																																																																				
$\beta^-\alpha$	$\beta^-$ delayed $\alpha$ emission																																																																				
$\beta^+\alpha$	$\beta^+$ delayed $\alpha$ emission																																																																				
$\beta^-d$	$\beta^-$ delayed deuteron emission																																																																				
IT	internal transition																																																																				
SF	spontaneous fission																																																																				
$\beta^+SF$	$\beta^+$ delayed fission																																																																				
$\beta^-SF$	$\beta^-$ delayed fission																																																																				
$^{24}\text{Ne}$	heavy cluster emission																																																																				
...	list is continued in a remark, at the end of the A-group																																																																				
IS	Isotopic abundance (from [2011Be53])																																																																				

\* A remark on the corresponding nuclide is given below the block of data corresponding to the same A.

*Remarks.* For nuclides marked with an asterisk at the end of the line, extra comments have been added. They are collected in groups at the end of each block of data corresponding to the same A. They start with a letter code, similar the ones following the reference key-number, as given above, indicating to which quantity the remark applies. They give:

- i) Continuation for the list of decays. In this case, the remark starts with three dots.
- ii) Information explaining how a value has been derived.
- iii) Reasons for changing a value or its uncertainty as given by the authors, or for rejecting it.
- iv) Complementary references to updated data.
- v) Separate values used in the adopted average.

TNN : Trends from neighboring nuclides.

$\tau$  : meanlife (or lifetime)  $T = \tau \times \ln 2$

**Table I. The NUBASE2016 table (Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
$^1_0\text{n}$	8071.3171	0.0005	613.9 s	0.6	$1/2^+$	06		$\beta^-$ =100	*	
$^1_1\text{H}$	7288.9706	0.0001	STABLE		$1/2^+$	06	11Be53 D	1920	IS=99.9885 70	
* $^1_0\text{n}$	T: also $^{15}\text{Ar}07=610.1(0.8)$ $\tau=880.2(1.2)$ , $^{13}\text{Yu}07=615.3(1.5)$ $\tau=887.7(2.2)$ ,								**	
* $^1_1\text{H}$	T: $^{12}\text{Ar}05=611.1(1.5)$ $\tau=881.6(2.1)$								**	
$^2_1\text{H}$	13135.7217	0.0001	STABLE		$1^+$	03		1932	IS=0.0115 70	
$^3_1\text{H}$	14949.8099	0.0002	12.32 y	0.02	$1/2^+$	00		1934	$\beta^-$ =100	
$^3_2\text{He}$	14931.2179	0.0002	STABLE		$1/2^+$	98		1934	IS=0.000134 3	
$^3_3\text{Li}$	28670#	2000#	p-unstable			98		1969	p?	*
* $^3_3\text{Li}$	I: identification in $^{69}\text{Wi}13$ not accepted, see ENSDF'98								**	
$^4_1\text{H}$	24620	100	139 ys	10	$2^-$	98	03Me11 T	1981	n=100	*
$^4_2\text{He}$	2424.9156	0.0001	STABLE		$0^+$	98		1908	IS=99.999866 3	
$^4_3\text{Li}$	25320	210	91 ys	9	$2^-$	98	65Ce02 T	1965	p=100	
* $^4_1\text{H}$	T: width=3.28(0.23) MeV; also $^{91}\text{Go}19=4.7(1.0)$ outweighed, not used								**	
$^5_1\text{H}$	32890	90	> 910 ys		$(1/2^+)$	02	03Go11 T	1987	2n=100	*
$^5_2\text{He}$	11231	20	700 ys	30	$3/2^-$	02		1937	n=100	
$^5_3\text{Li}$	11680	50	370 ys	30	$3/2^-$	02		1941	p=100	
$^5_4\text{Be}$	37140#	2000#			$1/2^+\#$	02			p?	
* $^5_1\text{H}$	T: from width < 0.5 MeV; conflicting with $^{01}\text{Ko}52=280(50)$ ys, width=1.9(0.4)								**	
* $^5_2\text{He}$	T: (same authors) but with instrumental resolution=1.3 MeV								**	
* $^5_3\text{Li}$	T: others $^{91}\text{Go}19=66(25)$ ys $^{95}\text{Al}31=110$ ys probably for higher state								**	
* $^5_4\text{Be}$	J: from angular distribution consistent with $l = 0$								**	
$^6_1\text{H}$	41880	250	290 ys	70	$2^\-\#$	02		1984	n?; 3n?	
$^6_2\text{He}$	17592.10	0.05	806.92 ms	0.24	$0^+$	02	15Pf01 D	1936	$\beta^-$ =100; $\beta^-$ -d=0.000278 18	*
$^6_3\text{Li}$	14086.8789	0.0014	STABLE		$1^+$	02		1921	IS=7.59 4	
$^6_4\text{Li}^i$	17649.76	0.10	3562.88	0.10	$0^+\text{T}=1$	02	81Ro02 E	1981	IT=100	
$^6_5\text{Be}$	18375	5	5.0 zs	0.3	$0^+$	02		1958	2p=100	
$^6_6\text{B}$	47320#	2000#	p-unstable#		$2^\-\#$				2p?	
* $^6_2\text{He}$	D: other $\beta^-$ -d from $^{09}\text{Ra}33=1.65(0.10)\text{e}-6$ but with 525 keV threshold								**	
* $^6_6\text{B}$	T: symmetrized from $^{12}\text{Kn}01=806.89(0.11)(+0.23-0.19)$								**	
$^7_1\text{H}$	49140#	1000#	500# ys		$1/2^+\#$			2003	2n?	
$^7_2\text{He}$	26073	8	2.51 zs	0.07	$(3/2)^-$	03	12Ca05 T	1967	n=100	*
$^7_3\text{Li}$	14907.105	0.004	STABLE		$3/2^-$	03		1921	IS=92.41 4	
$^7_4\text{Li}^i$	26150	30	11250	30	RQ				$3/2^-\text{T}=3/2$	
$^7_5\text{Be}$	15769.00	0.07	53.22 d	0.06	$3/2^-$	03		1938	$\epsilon$ =100	
$^7_6\text{Be}^i$	26750	30	10980	30	RQ				p?; $3\text{He}$ ?; $\alpha$ ?	
$^7_7\text{B}$	27677	25	570 ys	14	$(3/2)^-$	14	11Ch32 T	1967	p=100	*
* $^7_2\text{He}$	T: from $^{12}\text{Ca}05=182(5)$ keV								**	
* $^7_7\text{B}$	T: others $^{09}\text{Ak}03=190(30)$ $^{08}\text{De}29=125(+40-15)$ $^{02}\text{Me}07=150(80)$ $^{69}\text{St}02=160(30)$								**	
* $^7_7\text{B}$	T: from width $^{11}\text{Ch}32=801(20)$ keV $^{570}(14)$ ys								**	
$^8_2\text{He}$	31609.68	0.09	119.1 ms	1.2	$0^+$	05		1965	$\beta^-$ =100; $\beta^-$ -n=16 1; $\beta^-$ -t=0.9 1	
$^8_3\text{Li}$	20945.80	0.05	839.40 ms	0.36	$2^+$	05	10Fl01 T	1935	$\beta^-$ =100; $\beta^-$ - $\alpha$ =100	*
$^8_4\text{Li}^i$	31768	5	10822	5	RQ				$0^+\text{T}=2$	
$^8_5\text{Be}$	4941.67	0.04	81.9 as	3.7	$0^+$	05		1932	$\alpha$ =100	
$^8_6\text{Be}^i$	21568	3	16626	3	RQ			2004	$\alpha\approx 100$	*
$^8_7\text{Be}^j$	32436.0	2.0	27494.3	2.0	RQ				$0^+\text{T}=2$	*
$^8_8\text{B}$	22921.6	1.0	770 ms	3	$2^+$	05		1950	n=39.4; d=27.0; $3\text{H}=11.7$ ; $\alpha=7.9$ ; ...	*
$^8_9\text{Bx}^i$	33546	8	10624	8	RQ			1975	$\beta^+=100$ ; $\beta^+$ - $\alpha$ =100	*
$^8_{10}\text{C}$	35064	18	3.5 zs	1.4	$0^+$	05	11Ch32 T	1974	2p=100	*
* $^8_3\text{Li}$	D: $\beta^-$ decay to first $2^+$ state in $^8\text{Be}$ , which decays 100% in $2\alpha$								**	
* $^8_6\text{Be}^i$	E: strongest frg; other: $^{296}(3)$ higher $I(16626)/I(16922)=1.22$ in $^6\text{Li}(^6\text{Li},\alpha)$								**	
* $^8_6\text{Be}^i$	E: and 1.15 in $^{10}\text{B}(d,\alpha)$ ; see $^{04}\text{Ti}06$ p.213								**	
* $^8_7\text{Be}^j$	D: ...; p=6.9; $3\text{He}=6.6$ ; IT=0.60								**	
* $^8_8\text{B}$	D: $\beta^+$ to 2 excited states in $^8\text{Be}$ , then $\alpha$ and $\gamma$ , but not to $^8\text{Be}$ ground-state								**	
* $^8_{10}\text{C}$	T: from width $^{130}(50)$ keV $^{3.51}(1.35)$ zs								**	

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)		Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>9</sup> He	40940	50			2.5	zs 2.3	1/2 <sup>(+)</sup>	06	16Ub01	J	1987	n=100	*
<sup>9</sup> Li	24954.90	0.19			178.3	ms 0.4	3/2 <sup>-</sup>	06	95Re.A	D	1951	$\beta^-$ =100; $\beta^-$ n=50.8 2	
<sup>9</sup> Be	11348.45	0.08			STABLE		3/2 <sup>-</sup>	06			1921	IS=100.	
<sup>9</sup> Be <sup>i</sup>	25738.8	1.7	14390.3	1.7	RQ	1.25 as 0.10	3/2 <sup>-</sup> T=3/2	06			1976		
<sup>9</sup> B	12416.5	0.9				800 zs 300	3/2 <sup>-</sup>	06			1940	p=100	
<sup>9</sup> Bx <sup>i</sup>	27071.0	2.3	14654.5	2.5	RQ		3/2 <sup>-</sup> T=3/2	06					
<sup>9</sup> C	28911.0	2.1				126.5 ms 0.9	(3/2 <sup>-</sup> )	06			1964	$\beta^+$ =100; $\beta^+$ p=61.6; $\beta^+$ $\alpha$ =38.4	
* <sup>9</sup> He	T: derived from width 13Al14=180(100); other width 99Bo26=100(60)												**
<sup>10</sup> He	49200	90			3.1	zs 2.0	0 <sup>+</sup>	07			1994	2n=100	*
<sup>10</sup> Li	33053	13			2.0	zs 0.5	(1 <sup>-</sup> , 2 <sup>-</sup> )	07	94Yo01	TJ	1975	n=100	
<sup>10</sup> Li <sup>m</sup>	33250	40	200	40	RQ	3.7 zs 1.5	1 <sup>+</sup>	07	97Zi04	T	1994	IT=100	*
<sup>10</sup> Li <sup>n</sup>	33530	40	480	40	RQ	1.35 zs 0.24	(2 <sup>+</sup> )	07	94Yo01	T	1993	IT=100	*
<sup>10</sup> Be	12607.49	0.08				1.51 My 0.04	0 <sup>+</sup>	07			1935	$\beta^-$ =100	
<sup>10</sup> Be <sup>i</sup>	33787	21	21179	21	RQ		(2 <sup>-</sup> )T=2	07				n ?; p ?; 3H ?	
<sup>10</sup> B	12050.609	0.015				STABLE	3 <sup>+</sup>	07			1920	IS=19.9 7	
<sup>10</sup> B <sup>i</sup>	13790.66	0.04	1740.05	0.04			0 <sup>+</sup> T=1	07				IT=100	
<sup>10</sup> C	15698.67	0.07				19.3009 s 0.0017	0 <sup>+</sup>	07	16Du10	T	1949	$\beta^+$ =100	*
<sup>10</sup> N	38800	400				200 ys 140	(2 <sup>-</sup> )	07	02Le16	TJ	2002	p ?	
* <sup>10</sup> He	D: most probably 2 neutron emitter from S <sub>2n</sub> =-1440(90) keV												**
* <sup>10</sup> Li <sup>m</sup>	T: average 97Zi04=120(+100-50) 94Yo01=100(70) keV												**
* <sup>10</sup> Li <sup>n</sup>	T: average 94Yo01=358(23) 93Bo03=150(70) keV, Birge ratio B=2.8												**
* <sup>10</sup> C	T: also 16Du10(2)=19.2969(0.0074) 09Ba04=19.282(0.011) 08Ia01=19.310(0.004)												**
<sup>11</sup> Li	40728.3	0.6			8.75	ms 0.14	3/2 <sup>-</sup>	12	12Ke01	D	1966	$\beta^-$ =100; $\beta^-$ n=86.3 9; $\beta^-$ 2n=4.1 4; ...	*
<sup>11</sup> Be	20177.17	0.24			13.76	s 0.07	1/2 <sup>+</sup>	12	81Al03	D	1958	$\beta^-$ =100; $\beta^-$ $\alpha$ =2.9 4; ...	*
<sup>11</sup> Be <sup>i</sup>	41336	20	21158	20	RQ	0.93 zs 0.13	3/2 <sup>-</sup> T=5/2		MMC162	J	1997	IT ?	*
<sup>11</sup> B	8667.707	0.012				STABLE	3/2 <sup>-</sup>	12			1920	IS=80.1 7	
<sup>11</sup> B <sup>i</sup>	21228	9	12560	9	RQ		T=3/2	12			1963		
<sup>11</sup> B <sup>j</sup>	42230	80	33570	80	2p		3/2 <sup>-</sup> T=5/2		MMC146	J			
<sup>11</sup> C	10649.40	0.06				20.364 m 0.014	3/2 <sup>-</sup>	12			1934	$\beta^+$ =100	
<sup>11</sup> C <sup>i</sup>	22810	40	12160	40	RQ		1/2 <sup>+</sup> T=3/2	12	71Wa21	D	1971	p=?	
<sup>11</sup> N	24300	50				550 ys 20	1/2 <sup>+</sup>	12			1974	p=100	*
<sup>11</sup> N <sup>m</sup>	25040	80	740	60		690 ys 80	1/2 <sup>-</sup>	12	96Ax01	ETJ	1974	p=100	*
* <sup>11</sup> Li	D: ...; $\beta^-$ 3n=1.9 2; $\beta^-$ $\alpha$ =1.7 3; $\beta^-$ d=0.0130 13; $\beta^-$ t=0.0093 8												**
* <sup>11</sup> Li	D: total $\beta^-$ delayed neutron emission Pn=100.3(1.4)%												**
* <sup>11</sup> Be	D: ...; $\beta^-$ p=0.00083 9; $\beta^-$ n ?												**
* <sup>11</sup> Be <sup>i</sup>	T: from width 97Te07=490(70) keV												**
* <sup>11</sup> N	T: from ENSDF2012: width=830(30) keV												**
<sup>12</sup> Li	49010	30			< 10	ns		00	74Bo05	I	2008	n ?	
<sup>12</sup> Be	25077.8	1.9			21.50	ms 0.04	0 <sup>+</sup>	00	01Be53	T	1966	$\beta^-$ =100; $\beta^-$ n=0.50 3	*
<sup>12</sup> Be <sup>m</sup>	27328.8	2.1	2251	1		229 ns 8	0 <sup>+</sup>		07Sh34	EJT	2007	IT=100	
<sup>12</sup> B	13369.4	1.3				20.20 ms 0.02	1 <sup>+</sup>	00	66Sc23	D	1935	$\beta^-$ =100; $\beta^-$ $\alpha$ =1.6 3	
<sup>12</sup> B <sup>i</sup>	26088	19	12719	19	RQ		0 <sup>+</sup> T=2	00	08Ch28	J			*
<sup>12</sup> C	0.0	0.0				STABLE	0 <sup>+</sup>	00			1919	IS=98.93 8	
<sup>12</sup> C <sup>i</sup>	15108	3	15108	3	RQ		1 <sup>+</sup> T=1	00				IT=?; $\alpha$ ?	
<sup>12</sup> C <sup>j</sup>	27595.0	2.4	27595.0	2.4	RQ		0 <sup>+</sup> T=2	00					
<sup>12</sup> N	17338.1	1.0				11.000 ms 0.016	1 <sup>+</sup>	00	66Sc23	D	1949	$\beta^+$ =100; $\beta^+$ $\alpha$ =3.5 5	
<sup>12</sup> N <sup>i</sup>	29534	29	12195	29	2p		(0 <sup>+</sup> )T=2		MMC142	J			
<sup>12</sup> O	31915	24				> 6.3 zs	0 <sup>+</sup>	00	12Ja11	T	1978	2p=60 30	*
* <sup>12</sup> Be	D: from 99Be53; also 95Re.A=0.52(0.09)% outweighed, not used												**
* <sup>12</sup> B <sup>i</sup>	J: 08Ch28 "suggests that the 12.75-MeV, ... was a T=1 state, not the IAS"												**
* <sup>12</sup> O	T: from width 12Ja11 < 72 keV; others 09Su14=600(500)keV 95Kr03t=578(205)keV												**
<sup>13</sup> Li	56980	70			3.3	zs 1.2	3/2 <sup>-</sup> #		08Ak03	D	2008	2n=100	*
<sup>13</sup> Be	33659	10			1.0	zs 0.7	(1/2 <sup>-</sup> )		10Ko17	TJ	1983	n ?	*
<sup>13</sup> Be <sup>p</sup>	35160	50	1500	50	RQ		(5/2 <sup>+</sup> )				1992		
<sup>13</sup> B	16561.9	1.0				17.33 ms 0.17	3/2 <sup>-</sup>	00			1956	$\beta^-$ =100; $\beta^-$ n=0.28 4	
<sup>13</sup> C	3125.0088	0.0002				STABLE	1/2 <sup>-</sup>	01			1929	IS=1.07 8	
<sup>13</sup> C <sup>i</sup>	18233.8	1.1	15108.8	1.1	RQ		3/2 <sup>-</sup> T=3/2	00				IT=0.82 7; N ?; $\alpha$ ?	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>13</sup> N	5345.48	0.27							
<sup>13</sup> N <sup>i</sup>	20410.59	0.18	15065.1	0.3	RQ				$\beta^+=100$ IT=4.9 3; P ?; $\alpha$ ?
<sup>13</sup> O	23115	10							$\beta^+=100$ ; $\beta^+p=10.9$ 20
* <sup>13</sup> Li	T : from width 13Ko03=125(60-40) keV								
* <sup>13</sup> Be	T : from width 10Ko17=450(30) keV; other 95Pe12=300(200) keV								
* <sup>13</sup> Be	J : 1/2 <sup>+</sup> assigned to ground state in 01Th01 and 08Ch07, questioned in 10Ko17								
* <sup>13</sup> Be	J : see discussion in AME2012, Part I, Section 6.3, p.1313								
* <sup>13</sup> Be	J : and 14Ra07=1/2 <sup>+</sup>								
<sup>14</sup> Be	39950	130							$\beta^-=100$ ; $\beta^-n=98$ 2; $\beta^-2n=0.8$ 08; ... *
<sup>14</sup> Be <sup>p</sup>	41470	60	1520	150	RQ				
<sup>14</sup> B	23664	21							$\beta^-=100$ ; $\beta^-n=6.04$ 23; $\beta^-2n$ ?
<sup>14</sup> B <sup>i</sup>	40728	20	17065	29	RQ				IT ? *
<sup>14</sup> C	3019.893	0.004							$\beta^-=100$
<sup>14</sup> C <sup>i</sup>	25120	100	22100	100					IT=100
<sup>14</sup> N	2863.4167	0.0001			STABLE				IS=99.636 20
<sup>14</sup> N <sup>i</sup>	5176.007	0.010	2312.590	0.010					IT=100
<sup>14</sup> O	8007.781	0.025							$\beta^+=100$
<sup>14</sup> F	31960	40							p ? *
* <sup>14</sup> Be	D : ... ; $\beta^-3n=0.2$ 2; $\beta^-t=0.02$ 1; $\beta^-a<0.004$								
* <sup>14</sup> Be	D : supersedes 99Be53, same group								
* <sup>14</sup> B <sup>i</sup>	T : from width 01Ta23=110(50) keV								
* <sup>14</sup> O	T : average 13La23(beta)=70.610(0.030), 04Ba78=70.641(0.020),								
* <sup>14</sup> O	T : 78Wi04=70.613(0.025) and 73Cl12=70.590(0.030);								
* <sup>14</sup> O	T : others outweighed : 13La23( $\gamma$ )=70.632(0.094), 06Bu12=70.696(0.052)								
* <sup>14</sup> O	T : and 01Ga59=70.560(0.049)								
* <sup>14</sup> F	T : from width 10Go16=910(100) keV								
<sup>15</sup> Be	49830	170							n=100 *
<sup>15</sup> B	28958	21							$\beta^-=100$ ; $\beta^-n=93.6$ 12; $\beta^-2n=0.4$ 2 *
<sup>15</sup> C	9873.1	0.8							$\beta^-=100$
<sup>15</sup> N	101.4387	0.0006			STABLE				IS=0.364 20
<sup>15</sup> N <sup>i</sup>	11717	4	11615	4	RQ				n ?; p ?; IT=0.00523 19
<sup>15</sup> O	2855.6	0.5							$\beta^+=100$
<sup>15</sup> O <sup>i</sup>	14020#	40#	11165#	35#					p=100
<sup>15</sup> F	16567	14							p=100 *
<sup>15</sup> Ne	40220	70			2p				2p=100 *
* <sup>15</sup> Be	T : from width 13Sn02=575(200) keV								
* <sup>15</sup> B	D : $\beta^-2n$ intensity is from 89Re.A J : given in 91Aj01								
* <sup>15</sup> B	T : also 03Ye02=9.86(+0.15-0.19)								
* <sup>15</sup> F	T : from 16De15=370(70)(+200-0) keV								
* <sup>15</sup> Ne	T : from width 590(230) keV								
<sup>16</sup> Be	57450	170							2n=100 *
<sup>16</sup> B	37113	25							n ?
<sup>16</sup> C	13694	4							$\beta^-=100$ ; $\beta^-n=97.9$ 23
<sup>16</sup> N	5683.9	2.3							$\beta^-=100$ ; $\beta^-a=0.00145$ 8 *
<sup>16</sup> N <sup>m</sup>	5804.3	2.3	120.42	0.12					IT $\approx$ 100; $\beta^-=0.00040$ 4 *
<sup>16</sup> N <sup>i</sup>	15613	7	9929	7	RQ				
<sup>16</sup> O	-4737.0013	0.0001			STABLE				IS=99.757 16
<sup>16</sup> O <sup>i</sup>	8059	4	12796	4	RQ				IT=100
<sup>16</sup> O <sup>j</sup>	17984	4	22721	4	RQ				
<sup>16</sup> F	10680	8							p=100
<sup>16</sup> Ne	23987	20							2p=100 *
* <sup>16</sup> Be	T : from decay width 0.8(+0.1-0.2) MeV								
* <sup>16</sup> N	D : symmetrized from 16Re01=0.00149(5stat)(+0-10sys); other 74Ne10=0.00100 7								
* <sup>16</sup> N <sup>m</sup>	D : from B.Singh, average 3 results 83Mi20 83Ga18 (also 82Ga05) 75Pa01								
* <sup>16</sup> Ne	T : 14Br19 L<80 keV (3 $\sigma$ upper limit)								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
$^{17}\text{B}$	43720	200	5.08 ms	0.05	(3/2 <sup>-</sup> )	99 88Du09	D	1973	$\beta^- = 100$ ; $\beta^- n = 63$ 1; $\beta^- 2n = 11$ 7; ...	*
$^{17}\text{C}$	21032	17	193 ms	5	(3/2 <sup>+</sup> )	99 01Ma08	J	1968	$\beta^- = 100$ ; $\beta^- n = 28.4$ 13; $\beta^- 2n$ ?	*
$^{17}\text{N}$	7870	15	4.173 s	0.004	1/2 <sup>-</sup>	99 94Do08	D	1949	$\beta^- = 100$ ; $\beta^- n = 95$ 1; $\beta^- \alpha = 0.0025$ 4	
$^{17}\text{O}$	-808.7635	0.0007	STABLE		5/2 <sup>+</sup>	99		1925	IS=0.038 1	
$^{17}\text{O}^i$	10270.02	0.17	11078.78	0.17	RQ				$\beta^-$ ?; N ?; IT=0.42 14	
$^{17}\text{F}$	1951.70	0.25	64.370 s	0.027	5/2 <sup>+</sup>	99 16Br01	T	1934	$\beta^+ = 100$	*
$^{17}\text{F}^i$	13144.7	1.9	11193.0	1.9	RQ					
$^{17}\text{Ne}$	16500.4	0.4	109.2 ms	0.6	1/2 <sup>-</sup>	99 88Bo39	D	1963	$\beta^+ = 100$ ; $\beta^+ p = 96.0$ 9; $\beta^+ \alpha = 2.7$ 9	
$^{17}\text{Na}$	35170	1000			3/2 <sup>+</sup> #				p ?	
* $^{17}\text{B}$	D : ... ; $\beta^- 3n = 3.5$ 7; $\beta^- 4n = 0.4$ 3									**
* $^{17}\text{C}$	T : average 95Sc03=193(6) 95Re.A=188(10) 86Cu01=202(17)									**
* $^{17}\text{C}$	D : $\beta^- n$ intensity is from 95Re.A									**
* $^{17}\text{F}$	T : average 16Br01=64.402(0.042) 15Gr14=64.347(0.035)									**
$^{18}\text{B}$	51790	200	< 26 ns		(2 <sup>-</sup> )	16		2010	n ?	
$^{18}\text{C}$	24920	30	92 ms	2	0 <sup>+</sup>	96		1969	$\beta^- = 100$ ; $\beta^- n = 31.5$ 15; $\beta^- 2n$ ?	
$^{18}\text{N}$	13113	19	619.2 ms	1.9	1 <sup>-</sup>	96 05Li60	TD	1964	$\beta^- = 100$ ; $\beta^- n = 7.0$ 15; $\beta^- \alpha = 12.2$ 6; $\beta^- 2n$ ?	*
$^{18}\text{O}$	-782.8156	0.0007	STABLE		0 <sup>+</sup>	96		1929	IS=0.205 14	
$^{18}\text{O}^i$	15495	20	16278	20		1 <sup>-</sup> T=2	AHW	E		*
$^{18}\text{F}$	873.1	0.5	109.739 m	0.009	1 <sup>+</sup>	96 14Un01	T	1937	$\beta^+ = 100$	*
$^{18}\text{F}^{m1}$	1994.5	0.5	1121.36	0.15		5 <sup>+</sup>	96		IT=100	
$^{18}\text{F}^i$	1914.7	0.5	1041.55	0.08		0 <sup>+</sup> T=1	96		IT=100	
$^{18}\text{Ne}$	5317.6	0.4	1664.20 ms	0.47	0 <sup>+</sup>	96 15La19	T	1954	$\beta^+ = 100$	*
$^{18}\text{Na}$	25040	90	1.3 zs	0.4	1 <sup>-</sup> #	15 04Ze05	TD	2004	p=?	
* $^{18}\text{N}$	D : $\beta^- \alpha$ intensity from 89Zn04									**
* $^{18}\text{N}$	D : other $\beta^- n$ 94Sc01=2.2(0.4)% 95Re.A=10.9(0.9) 91Re02=14.3(2.0)(same group)									**
* $^{18}\text{N}$	T : average 05Li60=619(2) 99Og03=620(14) 82O101=624(12) 64Ch19=630(30)									**
* $^{18}\text{O}^i$	E : assuming 16399(5), 17025(10) levels to be IAS's of 114.90(0.18), 747(10)									**
* $^{18}\text{O}^i$	E : levels in $^{18}\text{N}$ (see 95Ti07)									**
* $^{18}\text{F}$	T : average 14Un01=109.770(0.018) 10Ga04=109.722(0.012) 04Sc04=109.748(0.021)									**
* $^{18}\text{Ne}$	T : average 15La19=1664.00(+0.57-0.48) 13Gr03=1664.8(1.1)									**
* $^{18}\text{Ne}$	T : 13Gr03 supersedes 07Gr18=1665.6(1.9) of same group									**
$^{19}\text{B}$	59770	530	2.92 ms	0.13	3/2 <sup>-</sup> #	96 03Yo02	TD	1984	$\beta^- = 100$ ; $\beta^- n = 71$ 9; $\beta^- 2n = 17$ 5; $\beta^- 3n < 9.1$	*
$^{19}\text{C}$	32410	100	46.2 ms	2.3	(1/2 <sup>+</sup> )	96 88Du09	TD	1974	$\beta^- = 100$ ; $\beta^- n = 47$ 3; $\beta^- 2n = 7$ 3	*
$^{19}\text{N}$	15856	16	336 ms	3	1/2 <sup>-</sup>	96 06Su12	TJD	1968	$\beta^- = 100$ ; $\beta^- n = 41.8$ 9	
$^{19}\text{O}$	3332.9	2.6	26.470 s	0.006	5/2 <sup>+</sup>	96 13Uj01	T	1936	$\beta^- = 100$	*
$^{19}\text{F}$	-1487.4442	0.0009	STABLE		1/2 <sup>+</sup>	96		1920	IS=100.	
$^{19}\text{F}^i$	6052.2	0.9	7539.6	0.9		5/2 <sup>+</sup> T=3/2	96		IT=100	
$^{19}\text{Ne}$	1752.05	0.16	17.274 s	0.010	1/2 <sup>+</sup>	96 14Br06	T	1939	$\beta^+ = 100$	*
$^{19}\text{Ne}^i$	9253	9	7501	9	RQ	(5/2 <sup>+</sup> ) <sup>+</sup> T=3/2	96 MMC127	J		*
$^{19}\text{Na}$	12929	11	> 1 as		(5/2 <sup>+</sup> )	15 10Mu12	T	1969	p=100	*
$^{19}\text{Mg}$	31830	50	5 ps	3	1/2 <sup>-</sup> #	14 09Mu17	TD	2007	2p=100	*
* $^{19}\text{B}$	D : symmetrized from 71.8(+8.3-9.1)% 16.0(+5.6-4.8)%									**
* $^{19}\text{C}$	T : average 88Du09=49(4) 95Re.A=44(4) 95Oz02=45.5(4.0)									**
* $^{19}\text{C}$	J : from 01Ma08, 99Na27 and 95Ba28									**
* $^{19}\text{O}$	T : average 13Uj01=26.476(0.009) 94It.A=26.464(0.009)									**
* $^{19}\text{Ne}$	T : unweighed average of 14Br06=17.283(0.008), 13Uj01=17.254(0.005),									**
* $^{19}\text{Ne}$	T : 12Tr06=17.262(0.007) and 94Ko.A=17.296(0.005)									**
* $^{19}\text{Ne}$	T : 92Ge08=18.5(0.6) for q=10 <sup>+</sup> (bare ion)									**
* $^{19}\text{Ne}^i$	J : if this is the IAS of $^{19}\text{O}$ ground-state 5/2 <sup>+</sup> ; not yet confirmed									**
* $^{19}\text{Na}$	T : from upper limit of 40 keV, dominated by resolution: <1 eV suggested									**
* $^{19}\text{Mg}$	T : symmetrized from 6(+2-4); supersedes 07Mu15=4.0(1.5) ps									**
$^{20}\text{B}$	68450#	800#							n ?; $\beta^- n$ ?; $\beta^- 2n$ ?	
$^{20}\text{C}$	37500	230	16 ms	3	0 <sup>+</sup>	98 90Mu06	TD	1981	$\beta^- = 100$ ; $\beta^- n = 70$ 11; $\beta^- 2n < 18.6$	*
$^{20}\text{N}$	21770	80	136 ms	3		98 06Su12	TD	1969	$\beta^- = 100$ ; $\beta^- n = 42.9$ 14; $\beta^- 2n$ ?	
$^{20}\text{O}$	3796.2	0.9	13.51 s	0.05	0 <sup>+</sup>	98		1959	$\beta^- = 100$	
$^{20}\text{F}$	-17.463	0.030	11.163 s	0.008	2 <sup>+</sup>	98 98Ti06	T	1935	$\beta^- = 100$	
$^{20}\text{F}^i$	6503	3	6521	3	RQ	0 <sup>+</sup> T=2	98			
$^{20}\text{Ne}$	-7041.9305	0.0016	STABLE			0 <sup>+</sup>	98		1913	IS=90.48 3
$^{20}\text{Ne}^i$	3230.5	2.0	10272.5	2.0	RQ	2 <sup>+</sup> T=1	98		IT=100	
$^{20}\text{Ne}^j$	9690.9	2.8	16732.8	2.8	RQ	0 <sup>+</sup> T=2	98		IT=100	
$^{20}\text{Na}$	6850.6	1.1	447.9 ms	2.3	2 <sup>+</sup>	98 89Cl02	D	1950	$\beta^+ = 100$ ; $\beta^+ \alpha = 25.0$ 4	
$^{20}\text{Na}^i$	13349.0	1.2	6498.4	0.5	p	0 <sup>+</sup> T=2	98		1979	p=100
$^{20}\text{Mg}$	17477.7	1.9	93 ms	5	0 <sup>+</sup>	16		1974	$\beta^+ = 100$ ; $\beta^+ p = 30.3$ 12	
* $^{20}\text{C}$	D : average $\beta^- n$ 03Yo02=65(+19-18)% 90Mu06=72(14)%									**
* $^{20}\text{C}$	T : average 90Mu06=14(+6-5) 95Re.A 16.7(3.5); also 03Yo02=21.8(+15.0-7.4)									**



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>21</sup> B	77330#	900#	<260 ns	3/2 <sup>-</sup> #		04 03Oz01 I		n ?
<sup>21</sup> C	45640#	600#	<30 ns	1/2 <sup>+</sup> #		04 93Po.A I		n ?
<sup>21</sup> N	25230	130	84 ms 7	(1/2 <sup>-</sup> )		15	1970	$\beta^-$ =100; $\beta^-$ n=90.5 42; $\beta^-$ 2n ?
<sup>21</sup> O	8062	12	3.42 s 0.10	(5/2 <sup>+</sup> )		04	1968	$\beta^-$ =100; $\beta^-$ n=0#
<sup>21</sup> F	-47.6	1.8	4.158 s 0.020	5/2 <sup>+</sup>		04	1955	$\beta^-$ =100
<sup>21</sup> Ne	-5731.78	0.04	STABLE	3/2 <sup>+</sup>		04	1928	IS=0.27 1
<sup>21</sup> Ne <sup>i</sup>	3127.4	1.4	8859.2	T=3/2		(3/2,5/2) <sup>+</sup>	04	
<sup>21</sup> Na	-2184.63	0.10	22.422 s 0.010	3/2 <sup>+</sup>		04 15Gr05 T	1940	$\beta^+$ =100
<sup>21</sup> Na <sup>i</sup>	6790	4	8975	4	p	5/2 <sup>+</sup> T=3/2	04	
<sup>21</sup> Mg	10903.8	0.8	118.6 ms 0.5	5/2 <sup>+</sup>		04 15Lu13 D	1963	$\beta^+$ =100; $\beta^+$ p=32.6 10; $\beta^+$ $\alpha$ =?; $\beta^+$ p $\alpha$ =0.016 3
<sup>21</sup> Al	26990#	600#	<35 ns	5/2 <sup>+</sup> #		04 93Po.A I		p ?
<sup>22</sup> C	53610	230	6.2 ms 1.3	0 <sup>+</sup>		15	1986	$\beta^-$ =100; $\beta^-$ n=61 14; $\beta^-$ 2n<37
<sup>22</sup> N	31760	210	23 ms 3	0 <sup>-</sup> #		15	1979	$\beta^-$ =100; $\beta^-$ n=34 3; $\beta^-$ 2n=12 3
<sup>22</sup> O	9280	60	2.25 s 0.09	0 <sup>+</sup>		15	1969	$\beta^-$ =100; $\beta^-$ n<22
<sup>22</sup> F	2793	12	4.23 s 0.04	(4 <sup>+</sup> )		15	1965	$\beta^-$ =100; $\beta^-$ n<11
<sup>22</sup> Ne	-8024.719	0.018	STABLE	0 <sup>+</sup>		15	1913	IS=9.25 3
<sup>22</sup> Ne <sup>i</sup>	5855	10	13880	10		4 <sup>+</sup> T=2	15 87Wi03 E	
<sup>22</sup> Na	-5181.51	0.17	2.6018 y 0.0022	3 <sup>+</sup>		15	1935	$\beta^+$ =100
<sup>22</sup> Na <sup>m</sup>	-4598.46	0.20	583.05	0.10		243 ns 2	15	IT=100
<sup>22</sup> Na <sup>i</sup>	-4524.51	0.22	657.00	0.14		19.6 ps 0.7	15	IT=100
<sup>22</sup> Mg	-399.9	0.3	3.8755 s 0.0012	0 <sup>+</sup>		15	1961	$\beta^+$ =100
<sup>22</sup> Mg <sup>i</sup>	13645	6	14044	6	p	(4 <sup>+</sup> )T=2	15 MMC12 J	$\alpha$ =?; p=?
<sup>22</sup> Al	18200#	400#	91.1 ms 0.5	(4 <sup>+</sup> )		15	1982	$\beta^+$ =100; $\beta^+$ p=55 3; $\beta^+$ 2p=1.10 11; ...
<sup>22</sup> Si	33340#	500#	29 ms 2	0 <sup>+</sup>		15	1987	$\beta^+$ =100; $\beta^+$ p=32 4
<sup>22</sup> C	T : symmetrized from 6.1(+1.4-1.2)		D : symmetrized from $\beta^-$ n=61(+14-13)%					**
<sup>22</sup> Ne <sup>i</sup>	E : 16Ma.A=13880(10) is preferred as IAS but not proven; 90En08=14060(20)							
<sup>22</sup> Mg <sup>i</sup>	J : IAS of <sup>22</sup> Al and <sup>22</sup> F ground-state							
<sup>22</sup> Al	D : ...; $\beta^+$ $\alpha$ =0.038 17							
<sup>23</sup> C	64170#	1000#				3/2 <sup>+</sup> #		n ?
<sup>23</sup> N	36720	420	13.9 ms 1.4	1/2 <sup>-</sup> #		07 03Yo02 TD	1985	$\beta^-$ =100; $\beta^-$ n=42 6; $\beta^-$ 2n=8 4; $\beta^-$ 3n<3.4
<sup>23</sup> O	14620	120	97 ms 8	1/2 <sup>+</sup>		07 07Su05 TD	1970	$\beta^-$ =100; $\beta^-$ n=7 2
<sup>23</sup> F	3290	30	2.23 s 0.14	5/2 <sup>+</sup>		07 95Re.A D	1970	$\beta^-$ =100; $\beta^-$ n<14
<sup>23</sup> Ne	-5154.05	0.10	37.140 s 0.028	5/2 <sup>+</sup>		07 15La19 T	1936	$\beta^-$ =100
<sup>23</sup> Na	-9529.8525	0.0018	STABLE	3/2 <sup>+</sup>		07	1921	IS=100.
<sup>23</sup> Na <sup>i</sup>	-1638.66	0.15	7891.19	0.15		5/2 <sup>+</sup> T=3/2	07	IT=100
<sup>23</sup> Na <sup>j</sup>	10060.6	2.0	19590.4	2.0		T=5/2	85Ev01 T	
<sup>23</sup> Mg	-5473.51	0.16	11.317 s 0.011	3/2 <sup>+</sup>		07	1939	$\beta^+$ =100
<sup>23</sup> Mg <sup>i</sup>	2328.7	1.4	7802.2	1.4		5/2 <sup>+</sup> T=3/2	07 00Pe28 D	IT $\approx$ 100; p=0.17 8
<sup>23</sup> Al	6748.1	0.3	470 ms 30	5/2 <sup>+</sup>		07	1969	$\beta^+$ =100; $\beta^+$ p=0.46 23
<sup>23</sup> Al <sup>i</sup>	18530	60	11780	60	p	(5/2 <sup>+</sup> )T=5/2	07	p=0.10 5; 2p=3.6 4
<sup>23</sup> Si	23700#	500#	42.3 ms 0.4	3/2 <sup>+</sup> #		07 97Bi04 TD	1986	$\beta^+$ =100; $\beta^+$ p $\approx$ 88; $\beta^+$ 2p=3.6 3
<sup>23</sup> N	T : symmetrized from 14.1(+1.2-1.5)							
<sup>23</sup> N	D : symmetrized from 42.2(+6.3-6.5)% 8.0(+3.8-3.4)%							
<sup>23</sup> Ne	T : average 15La19=37.148(0.032) 07Gr18=37.11(0.06); other 74Al03=37.24(0.12)							
<sup>23</sup> Na <sup>j</sup>	T : from width=1.9(0.8) keV							
<sup>24</sup> N	46940#	400#	<52 ns			07 93Po.A I		n ?
<sup>24</sup> O	18500	160	77.4 ms 4.5	0 <sup>+</sup>		07 15Ca09 TD	1970	$\beta^-$ =100; $\beta^-$ n=43 4
<sup>24</sup> F	7540	100	384 ms 16	3 <sup>+</sup>		07 07Su05 T	1970	$\beta^-$ =100; $\beta^-$ n<5.9
<sup>24</sup> Ne	-5951.6	0.5	3.38 m 0.02	0 <sup>+</sup>		07	1956	$\beta^-$ =100
<sup>24</sup> Na	-8417.901	0.017	14.957 h 0.004	4 <sup>+</sup>		07 14Un01 T	1934	$\beta^-$ =100
<sup>24</sup> Na <sup>m</sup>	-7945.694	0.017	472.2074	0.0008		1 <sup>+</sup>	07	IT $\approx$ 100; $\beta^-$ =0.05
<sup>24</sup> Na <sup>i</sup>	-2450.53	0.13	5967.37	0.13		0 <sup>+</sup> T=2	07	
<sup>24</sup> Mg	-13933.569	0.013	STABLE	0 <sup>+</sup>		07	1920	IS=78.99 4
<sup>24</sup> Mg <sup>i</sup>	-4417.29	0.04	9516.28	0.04		(4 <sup>+</sup> )T=1	07	
<sup>24</sup> Mg <sup>j</sup>	1502.8	0.6	15436.4	0.6		0 <sup>+</sup> T=2	07	
<sup>24</sup> Al	-48.86	0.23	2.053 s 0.004	4 <sup>+</sup>		07	1953	$\beta^+$ =100; $\beta^+$ $\alpha$ =0.035 6; $\beta^+$ p=0.0016 3
<sup>24</sup> Al <sup>m</sup>	376.94	0.25	425.8	0.1		1 <sup>+</sup>	07	IT=82.5 30; $\beta^+$ =17.5 30; $\beta^+$ $\alpha$ =0.028 6
<sup>24</sup> Al <sup>i</sup>	5900	3	5949	3	p	0 <sup>+</sup> T=2	07	
<sup>24</sup> Si	10745	19	140 ms 8	0 <sup>+</sup>		07 98Cz01 D	1979	$\beta^+$ =100; $\beta^+$ p=37.6 25
<sup>24</sup> P	33320#	500#		1 <sup>+</sup> #				p ?; $\beta^+$ ?; $\beta^+$ p ?
<sup>24</sup> O	T : average 15Ca09=80(5) 01Pe14=67(10)							
<sup>24</sup> F	J : 15Ca09=3 <sup>+</sup>							
<sup>24</sup> Na	T : average 14Un01=14.955(0.007) 83Wa26=14.958(0.005) and							
<sup>24</sup> Na	T : 80Ho17=14.9590(0.0150); all values are from standard labs							
<sup>24</sup> Na	T : original unc of 80Ho17=0.0012 increased to 0.1% by evaluator							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
<sup>25</sup> N	55980#	500#	<260 ns	1/2 <sup>-</sup> #	09	99Sa06	ID	n ?; 2n ?; β <sup>-</sup> ?	*		
<sup>25</sup> O	27330	170	5.18 zs	0.35	3/2 <sup>+</sup> #	09	16Ko11	T	2008	n=100	*
<sup>25</sup> F	11330	100	80 ms	9	(5/2 <sup>+</sup> )	09			1970	β <sup>-</sup> =100; β <sup>-</sup> n=23.1 45; β <sup>-</sup> 2n=0#	
<sup>25</sup> Ne	-2036	29	602 ms	8	1/2 <sup>+</sup>	09			1970	β <sup>-</sup> =100	
<sup>25</sup> Na	-9357.8	1.2	59.1 s	0.6	5/2 <sup>+</sup>	09			1943	β <sup>-</sup> =100	
<sup>25</sup> Mg	-13192.78	0.05	STABLE		5/2 <sup>+</sup>	09			1920	IS=10.00 1	
<sup>25</sup> Mg <sup>i</sup>	-5405.8	0.3	7787.0	0.3	5/2 <sup>+</sup> T=3/2	09					
<sup>25</sup> Al	-8915.97	0.06	7.183 s	0.012	5/2 <sup>+</sup>	09			1953	β <sup>+</sup> =100	
<sup>25</sup> Al <sup>i</sup>	-1014.9	1.8	7901.1	1.8	RQ	5/2 <sup>+</sup> T=3/2	09				
<sup>25</sup> Si	3827	10	220 ms	3	5/2 <sup>+</sup>	09			1963	β <sup>+</sup> =100; β <sup>+</sup> p=35 2	
<sup>25</sup> P	19740#	400#	<30 ns		1/2 <sup>+</sup> #	09	93Po.A	I		p ?	
* <sup>25</sup> N	D : in 99Sa06 experiment, 240 <sup>25</sup> N events expected, none observed									**	
* <sup>25</sup> O	T : from decay width 16Ko11=88(6) keV; other 13Ca18=20(+60-20) keV									**	
<sup>26</sup> O	34660	160	4.2 ps	3.3	0 <sup>+</sup>	16	13Ko10	T	2012	2n=100	*
<sup>26</sup> F	18650	110	8.2 ms	0.9	1 <sup>+</sup>	16			1979	β <sup>-</sup> =100; β <sup>-</sup> n=13.5 40; β <sup>-</sup> 2n=0.4#	
<sup>26</sup> F <sup>m</sup>	19290	110	2.2 ms	0.1	(4 <sup>+</sup> )	16			2013	IT=82 11; β <sup>-</sup> =?; β <sup>-</sup> n=12 8	
<sup>26</sup> Ne	481	18	197 ms	2	0 <sup>+</sup>	16			1970	β <sup>-</sup> =100; β <sup>-</sup> n=0.13 3	
<sup>26</sup> Na	-6861	4	1071.28 ms	0.25	3 <sup>+</sup>	16			1958	β <sup>-</sup> =100	
<sup>26</sup> Na <sup>m</sup>	-6779	4	9 μs	2	1 <sup>+</sup>	16			1987	IT=100	
<sup>26</sup> Mg	-16214.542	0.030	STABLE		0 <sup>+</sup>	16			1920	IS=11.01 3	
<sup>26</sup> Al	-12210.15	0.07	717 ky	24	5 <sup>+</sup>	16			1934	β <sup>+</sup> =100	
<sup>26</sup> Al <sup>m</sup>	-11981.85	0.07	228.306	0.013	MD	6346.0 ms	0.8	0 <sup>+</sup> T=1	16	1934	β <sup>+</sup> =100
<sup>26</sup> Si	-7141.02	0.11	2.2453 s	0.0007	0 <sup>+</sup>	16			1960	β <sup>+</sup> =100	
<sup>26</sup> Si <sup>i</sup>	5926	11	13068	11	p	(3 <sup>+</sup> )T=2	16				
<sup>26</sup> P	10970#	200#	43.7 ms	0.6	(3 <sup>+</sup> )	16			1983	β <sup>+</sup> =100; β <sup>+</sup> p=36.8 20; ...	*
<sup>26</sup> P <sup>m</sup>	11130#	200#	120 ns	9	0 <sup>+</sup>	16			2014	IT=100	
<sup>26</sup> S	27080#	600#	<79 ns		0 <sup>+</sup>	16				2p ?	
* <sup>26</sup> O	T : symmetrized from 13Ko10=4.5(+1.1-1.5 stat)(3 systematics)									**	
* <sup>26</sup> P	D : ...; β <sup>+</sup> 2p=2.16 24 D : β <sup>+</sup> p + β <sup>+</sup> 2p=39(2)									**	
<sup>27</sup> O	44670#	500#	<260 ns		3/2 <sup>+</sup> #	99	Sa06	I		n ?; 2n ?	
<sup>27</sup> F	25450	390	4.9 ms	0.2	5/2 <sup>+</sup> #	11	98No.A	T	1981	β <sup>-</sup> =100; β <sup>-</sup> n=77 21; β <sup>-</sup> 2n=5#	*
<sup>27</sup> Ne	7050	90	31.5 ms	1.3	(3/2 <sup>+</sup> )	11			1977	β <sup>-</sup> =100; β <sup>-</sup> n=2.0 5; β <sup>-</sup> 2n=0#	
<sup>27</sup> Na	-5518	4	301 ms	6	5/2 <sup>+</sup>	11			1968	β <sup>-</sup> =100; β <sup>-</sup> n=0.13 4	
<sup>27</sup> Mg	-14586.61	0.05	9.435 m	0.027	1/2 <sup>+</sup>	11	15ZaZY	T	1934	β <sup>-</sup> =100	*
<sup>27</sup> Al	-17196.86	0.05	STABLE		5/2 <sup>+</sup>	11			1922	IS=100.	
<sup>27</sup> Al <sup>i</sup>	-10383.1	0.7	6813.8	0.7		1/2 <sup>+</sup> T=3/2	11			IT=100	
<sup>27</sup> Si	-12384.50	0.11	4.15 s	0.04	5/2 <sup>+</sup>	11			1939	β <sup>+</sup> =100	
<sup>27</sup> Si <sup>i</sup>	-5759.5	2.3	6625.0	2.3	RQ	1/2 <sup>+</sup> T=3/2	11		1977	IT ?	
<sup>27</sup> P	-722	26	260 ms	80	1/2 <sup>+</sup>	11			1977	β <sup>+</sup> =100; β <sup>+</sup> p=0.07	
<sup>27</sup> P <sup>i</sup>	12010	30	12730	40	p	5/2 <sup>+</sup> T=5/2	11		1991	IT ?	
<sup>27</sup> S	17030#	400#	15.5 ms	1.5	(5/2 <sup>+</sup> )	11			1986	β <sup>+</sup> =100; β <sup>+</sup> p=2.3 9; β <sup>+</sup> 2p=1.1 5	
* <sup>27</sup> F	T : others not used: 99Re16=6.5(1.1) 97Ta22=5.3(0.9) outweighed; and									**	
* <sup>27</sup> F	T : 99Di01=5.2(0.3) same data as in 99Re16									**	
* <sup>27</sup> Mg	T : average 15ZaZY=9.408 (0.012) 70Re13=9.462 (0.012); Birge ratio=3.18									**	
<sup>28</sup> O	52080#	700#	<100 ns		0 <sup>+</sup>	13	98Po.A	I		n ?; 2n ?; β <sup>-</sup> =0	*
<sup>28</sup> F	33740	390	46 zs			13				n ?	
<sup>28</sup> Ne	11300	130	20 ms	1	0 <sup>+</sup>	13			1979	β <sup>-</sup> =100; β <sup>-</sup> n=12 1; β <sup>-</sup> 2n=3.7 5	
<sup>28</sup> Na	-988	10	30.5 ms	0.4	1 <sup>+</sup>	13			1969	β <sup>-</sup> =100; β <sup>-</sup> n=0.58 12	
<sup>28</sup> Mg	-15018.8	2.0	20.915 h	0.009	0 <sup>+</sup>	13			1953	β <sup>-</sup> =100	
<sup>28</sup> Al	-16850.64	0.08	2.245 m	0.005	3 <sup>+</sup>	13			1934	β <sup>-</sup> =100	
<sup>28</sup> Al <sup>i</sup>	-10858.06	0.13	5992.58	0.10		0 <sup>+</sup> T=2	13				
<sup>28</sup> Si	-21492.7943	0.0005	STABLE		0 <sup>+</sup>	13			1920	IS=92.223 19	
<sup>28</sup> Si <sup>r</sup>	-8951.75	0.05	12541.04	0.05	RQ	(3 <sup>+</sup> )	13				
<sup>28</sup> Si <sup>i</sup>	-12176.87	0.10	9315.92	0.10		3 <sup>+</sup> T=1	13				
<sup>28</sup> Si <sup>j</sup>	-6265.8	1.0	15227	1		(0 <sup>+</sup> )T=2	13	68Mc12	D	1968	α=90 11; p=10 11
<sup>28</sup> P	-7147.7	1.2	270.3 ms	0.5		3 <sup>+</sup>	13	79Ho27	D	1953	β <sup>+</sup> =100; β <sup>+</sup> p=0.0013 4; β <sup>+</sup> α=0.00086 25
<sup>28</sup> P <sup>i</sup>	-1261	20	5887	20	p	0 <sup>+</sup> T=2	13				
<sup>28</sup> S	4070	160	125 ms	10	0 <sup>+</sup>	13			1982	β <sup>+</sup> =100; β <sup>+</sup> p=20.7 19	
<sup>28</sup> Cl	27520#	600#			1 <sup>+</sup> #					p ?	
* <sup>28</sup> O	D : in 97Ta22 and 99Sa06, 11 and 37 <sup>28</sup> O events expected, none observed									**	

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>29</sup> F	40150	530	2.5 ms 0.3	5/2 <sup>+</sup> #	12	99DI01	D	1989	$\beta^-$ =100; $\beta^-$ n=60 40; $\beta^-$ 2n=5#	*
<sup>29</sup> Ne	18400	150	14.7 ms 0.4	(3/2 <sup>-</sup> )	12	05Tr13	T	1985	$\beta^-$ =100; $\beta^-$ n=28 5; $\beta^-$ 2n=4 1	*
<sup>29</sup> Na	2680	7	44.1 ms 0.9	3/2 <sup>(+)</sup> #	12	95Re.A	D	1969	$\beta^-$ =100; $\beta^-$ n=25.9 23; $\beta^-$ 2n=0#	*
<sup>29</sup> Mg	-10603	11	1.30 s 0.12	3/2 <sup>+</sup>	12			1971	$\beta^-$ =100	
<sup>29</sup> Al	-18207.8	0.3	6.56 m 0.06	5/2 <sup>+</sup>	12			1939	$\beta^-$ =100	
<sup>29</sup> Si	-21895.0784	0.0006	STABLE	1/2 <sup>+</sup>	12			1920	IS=4.685 8	
<sup>29</sup> Si <sup>i</sup>	-13605	5	8290 5	5/2 <sup>+</sup> T=3/2	12				IT=100	
<sup>29</sup> P	-16952.8	0.4	4.142 s 0.015	1/2 <sup>+</sup>	12			1941	$\beta^+$ =100	
<sup>29</sup> P <sup>i</sup>	-8571.0	2.5	8381.8 2.4 RQ	5/2 <sup>+</sup> T=3/2	12			1969	IT=100	
<sup>29</sup> S	-3160	50	188 ms 4	5/2 <sup>+</sup> #	12	79Vi01	D	1964	$\beta^+$ =100; $\beta^+$ p=46.4 10	
<sup>29</sup> Cl	13160	190	<10ps	(1/2 <sup>+</sup> )	16	15Mu13	I		p=100	
* <sup>29</sup> F	D : $\beta^-$ n from 99DI01=100(80)%									**
* <sup>29</sup> Ne	T : average 05Tr13=13.8(0.5) 97No.A=15.6(0.5); others outweighed, not used:									**
* <sup>29</sup> Ne	T : 06Tr02=15.1(2.6) 16.4(1.3) 99DI01=15(4) 99Re16=19(9) 97Ta22=15(3)									**
* <sup>29</sup> Ne	J : 16Ko05=(3/2 <sup>-</sup> )									**
* <sup>29</sup> Na	D : $\beta^-$ n: average 95Re.A=27.1(1.6)% 84La03=21.5(3.0)%									**
<sup>30</sup> F	48110#	600#	<260 ns		10	99Sa06	I		n ?	
<sup>30</sup> Ne	23280	250	7.22 ms 0.18	0 <sup>+</sup>	10	15St14	T	1985	$\beta^-$ =100; $\beta^-$ n=13 4; $\beta^-$ 2n=8.9 23	*
<sup>30</sup> Na	8475	5	48.4 ms 1.7	2 <sup>+</sup>	10	99DI01	T	1969	$\beta^-$ =100; $\beta^-$ n=30 4; ...	*
<sup>30</sup> Mg	-8884	3	313 ms 4	0 <sup>+</sup>	10	84La03	D	1971	$\beta^-$ =100; $\beta^-$ n<0.06	*
<sup>30</sup> Al	-15864.8	2.9	3.62 s 0.06	3 <sup>+</sup>	10			1961	$\beta^-$ =100	
<sup>30</sup> Si	-24432.960	0.022	STABLE	0 <sup>+</sup>	10			1924	IS=3.092 11	
<sup>30</sup> P	-20200.85	0.07	2.498 m 0.004	1 <sup>+</sup> T=0	10			1934	$\beta^+$ =100	*
<sup>30</sup> P <sup>i</sup>	-19523.84	0.08	677.01 0.03	0 <sup>+</sup> T=1	10					
<sup>30</sup> S	-14059.25	0.21	1.1759 s 0.0017	0 <sup>+</sup>	10	11So11	T	1961	$\beta^+$ =100	
<sup>30</sup> Cl	4440#	200#	<30 ns	3 <sup>+</sup> #	10	93Po.A	I		p ?	
<sup>30</sup> Ar	20930	210	<10ps	0 <sup>+</sup>	16			2015	2p=100	
* <sup>30</sup> Ne	T : average 15St14=7.18(0.22) 07Tr08=7.3(0.3)									**
* <sup>30</sup> Na	D : ... ; $\beta^-$ 2n=1.15 25; $\beta^-$ $\alpha$ =5.5e-5 2									**
* <sup>30</sup> Na	T : average 99DI01=50(4) 97Ta22=48(5) 84La02=48(2)									**
* <sup>30</sup> Mg	T : average 08Hi05=314(5) and 312(7)									**
* <sup>30</sup> P	D : first observed radionuclide, in 1934									**
<sup>31</sup> F	56140#	550#	1# ms (>260 ns)	5/2 <sup>+</sup> #	13			1999	$\beta^-$ ?; $\beta^-$ n=10#; $\beta^-$ 2n=0#	
<sup>31</sup> Ne	31180	270	3.4 ms 0.8	(3/2 <sup>-</sup> )	13			1996	$\beta^-$ =100; $\beta^-$ n=10#; $\beta^-$ 2n=4#	
<sup>31</sup> Na	12246	14	17.35 ms 0.40	3/2 <sup>(+)</sup> #	13	93KI02	J	1969	$\beta^-$ =100; $\beta^-$ n=37.3 54; ...	*
<sup>31</sup> Mg	-3122	3	236 ms 20	1/2 <sup>(+)</sup>	13			1977	$\beta^-$ =100; $\beta^-$ n=6.2 19	*
<sup>31</sup> Al	-14950.7	2.2	644 ms 25	5/2 <sup>(+)</sup>	13			1971	$\beta^-$ =100; $\beta^-$ n<1.6	
<sup>31</sup> Si	-22949.04	0.04	157.36 m 0.26	3/2 <sup>+</sup>	13			1934	$\beta^-$ =100	
<sup>31</sup> P	-24440.5410	0.0007	STABLE	1/2 <sup>+</sup>	13			1920	IS=100.	
<sup>31</sup> P <sup>i</sup>	-18059.7	2.0	6380.8 2.0	3/2 <sup>+</sup> T=3/2	13				IT=100	
<sup>31</sup> S	-19042.52	0.23	2.5534 s 0.0018	1/2 <sup>+</sup>	13			1940	$\beta^+$ =100	
<sup>31</sup> Si <sup>i</sup>	-12761.9	0.6	6280.60 0.60	3/2 <sup>+</sup> T=3/2	13					
<sup>31</sup> Cl	-7035	3	190 ms 1	3/2 <sup>+</sup>	13			1977	$\beta^+$ =100; $\beta^+$ p=2.4 2	
<sup>31</sup> Cl <sup>i</sup>	5256	3	12291 5 RQ	3/2 <sup>+</sup> T=5/2						
<sup>31</sup> Ar	11330#	200#	15.1 ms 0.3	5/2 <sup>+</sup>	13	14Ko17	T	1986	$\beta^+$ =100; $\beta^+$ p=68.3 3; $\beta^+$ 2p=9.0 2; ...	*
* <sup>31</sup> Na	D : ... ; $\beta^-$ 2n=0.87 24; $\beta^-$ 3n<0.05									**
* <sup>31</sup> Mg	D : strongly conflicting with earlier 84La03=1.7(0.3)%									**
* <sup>31</sup> Ar	D : ... ; $\beta^+$ $\alpha$ <0.38; $\beta^+$ 3p=0.07 2; $\beta^+$ $\alpha$ <0.03; 2p<0.0006									**
<sup>32</sup> Ne	37000#	500#	3.5 ms 0.9	0 <sup>+</sup>	11			1990	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=7#	
<sup>32</sup> Na	18640	40	12.9 ms 0.3	(3 <sup>-</sup> )	11	08Tr04	TJ	1972	$\beta^-$ =100; $\beta^-$ n=24 7; $\beta^-$ 2n=8 2	*
<sup>32</sup> Mg	-829	3	86 ms 5	0 <sup>+</sup>	11			1977	$\beta^-$ =100; $\beta^-$ n=5.5 5	
<sup>32</sup> Al	-11099	7	33.0 ms 0.2	1 <sup>+</sup>	11			1971	$\beta^-$ =100; $\beta^-$ n=0.7 5	
<sup>32</sup> Al <sup>m</sup>	-10142	7	956.6 0.5	(4 <sup>+</sup> )	11			1996	IT=100	
<sup>32</sup> Si	-24077.69	0.30	153 y 19	0 <sup>+</sup>	11			1953	$\beta^-$ =100	
<sup>32</sup> P	-24304.87	0.04	14.268 d 0.005	1 <sup>+</sup>	11			1934	$\beta^-$ =100	*
<sup>32</sup> P <sup>i</sup>	-19232.43	0.07	5072.44 0.06	0 <sup>+</sup> T=2	11				IT=100	
<sup>32</sup> S	-26015.5336	0.0013	STABLE	0 <sup>+</sup>	11			1920	IS=94.99 26	
<sup>32</sup> Si <sup>i</sup>	-19014.1	0.4	7001.4 0.4	1 <sup>+</sup> T=1	11				IT=100	
<sup>32</sup> Si <sup>j</sup>	-13967.57	0.28	12047.96 0.28	0 <sup>+</sup> T=2	11				IT=100	
<sup>32</sup> Cl	-13334.7	0.6	298 ms 1	1 <sup>+</sup>	11			1953	$\beta^+$ =100; $\beta^+$ $\alpha$ =0.054 8; $\beta^+$ p=0.026 5	
<sup>32</sup> Cl <sup>i</sup>	-8288.4	0.7	5046.3 0.3	0 <sup>+</sup> T=2	11				IT=100	
<sup>32</sup> Ar	-2200.4	1.8	98 ms 2	0 <sup>+</sup>	11			1977	$\beta^+$ =100; $\beta^+$ p=35.58 0.22	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>32</sup> K	21100#	400#		1+#				p ?
<sup>32</sup> K <sup>m</sup>	22050#	410#	950# 100#	4+#		Mirror I		p ?
* <sup>32</sup> Na	T : average 08Tr04=13.1(0.5) and 11.5(1.2) 98No.A=11.5(0.8) 84La03=13.2(0.4)							
* <sup>32</sup> P	T : also 14Un01=14.263(0.035)							
<sup>33</sup> Ne	46000#	600#		<260 ns	7/2 <sup>-</sup> #	11 02No11 I		n ?
<sup>33</sup> Na	23780	450		8.2 ms 0.4	(3/2 <sup>+</sup> )	11	1972	$\beta^-$ =100; $\beta^-$ n=47 6; $\beta^-$ 2n=13 3
<sup>33</sup> Mg	4962.3	2.9		90.5 ms 1.6	3/2 <sup>-</sup>	11	1979	$\beta^-$ =100; $\beta^-$ n=14 2; $\beta^-$ 2n=3#
<sup>33</sup> Al	-8497	7		41.7 ms 0.2	5/2 <sup>+</sup>	11 06Hi18 J	1971	$\beta^-$ =100; $\beta^-$ n=8.5 7
<sup>33</sup> Si	-20514.3	0.7		6.18 s 0.18	3/2 <sup>+</sup>	11	1971	$\beta^-$ =100
<sup>33</sup> P	-26337.3	1.1		25.35 d 0.11	1/2 <sup>+</sup>	11	1951	$\beta^-$ =100
<sup>33</sup> S	-26585.8543	0.0014		STABLE	3/2 <sup>+</sup>	11	1926	IS=0.75 2
<sup>33</sup> S <sup>i</sup>	-21106.06	0.13	5479.79	0.13		1/2 <sup>+</sup> T=3/2	11	IT=100
<sup>33</sup> Cl	-21003.3	0.4		2.5038 s 0.0022	3/2 <sup>+</sup>	11 15Gr14 T	1940	$\beta^+$ =100
<sup>33</sup> Cl <sup>i</sup>	-15454.9	0.5	5548.4	0.4	RQ	1/2 <sup>+</sup> T=3/2	11	IT=100
<sup>33</sup> Ar	-9384.3	0.4		173.0 ms 2.0	1/2 <sup>+</sup>	11	1964	$\beta^+$ =100; $\beta^+$ p=38.7 10
<sup>33</sup> K	7040#	200#		<25 ns	3/2 <sup>+</sup> #	11 93Po.A I		p ?
* <sup>33</sup> Ne	T : estimated half-life 1# ms for $\beta^-$ decay I : also 02Le.A < 1.5 $\mu$ s							
<sup>34</sup> Ne	52840#	510#		1# ms (>1.5 $\mu$ s)	0 <sup>+</sup>	12 02Le.A I	2002	$\beta^-$ ?; $\beta^-$ 2n=40#; $\beta^-$ n=1#
<sup>34</sup> Na	31680	600		5.5 ms 1.0	1 <sup>+</sup>	12 GAu03 D	1983	$\beta^-$ =100; $\beta^-$ 2n $\approx$ 50; $\beta^-$ n $\approx$ 15
<sup>34</sup> Mg	8323	29		20 ms 10	0 <sup>+</sup>	12	1979	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=0.4#
<sup>34</sup> Al	-3000	3		56.3 ms 0.5	(4 <sup>-</sup> )	12	1977	$\beta^-$ =100; $\beta^-$ n=26 4; $\beta^-$ 2n=0.4#
<sup>34</sup> Al <sup>m</sup>	-2450#	100#	550# 100#	26 ms 1	(1 <sup>+</sup> )	12Ro25 TJ	2012	$\beta^-$ $\approx$ 100; $\beta^-$ n=30#; $\beta^-$ 2n=0.4#
<sup>34</sup> Si	-19957	14		2.77 s 0.20	0 <sup>+</sup>	12	1971	$\beta^-$ =100
<sup>34</sup> Si <sup>m</sup>	-15701	14	4256.1	0.4		< 210 ns	12	IT=100
<sup>34</sup> P	-24548.7	0.8		12.43 s 0.10	1 <sup>+</sup>	12	1945	$\beta^-$ =100
<sup>34</sup> S	-29931.69	0.04		STABLE	0 <sup>+</sup>	12	1926	IS=4.25 24
<sup>34</sup> Cl	-24440.08	0.05		1.5266 s 0.0004	0 <sup>+</sup> T=1	12	1934	$\beta^+$ =100
<sup>34</sup> Cl <sup>m</sup>	-24293.72	0.05	146.360	0.027	MD	3 <sup>+</sup> T=0	12	$\beta^+$ =55.4 6; IT=44.6 6
<sup>34</sup> Ar	-18378.29	0.08		843.8 ms 0.4	0 <sup>+</sup>	12	1966	$\beta^+$ =100
<sup>34</sup> Ar <sup>i</sup>	-10444	5	7934	5	RQ	1 <sup>+</sup> #T=2	12	$\beta^+$ ?; IT ?
<sup>34</sup> K	-1220#	200#		<40 ns	1 <sup>+</sup> #	12 93Po.A I		p ?
<sup>34</sup> Ca	13850#	300#		<35 ns	0 <sup>+</sup>	12 93Po.A I		2p ?
* <sup>34</sup> Na	D : $\beta^-$ n $\approx$ 15%, $\beta^-$ 2n $\approx$ 50% estimated from $P_n = \beta^-n + 2 \times \beta^-2n = 115(20)\%$ in 84La03							
* <sup>34</sup> Na	D : assuming $\beta^-n/\beta^-2n=0.3$ from trends in the <sup>30</sup> Na- <sup>33</sup> Na series: 26 41 3 4							
<sup>35</sup> Na	38230#	670#		1.5 ms 0.5	3/2 <sup>+</sup> #	11	1983	$\beta^-$ =100; $\beta^-$ n=60#; $\beta^-$ 2n=10#
<sup>35</sup> Mg	15640	270		70 ms 40	7/2 <sup>-</sup> #	11	1989	$\beta^-$ =100; $\beta^-$ n=52 46; $\beta^-$ 2n=20#
<sup>35</sup> Al	-224	7		37.2 ms 0.8	5/2 <sup>+</sup> #	11	1979	$\beta^-$ =100; $\beta^-$ n=38 2; $\beta^-$ 2n=0.2#
<sup>35</sup> Si	-14390	40		780 ms 120	7/2 <sup>-</sup> #	15 95Re.A D	1971	$\beta^-$ =100; $\beta^-$ n<5
<sup>35</sup> P	-24857.8	1.9		47.3 s 0.8	1/2 <sup>+</sup>	11	1971	$\beta^-$ =100
<sup>35</sup> S	-28846.21	0.04		87.37 d 0.04	3/2 <sup>+</sup>	11	1936	$\beta^-$ =100
<sup>35</sup> S <sup>i</sup>	-19691	10	9155	10	RQ	T=5/2	11	1975
<sup>35</sup> Cl	-29013.53	0.04		STABLE	3/2 <sup>+</sup>	11	1919	IS=75.76 10
<sup>35</sup> Cl <sup>i</sup>	-23359.05	0.22	5654.48	0.22		3/2 <sup>+</sup> T=3/2	11	IT=100
<sup>35</sup> Ar	-23047.3	0.7		1.7756 s 0.0010	3/2 <sup>+</sup>	11	1940	$\beta^+$ =100
<sup>35</sup> Ar <sup>i</sup>	-17474.6	0.7	5572.66	0.15		3/2 <sup>+</sup> T=3/2	11	IT=100
<sup>35</sup> K	-11172.9	0.5		178 ms 8	3/2 <sup>+</sup>	11 06Me04 J	1976	$\beta^+$ =100; $\beta^+$ p=0.37 15
<sup>35</sup> K <sup>i</sup>	-2110	40	9060	40	2p	3/2 <sup>+</sup> T=5/2		
<sup>35</sup> Ca	4790#	200#		25.7 ms 0.2	1/2 <sup>+</sup> #	11	1985	$\beta^+$ =100; $\beta^+$ p=95.9 14; $\beta^+$ 2p=4.1 6
* <sup>35</sup> Na	D : $\beta^-$ n has been observed by 83La12 but not quantified							
<sup>36</sup> Na	46300#	680#		<180 ns		12		n ?
<sup>36</sup> Mg	20380	690		3.9 ms 1.3	0 <sup>+</sup>	12	1989	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=3#
<sup>36</sup> Al	5950	150		90 ms 40		12	1979	$\beta^-$ =100; $\beta^-$ n<30; $\beta^-$ 2n=7#
<sup>36</sup> Si	-12440	70		450 ms 60	0 <sup>+</sup>	12 95Re.A D	1971	$\beta^-$ =100; $\beta^-$ n=12 5
<sup>36</sup> P	-20251	13		5.6 s 0.3	4 <sup>-</sup>	12 15Ch56 J	1971	$\beta^-$ =100; $\beta^-$ n=0#
<sup>36</sup> S	-30664.13	0.19		STABLE	0 <sup>+</sup>	12	1938	IS=0.01 1
<sup>36</sup> Cl	-29522.01	0.04		301.3 ky 1.5	2 <sup>+</sup>	12	1941	$\beta^-$ =98.1 1; $\beta^+$ =1.9 1
<sup>36</sup> Cl <sup>i</sup>	-25222.34	0.04	4299.667	0.014		(0 <sup>+</sup> )T=2	12	IT=100
<sup>36</sup> Ar	-30231.540	0.027		STABLE	0 <sup>+</sup>	12	1920	IS=0.3336 21; 2 $\beta^+$ ?
<sup>36</sup> Ar <sup>i</sup>	-23620.5	0.3	6611.0	0.3		2 <sup>+</sup> T=1	12	IT=100
<sup>36</sup> Ar <sup>j</sup>	-19379.4	1.2	10852.2	1.2	RQ	0 <sup>+</sup> T=2	12	IT=100
<sup>36</sup> K	-17417.1	0.3		341 ms 3	2 <sup>+</sup>	12	1967	$\beta^+$ =100; $\beta^+$ p=0.048 14; $\beta^+$ $\alpha$ =0.0034 13
<sup>36</sup> K <sup>i</sup>	-13134.5	2.4	4282.6	2.4	p	0 <sup>+</sup> T=2	12	p=100
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...										
<sup>36</sup> Ca	-6450	40	101.2 ms	1.5	0 <sup>+</sup>	12 07Do17	T 1977	$\beta^+=100$ ; $\beta^+p=51.2$ 10	*	
<sup>36</sup> Sc	15350#	300#						p ?	**	
* <sup>36</sup> K <sup>i</sup>	E : ENSDF2012 finds 4281.9(0.8) as IAS of <sup>36</sup> Ca ground-state									
* <sup>36</sup> Ca	T : average 07Do17=100.1(2.3) 95Tr02=102(2)									
<sup>37</sup> Na	53530#	690#	1# ms	(>1.5 $\mu$ s)	3/2 <sup>+</sup> #	12 02Le.A	I 2002	$\beta^-$ ?; $\beta^-n=100$ #; $\beta^-2n=50$ #		
<sup>37</sup> Mg	28210	700	8 ms	4	(3/2 <sup>-</sup> )	12 14Ko14	J 1996	$\beta^-$ ?; $\beta^-n=80$ #; $\beta^-2n=20$ #		
<sup>37</sup> Al	9810	180	11.5 ms	0.4	5/2 <sup>+</sup> #	12 15St14	TD 1979	$\beta^-n=100$ ; $\beta^-n>29$ 3; $\beta^-2n>1$ 1		
<sup>37</sup> Si	-6570	110	90 ms	60	7/2 <sup>-</sup> #	15		$\beta^-n=100$ ; $\beta^-n=17$ 13; $\beta^-2n=0.2$ #		
<sup>37</sup> P	-19000	40	2.31 s	0.13	(1/2 <sup>+</sup> )	12 15Ch56	J 1971	$\beta^-n=100$ ; $\beta^-n=0.02$ #		
<sup>37</sup> S	-26896.42	0.20	5.05 m	0.02	7/2 <sup>-</sup>	12		$\beta^-n=100$		
<sup>37</sup> Cl	-31761.54	0.05	STABLE		3/2 <sup>+</sup>	12		IS=24.24 10		
<sup>37</sup> Cl <sup>i</sup>	-21539.7	0.3	10221.8	0.3	RQ			IT=100		
<sup>37</sup> Ar	-30947.66	0.21	35.011 d	0.019	3/2 <sup>+</sup>	12		$\epsilon=100$		
<sup>37</sup> Ar <sup>j</sup>	-25956	6	4992	6	RQ					
<sup>37</sup> K	-24800.20	0.09	1.2365 s	0.0009	3/2 <sup>+</sup>	12 14Kr04	J 1973	$\beta^+=100$	*	
<sup>37</sup> K <sup>i</sup>	-19749.9	0.8	5050.3	0.8	RQ			IT=100		
<sup>37</sup> Ca	-13136.1	0.6	181.1 ms	1.0	3/2 <sup>+</sup> #	12		$\beta^+=100$ ; $\beta^+p=82.1$ 7	*	
<sup>37</sup> Sc	3520#	300#			7/2 <sup>-</sup> #			p ?	**	
* <sup>37</sup> K	T : more precisely 14Sh25=1.23651(0.00094)									
* <sup>37</sup> Ca	TD : also 07Do17=181.7(3.6) ms; 72.2(4.3)%; also $\beta^+p=74.5(0.7)$ % from 95Tr03									
<sup>38</sup> Mg	34070#	500#	1# ms	(>260 ns)	0 <sup>+</sup>	13		$\beta^-n=100$ #; $\beta^-n=80$ #; $\beta^-2n=7$ #		
<sup>38</sup> Al	16210	370	9.0 ms	0.7	0 <sup>+</sup>	08 15St14	T 1989	$\beta^-n=100$ ; $\beta^-n=0$ #; $\beta^-2n=10$ #	*	
<sup>38</sup> Si	-4170	100	90# ms	(>1 $\mu$ s)	0 <sup>+</sup>	08		$\beta^-n=100$ #; $\beta^-n=30$ #		
<sup>38</sup> P	-14620	70	640 ms	140	(2 <sup>-</sup> )	08 15Ch56	J 1971	$\beta^-n=100$ ; $\beta^-n=12$ 5		
<sup>38</sup> S	-26861	7	170.3 m	0.7	0 <sup>+</sup>	08		$\beta^-n=100$		
<sup>38</sup> Cl	-29798.10	0.10	37.24 m	0.05	2 <sup>-</sup>	08		$\beta^-n=100$		
<sup>38</sup> Cl <sup>m</sup>	-29126.73	0.10	671.365	0.008	715 ms	3		IT=100		
<sup>38</sup> Cl <sup>i</sup>	-21590	24	8208	24	RQ			0 <sup>+</sup> T=3		
<sup>38</sup> Ar	-34714.82	0.19	STABLE		0 <sup>+</sup>	08		1934	IS=0.0629 7	
<sup>38</sup> Ar <sup>i</sup>	-24083.9	0.9	10630.9	0.9		(2 <sup>-</sup> )T=2	08			
<sup>38</sup> Ar <sup>j</sup>	-15940	30	18780	30	RQ			0 <sup>+</sup> T=3		
<sup>38</sup> K	-28800.75	0.20	7.636 m	0.018	3 <sup>+</sup> T=0	08 14Kr04	J 1937	$\beta^+=100$		
<sup>38</sup> K <sup>m</sup>	-28670.61	0.20	130.15	0.04	MD	924.46 ms	0.14	0 <sup>+</sup> T=1	08 10Ba43	T 1953
<sup>38</sup> K <sup>n</sup>	-25342.61	0.26	3458.14	0.17		21.95 $\mu$ s	0.11	(7 <sup>+</sup> )	08	1971
<sup>38</sup> Ca	-22058.50	0.19	443.70 ms	0.25	0 <sup>+</sup>	08 15Bi02	T 1966	$\beta^+=100$	*	
<sup>38</sup> Sc	-4250#	200#		<300 ns	2 <sup>-</sup> #	08 94B110	I	p ?		
<sup>38</sup> Sc <sup>m</sup>	-3580#	220#	670#	100#		5 <sup>-</sup> #	Mirror	I	IT ?; p ?	
<sup>38</sup> Ti	10870#	300#		<120 ns	0 <sup>+</sup>	08 96B121	I	2p ?		
* <sup>38</sup> Al	T : other 04Gr20=7.6(0.6) without $\gamma$ -correlation I : 89Gu03>200ns									
* <sup>38</sup> Ca	T : average 15Bi02=443.63(0.35) 11Pa38=443.77(0.36)									
<sup>39</sup> Mg	42280#	510#		<180 ns	7/2 <sup>-</sup> #	07		n ?	*	
<sup>39</sup> Al	20650#	400#	7.6 ms	1.6	5/2 <sup>+</sup> #	11		1989	$\beta^-n=100$ ; $\beta^-n=90$ #; $\beta^-2n=1$ #	
<sup>39</sup> Si	2320	140	47.5 ms	2.0	5/2 <sup>-</sup> #	15		1979	$\beta^-n=100$ ; $\beta^-n=25$ #; $\beta^-2n=2$ #	
<sup>39</sup> P	-12770	110	282 ms	24	1/2 <sup>+</sup> #	06 04Gr20	T 1977	$\beta^-n=100$ ; $\beta^-n=26$ 8	*	
<sup>39</sup> S	-23160	50	11.5 s	0.5	(7/2 <sup>-</sup> )	06		1971	$\beta^-n=100$	
<sup>39</sup> Cl	-29800.2	1.7	56.2 m	0.6	3/2 <sup>+</sup>	06		1949	$\beta^-n=100$	
<sup>39</sup> Ar	-33242	5	269 y	3	7/2 <sup>-</sup>	06		1950	$\beta^-n=100$	
<sup>39</sup> Ar <sup>i</sup>	-24161	7	9081	9	RQ			T=5/2		
<sup>39</sup> K	-33807.190	0.005	STABLE		3/2 <sup>+</sup>	06 MMC149	J 1921	IS=93.2581 44	*	
<sup>39</sup> K <sup>i</sup>	-27261.2	2.0	6546	2		7/2 <sup>-</sup> T=3/2	06	IT=100		
<sup>39</sup> Ca	-27282.7	0.6	860.3 ms	0.8	3/2 <sup>+</sup>	06 10Bi09	T 1943	$\beta^+=100$	*	
<sup>39</sup> Ca <sup>i</sup>	-20917#	9#	6366#	9#		3/2 <sup>+</sup> T=3/2	Imme	E		
<sup>39</sup> Sc	-14173	24	< 300 ns		7/2 <sup>-</sup> #	06 GAu128	D 1988	p=100	*	
<sup>39</sup> Sc <sup>i</sup>	-5050	40	9120	50	2p	(3/2 <sup>+</sup> )T=5/2	06			
<sup>39</sup> Ti	2200#	200#	28.5 ms	0.9	3/2 <sup>+</sup> #	06 07Do17	TD 1990	$\beta^+=100$ ; $\beta^+p=93.7$ 28; ...	*	
* <sup>39</sup> Mg	T : estimated half-life 1# ms for $\beta^-$ decay									
* <sup>39</sup> P	T : average 04Gr20=250(80) 98Wi.A=320(30) 95Re.A=190(50)									
* <sup>39</sup> Ar <sup>j</sup>	J : due to IAS appartenance; was (3/2,5/2) <sup>+</sup> in ENSDF2006									
* <sup>39</sup> Ca	T : average 10Bi09=860.7(1.0) 77Az01=859.4(1.6) 73Al11=860.4(3.0)									
* <sup>39</sup> Sc	D : most probably proton emitter from S <sub>p</sub> =-597(24) keV									
* <sup>39</sup> Ti	D : ...; $\beta^+2p=15$ # D : $\beta^+2p$ decay observed in 92Mo15									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>40</sup> Mg	48350#	500#	1# ms (>170 ns)	0 <sup>+</sup>	07	14Cr02 I	2007	$\beta^-$ ?; $\beta^-$ -n=100#; $\beta^-$ -2n=50#	*
<sup>40</sup> Al	27590#	400#	10# ms (>260 ns)		04		2002	$\beta^-$ ?; $\beta^-$ -n=0#; $\beta^-$ -2n=90#	
<sup>40</sup> Si	5430	350	33.0 ms 1.0	0 <sup>+</sup>	06	04Gr20 TD	1989	$\beta^-$ =100; $\beta^-$ -n=40#; $\beta^-$ -2n=60#	
<sup>40</sup> P	-8110	150	150 ms 8	(2 <sup>-</sup> , 3 <sup>-</sup> )	04		1979	$\beta^-$ =100; ; $\beta^-$ -n=15.8 21; $\beta^-$ -2n=2#	
<sup>40</sup> S	-22838	4	8.8 s 2.2	0 <sup>+</sup>	04		1971	$\beta^-$ =100	
<sup>40</sup> Cl	-27560	30	1.35 m 0.02	2 <sup>-</sup>	04		1956	$\beta^-$ =100	
<sup>40</sup> Ar	-35039.8946	0.0022	STABLE	0 <sup>+</sup>	04		1920	IS=99.6035 25	
<sup>40</sup> K	-33535.49	0.06	1.248 Gy 0.003	4 <sup>-</sup>	04	14Kr04 J	1935	IS=0.0117 1; $\beta^-$ =89.28 13; $\beta^+$ =10.72 13	
<sup>40</sup> K <sup>m</sup>	-31891.85	0.06	1643.639 0.011	336 ns 12	0 <sup>+</sup>	04	1968	IT=100	
<sup>40</sup> K <sup>i</sup>	-29151.5	0.3	4384.0 0.3		0 <sup>+</sup> T=2	04		IT=100	
<sup>40</sup> Ca	-34846.384	0.021	STABLE (>5.9 Zy)	0 <sup>+</sup>	04	99Be64 T	1922	IS=96.94 16; 2 $\beta^+$ ?	
<sup>40</sup> Ca <sup>i</sup>	-27188.20	0.05	7658.18 0.05		4 <sup>-</sup> T=1	04 AHW E		IT=100	*
<sup>40</sup> Ca <sup>j</sup>	-22858.4	1.0	11988 1		0 <sup>+</sup> T=2	04		IT=100	
<sup>40</sup> Sc	-20523.3	2.8		182.3 ms 0.7	4 <sup>-</sup>	04	1955	$\beta^+$ =100; $\beta^+$ p=0.44 7; $\beta^+$ $\alpha$ =0.017 5	
<sup>40</sup> Sc <sup>i</sup>	-16164	6	4359 6 RQ		0 <sup>+</sup> T=2	04		IT=100	
<sup>40</sup> Ti	-8850	160		52.4 ms 0.3	0 <sup>+</sup>	04 07Do17 TD	1982	$\beta^+$ =100; $\beta^+$ p=95.8 13	
<sup>40</sup> V	12170#	300#			2 <sup>-</sup> #			p ?	
* <sup>40</sup> Mg	I : 14Cr02 5 events observed in direct two-proton removal from 42Si								**
* <sup>40</sup> Ca <sup>i</sup>	E : Original 7658.23(0.05) recalibrated -0.05 keV for <sup>27</sup> Al+p resonances								**
<sup>41</sup> Al	33420#	500#	2# ms (>260 ns)	5/2 <sup>+</sup> #	16		2002	$\beta^-$ ?; $\beta^-$ -n=50#; $\beta^-$ -2n=10#	
<sup>41</sup> Si	12120	550	20.0 ms 2.5	7/2 <sup>-</sup> #	16		1989	$\beta^-$ =100; $\beta^-$ -n=45#; $\beta^-$ -2n=10#	
<sup>41</sup> P	-4980	120	101 ms 5	1/2 <sup>+</sup> #	16		1979	$\beta^-$ =100; $\beta^-$ -n=30 10; $\beta^-$ -2n=0.2#	
<sup>41</sup> S	-19009	4	1.99 s 0.05	7/2 <sup>-</sup> #	16		1979	$\beta^-$ =100; $\beta^-$ -n=0.1#	
<sup>41</sup> Cl	-27310	70	38.4 s 0.8	(1/2 <sup>+</sup> )	16		1971	$\beta^-$ =100	
<sup>41</sup> Ar	-33067.5	0.3		109.61 m 0.04	7/2 <sup>-</sup>	16	1936	$\beta^-$ =100	
<sup>41</sup> K	-35559.543	0.004	STABLE	3/2 <sup>+</sup>	16	14Kr04 J	1921	IS=6.7302 44	
<sup>41</sup> K <sup>i</sup>	-27210	15	8349 15 RQ		7/2 <sup>-</sup> T=5/2	16 75Me10 J	1975		*
<sup>41</sup> Ca	-35137.89	0.14		99.4 ky 1.5	7/2 <sup>-</sup>	16	1939	$\epsilon$ =100	
<sup>41</sup> Ca <sup>i</sup>	-29320.8	0.5	5817.1 0.5	< 28 fs	3/2 <sup>+</sup> T=3/2	16		IT=100	
<sup>41</sup> Sc	-28642.41	0.08		596.3 ms 1.7	7/2 <sup>-</sup>	16	1941	$\beta^+$ =100	
<sup>41</sup> Sc <sup>r</sup>	-25760.09	0.09	2882.32 0.05 RQ		7/2 <sup>+</sup>	16		P=59 2; IT=41 2	
<sup>41</sup> Sc <sup>i</sup>	-22704	3	5939 3 RQ		3/2 <sup>+</sup> T=3/2	16		p=100	
<sup>41</sup> Ti	-15698	28		81.9 ms 0.5	3/2 <sup>+</sup>	16 07Do17 D	1964	$\beta^+$ =100; $\beta^+$ p=91.1 6	
<sup>41</sup> V	320#	200#			7/2 <sup>-</sup> #			p ?	
* <sup>41</sup> K <sup>i</sup>	I : ENSDF=5/2 <sup>-</sup> , 7/2 <sup>-</sup> and T=3/2; NUBASE adopts this level as IAS of <sup>41</sup> Ar ground-state								**
<sup>42</sup> Al	40100#	600#	1# ms (>170 ns)		16		2007	$\beta^-$ ?; $\beta^-$ -n=30#; $\beta^-$ -2n=40#	
<sup>42</sup> Si	16470#	500#	12.5 ms 3.5	0 <sup>+</sup>	16		1990	$\beta^-$ =100; $\beta^-$ -n=40#; $\beta^-$ -2n=60#	
<sup>42</sup> P	1010	310	48.5 ms 1.5		16		1979	$\beta^-$ =100; $\beta^-$ -n=50 20; $\beta^-$ -2n=20#	
<sup>42</sup> S	-17637.7	2.8	1.016 s 0.015	0 <sup>+</sup>	16		1979	$\beta^-$ =100; $\beta^-$ -n<4	
<sup>42</sup> Cl	-24830	60	6.8 s 0.3	(2 <sup>-</sup> )	16		1971	$\beta^-$ =100; $\beta^-$ -n=0#	
<sup>42</sup> Ar	-34423	6	32.9 y 1.1	0 <sup>+</sup>	16		1952	$\beta^-$ =100	
<sup>42</sup> K	-35022.03	0.11	12.355 h 0.007	2 <sup>-</sup>	16	14Kr04 J	1935	$\beta^-$ =100	
<sup>42</sup> K <sup>i</sup>	-28570	100	6450 100		(0 <sup>+</sup> )T=3	16			
<sup>42</sup> Ca	-38547.24	0.15	STABLE	0 <sup>+</sup>	16		1934	IS=0.647 23	
<sup>42</sup> Ca <sup>i</sup>	-28797	10	9750 10		(2 <sup>-</sup> )T=2	16			
<sup>42</sup> Sc	-32121.15	0.17		680.79 ms 0.28	0 <sup>+</sup> T=1	16	1955	$\beta^+$ =100	
<sup>42</sup> Sc <sup>m</sup>	-31504.83	0.18	616.32 0.06 MD		7 <sup>+</sup>	16	1963	$\beta^+$ =100	*
<sup>42</sup> Sc <sup>r</sup>	-26044.89	0.17	6076.26 0.07 RQ		(2 <sup>+</sup> , 3 <sup>+</sup> , 4 <sup>+</sup> )	16		IT=100	
<sup>42</sup> Ti	-25104.67	0.28		208.65 ms 0.80	0 <sup>+</sup>	16	1964	$\beta^+$ =100	
<sup>42</sup> V	-7620#	200#		<55 ns	2 <sup>-</sup> #	16 92Bo37 I		p ?	
<sup>42</sup> Cr	6730#	400#		13.3 ms 1.0	0 <sup>+</sup>	16	1996	$\beta^+$ $\approx$ 100; $\beta^+$ p=94.4 50; 2p ?	
* <sup>42</sup> Sc <sup>m</sup>	J : 5 <sup>+</sup> , 6 <sup>+</sup> , 7 <sup>+</sup> from $\beta^+$ decay to 6 <sup>+</sup> level; 7 <sup>+</sup> is most likely from shell model								**
<sup>43</sup> Al	47020#	800#	1# ms (>170 ns)	5/2 <sup>+</sup> #	15		2007	$\beta^-$ ?; $\beta^-$ -n=100#; $\beta^-$ -2n=50#	
<sup>43</sup> Si	23100#	600#	15# ms (>260 ns)	3/2 <sup>-</sup> #	15	02No11 I	2002	$\beta^-$ ?; $\beta^-$ -n=40#; $\beta^-$ -2n=30#	
<sup>43</sup> P	4680	550	35.8 ms 1.3	(1/2 <sup>+</sup> )	15	04Gr20 T	1989	$\beta^-$ =100; $\beta^-$ -n=100; $\beta^-$ -2n=10#	*
<sup>43</sup> S	-12195	5	265 ms 13	3/2 <sup>-</sup> #	15		1979	$\beta^-$ =100; $\beta^-$ -n=40 10	
<sup>43</sup> S <sup>m</sup>	-11874	5	320.7 0.5	415.0 ns 2.6	(7/2 <sup>-</sup> )	15 09Ga05 J	2000	IT=100	*
<sup>43</sup> Cl	-24160	60	3.13 s 0.09	(3/2 <sup>+</sup> )	15	06Wi10 J	1976	$\beta^-$ =100; $\beta^-$ -n=2#	
<sup>43</sup> Ar	-32010	5	5.37 m 0.06	5/2 <sup>(-)</sup>	15		1969	$\beta^-$ =100	
<sup>43</sup> K	-36575.4	0.4	22.3 h 0.1	3/2 <sup>+</sup>	01	14Kr04 J	1949	$\beta^-$ =100	
<sup>43</sup> K <sup>m</sup>	-35837.1	0.4	738.30 0.06	200 ns 5	7/2 <sup>-</sup>	15	1978	IT=100	
<sup>43</sup> Ca	-38408.82	0.23	STABLE	7/2 <sup>-</sup>	15		1934	IS=0.135 10	
<sup>43</sup> Ca <sup>i</sup>	-30414	14	7995 14 RQ		(3/2 <sup>+</sup> )T=5/2	15			
<sup>43</sup> Ca <sup>j</sup>	-36188.1	1.9		3.891 h 0.012	7/2 <sup>-</sup>	15	1935	$\beta^+$ =100	
<sup>43</sup> Sc <sup>m</sup>	-36036.3	1.9	151.79 0.08	438 $\mu$ s 5	3/2 <sup>+</sup>	15 77Mi10 T	1964	IT=100	*
<sup>43</sup> Sc <sup>n</sup>	-33064.4	1.9	3123.73 0.15	472 ns 3	19/2 <sup>-</sup>	15 08Fe02 T	1978	IT=100	*
<sup>43</sup> Sc <sup>i</sup>	-31956	3	4232 4 RQ		7/2 <sup>-</sup> T=3/2	15			

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...											
<sup>43</sup> Ti	-29321	7		509 ms	5		15	1948	$\beta^+=100; \beta^+p?$		
<sup>43</sup> Ti <sup>m</sup>	-29008	7	313.0	11.9 $\mu$ s	0.3		15	1978	IT=100		
<sup>43</sup> Ti <sup>n</sup>	-26255	7	3066.4	556 ns	6		15	1978	IT=100		
<sup>43</sup> Ti <sup>i</sup>	-24610#	50#	4710#	50#							
<sup>43</sup> V	-17920	40		79.3 ms	2.4		15 07Do17 D	1987	$\beta^+=100; \beta^+p<2.5$		
<sup>43</sup> Vi	-9705	15	8210			RQ					
<sup>43</sup> Cr	-1970#	400#		21.1 ms	0.3			1992	$\beta^+=100; \beta^+p=79.3\ 30; \beta^+2p=11.6\ 10; \dots$ *		
<sup>43</sup> P	T : average 04Gr20=36.5(1.5) 95So03=33(3) **										
<sup>43</sup> Sc <sup>m</sup>	T : average 12Ch16=415(3) 09Ga05=415(5) **										
<sup>43</sup> Sc <sup>m</sup>	T : average 77Mi10=438(7) 65De15=470(20) 64Ho14=435(7) **										
<sup>43</sup> Sc <sup>n</sup>	T : average 08Fe02=481(9) 81Da06=469(4) 78Ha07=473(5) **										
<sup>43</sup> Cr	D : ... ; $\beta^+3p=0.13+18-8; \beta^+\alpha?$ **										
<sup>43</sup> Cr	T : average 11Po01=20.6(0.9) 07Do17=21.1(0.4) 01Gi01=21.6(0.7) **										
<sup>44</sup> Si	28510#	600#		10# ms	(>360 ns)		0 <sup>+</sup>	11	2007	$\beta^-?$ ; $\beta^-n=100\#; \beta^-2n=50\#$	
<sup>44</sup> P	10450#	500#		18.5 ms	2.5			11	1989	$\beta^-n=100; \beta^-n=20\#; \beta^-2n=70\#$	
<sup>44</sup> S	-9204	5		100 ms	1		0 <sup>+</sup>	11	1979	$\beta^-n=100; \beta^-n=18\ 3$	
<sup>44</sup> S <sup>m</sup>	-7839	5	1365.0	0.8			0 <sup>+</sup>	11	2005	IT=100	
<sup>44</sup> Cl	-20380	140		2.619 $\mu$ s	0.026		0 <sup>+</sup>	11	1979	$\beta^-n=100; \beta^-n<8$	
<sup>44</sup> Ar	-32673.3	1.6		11.87 m	0.05		0 <sup>+</sup>	11	1969	$\beta^-n=100$	
<sup>44</sup> K	-35781.5	0.4		22.13 m	0.19		2 <sup>-</sup>	11 14Kr04 J	1954	$\beta^-n=100$	
<sup>44</sup> Ca	-41468.7	0.3				STABLE	0 <sup>+</sup>	11	1922	IS=2.09 11	
<sup>44</sup> Ca <sup>i</sup>	-29619	10	11850	10			2 <sup>-</sup> T=3	11		*	
<sup>44</sup> Sc	-37816.0	1.8		4.0420 h	0.0025		2 <sup>+</sup>	11 16Ga24 T	1937	$\beta^+=100$	
<sup>44</sup> Sc <sup>m</sup>	-37748.1	1.8	67.8679	0.0014			1 <sup>-</sup>	11	1967	IT=100	
<sup>44</sup> Sc <sup>n</sup>	-37669.8	1.8	146.1914	0.0020			0 <sup>-</sup>	11	1963	IT=100	
<sup>44</sup> Sc <sup>p</sup>	-37544.8	1.8	271.240	0.010			6 <sup>+</sup>	11	1940	IT=98.80 7; $\beta^+=1.20\ 7$	
<sup>44</sup> Sc <sup>i</sup>	-35038.2	2.5	2778	3		RQ	0 <sup>+</sup> T=2	11			
<sup>44</sup> Ti	-37548.6	0.7		59.1 y	0.3		0 <sup>+</sup>	11	1954	$\epsilon=100$	
<sup>44</sup> Ti <sup>i</sup>	-30942.2	0.9	6606.4	0.5			2 <sup>+</sup> T=1	11		IT=100	
<sup>44</sup> Ti <sup>j</sup>	-28210.6	2.1	9338	2			0 <sup>+</sup> frg.T=2	11		IT=100	
<sup>44</sup> V	-24120	180		111 ms	7	*	(2) <sup>+</sup>	11	1971	$\beta^+=100; \beta^+\alpha=?; \beta^+p?$	
<sup>44</sup> V <sup>m</sup>	-23850#	210#	270#	100#	*		(6) <sup>+</sup>	11	1997	$\beta^+=100$	
<sup>44</sup> V <sup>n</sup>	-23970#	210#	150#	100#			0 <sup>-</sup> #			Mirror I	
<sup>44</sup> Vi	-21124	13	2990	180		p	0 <sup>+</sup> #T=2		92Bo37 D	1992	p=100
<sup>44</sup> Cr	-13360#	300#		42.8 ms	0.6		0 <sup>+</sup>	11 07Do17 D	1987	$\beta^+=100; \beta^+p=14.0\ 9$	
<sup>44</sup> Mn	7030#	500#		<105 ns			2 <sup>-</sup> #	11		p?	
<sup>44</sup> Ca <sup>i</sup>	J : ENSDF no $J^\pi$ ; data from (e,e <sup>-</sup> ) scattering 84Ra04; IAS candidate **										
<sup>44</sup> Sc	T : 16Ga24=242.52(0.15) min, but the authors quote in error 4.042(0.025) h **										
<sup>44</sup> Ti <sup>j</sup>	E : strongest fragment 9338(2); other 40(2) lower **										
<sup>45</sup> Si	37490#	700#		1# ms			3/2 <sup>-</sup> #			$\beta^-?$ ; $\beta^-n=100\#; \beta^-2n=50\#$	
<sup>45</sup> P	15600#	500#		8# ms	(>200 ns)		1/2 <sup>+</sup> #	08	1990	$\beta^-?$ ; $\beta^-n=30\#; \beta^-2n=30\#$	
<sup>45</sup> S	-3990	1040		68 ms	2		3/2 <sup>-</sup> #	08	1989	$\beta^-n=100; \beta^-n=54; \beta^-2n=4\#$	
<sup>45</sup> Cl	-18260	140		413 ms	25		(3/2 <sup>+</sup> )	08 12Ri08 J	1979	$\beta^-n=100; \beta^-n=24\ 4$	
<sup>45</sup> Ar	-29770.8	0.5		21.48 s	0.15		(5/2 <sup>-</sup> , 7/2 <sup>-</sup> )	08	1974	$\beta^-n=100$	
<sup>45</sup> K	-36615.6	0.5		17.8 m	0.6		3/2 <sup>+</sup>	08 14Kr04 J	1964	$\beta^-n=100$	
<sup>45</sup> Ca	-40812.2	0.4		162.61 d	0.09		7/2 <sup>-</sup>	08	1940	$\beta^-n=100$	
<sup>45</sup> Sc	-41071.9	0.7				STABLE	7/2 <sup>-</sup>	08	1923	IS=100.	
<sup>45</sup> Sc <sup>m</sup>	-41059.5	0.7	12.40	0.05			3/2 <sup>+</sup>	08	1964	IT=100	
<sup>45</sup> Sc <sup>i</sup>	-34373	15	6699	15			7/2 <sup>-</sup> T=5/2	08			
<sup>45</sup> Ti	-39009.8	0.8		184.8 m	0.5		7/2 <sup>-</sup>	08	1941	$\beta^+=100$	
<sup>45</sup> Ti <sup>m</sup>	-38973.3	0.8	36.53	0.15			3/2 <sup>-</sup>	08	2006	IT=100	
<sup>45</sup> Ti <sup>i</sup>	-34291	3	4719	3		RQ	7/2 <sup>-</sup> T=3/2	08			
<sup>45</sup> V	-31886.0	0.9		547 ms	6		7/2 <sup>-</sup>	08	1975	$\beta^+=100$	
<sup>45</sup> V <sup>m</sup>	-31829.2	1.1	56.8	0.6			(3/2 <sup>-</sup> )	08 11Ho02 T	1980	IT=100	
<sup>45</sup> Vi	-27090	9	4796	9		RQ	7/2 <sup>-</sup> T=3/2	08		p=100	
<sup>45</sup> Cr	-19510	40		60.9 ms	0.4	*	7/2 <sup>-</sup> #	08	1974	$\beta^+=100; \beta^+p=34.4\ 8$	
<sup>45</sup> Cr <sup>m</sup>	-19400	40	107	1	*		(3/2 <sup>-</sup> )	11 11Ho02 ETJ	2011	IT=100	
<sup>45</sup> Mn	-5250#	400#		<70 ns			7/2 <sup>-</sup> #	08 92Bo37 I		p?	
<sup>45</sup> Fe	13760#	400#		2.2 ms	0.3		3/2 <sup>+</sup> #	08 05Do20 T	1996	2p=57 10; $\beta^+<43; \beta^+p<43; \beta^+p=25\ 5$ *	
<sup>45</sup> V <sup>m</sup>	T : average 11Ho02=468(23) 87Ha.B=430(80) 82Ho11=539(18) 82Al.C=610(80) and **										
<sup>45</sup> V <sup>m</sup>	T : 80Gr.A=510(50) **										
<sup>45</sup> Fe	T : average 05Do20=1.6(+0.5-0.3) 02Gi09=4.7(+3.4-1.4) 02Pf02=3.2(+2.6-1.0) **										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>46</sup> P	22970# 700#		4# ms (>200 ns)			00 90Le03 I	1990	$\beta^- ?$ ; $\beta^- n=0\#$ ; $\beta^- 2n=90\#$
<sup>46</sup> S	340# 500#		50 ms 8	0 <sup>+</sup>	10		1989	$\beta^- =100$ ; $\beta^- n=70\#$ ; $\beta^- 2n=3\#$
<sup>46</sup> Cl	-13860 210		232 ms 2	2 <sup>-</sup> #	12		1989	$\beta^- =100$ ; $\beta^- n=60 9$ ; $\beta^- 2n=0.3\#$
<sup>46</sup> Ar	-29772.9 1.1		8.4 s 0.6	0 <sup>+</sup>	00		1974	$\beta^- =100$
<sup>46</sup> K	-35413.9 0.7		105 s 10	2 <sup>-</sup>	00	14Pa45 J	1965	$\beta^- =100$
<sup>46</sup> Ca	-43139.4 2.2		STABLE	0 <sup>+</sup>	00		1938	IS=0.004 3; $2\beta^- ?$
<sup>46</sup> Sc	-41761.2 0.7		83.80 d 0.03	4 <sup>+</sup>	00	14Un01 T	1936	$\beta^- =100$
<sup>46</sup> Sc <sup>m</sup>	-41709.2 0.7	52.011 0.001	9.4 $\mu$ s 0.8	6 <sup>+</sup>	00		1966	IT=100
<sup>46</sup> Sc <sup>n</sup>	-41618.7 0.7	142.528 0.007	18.75 s 0.04	1 <sup>-</sup>	00		1948	IT=100
<sup>46</sup> Sc <sup>i</sup>	-36748 4	5013 4 RQ		0 <sup>+</sup> T=3	00			
<sup>46</sup> Ti	-44127.80 0.16		STABLE	0 <sup>+</sup>	00		1934	IS=8.25 3
<sup>46</sup> Ti <sup>i</sup>	-34962 7	9166 7 RQ		4 <sup>+</sup> T=2	00			
<sup>46</sup> Ti <sup>j</sup>	-29977 6	14151 6 RQ		0 <sup>+</sup> T=3	00			
<sup>46</sup> V	-37075.35 0.20		422.64 ms 0.05	0 <sup>+</sup> T=1	00	12Pa07 T	1952	$\beta^+ =100$
<sup>46</sup> V <sup>m</sup>	-36273.89 0.22	801.46 0.10	1.02 ms 0.07	3 <sup>+</sup> T=0	00		1962	IT=100
<sup>46</sup> Cr	-29472 11		224.3 ms 1.3	0 <sup>+</sup>	10	15Mo01 T	1972	$\beta^+ =100$
<sup>46</sup> Cr <sup>i</sup>	-20328 13	9144 17 RQ		(4 <sup>+</sup> )T=2	10			p=?
<sup>46</sup> Mn	-12570# 400#		36.2 ms 0.4	(4 <sup>+</sup> )	10		1987	$\beta^+ =100$ ; $\beta^+ p=57.0 8$ ; $\beta^+ 2p\approx 18$ ; $\beta^+ \alpha ?$
<sup>46</sup> Mn <sup>m</sup>	-12420# 410#	150# 100#	1# ms	1 <sup>-</sup> #				$\beta^+ ?$
<sup>46</sup> Mn <sup>i</sup>	-7390 50	5180# 400# p		T=3				
<sup>46</sup> Fe	910# 500#		13.0 ms 2.0	0 <sup>+</sup>	10	07Do17 TD	1992	$\beta^+ =100$ ; $\beta^+ p=78.7 38$ ; $\beta^+ 2p=?$
<sup>46</sup> Ca	T: 99Be64: 0v- $\beta\beta > 100$ Ey							
<sup>46</sup> Sc	T: average 14Un01=83.84(0.08) 83Wa26=83.73(0.11) 80Ho17=83.819(0.080);							
<sup>46</sup> Sc	T: all values are from standard labs							
<sup>46</sup> Sc	T: original unc of 80Ho17=0.006 increased to 0.1% by evaluator							
<sup>46</sup> V	T: average 12Pa07=422.66(0.06) 97Ko65=422.57(0.13)							
<sup>46</sup> Mn	T: others 92Bo37=41(+7-6) 01Gi01=34.0(+4.5-3.5)							
<sup>46</sup> Mn	D: $\beta^+ 2p\approx 18\%$ estimated from $P_p = \beta^+ p + 2 \times \beta^+ 2p=57(1)\%$							
<sup>46</sup> Fe	T: average 14Po05=16.4(+4.2-2.8) 07Do17=13.0(2.0) 01Gi01=9.7(+3.5-4.3)							
<sup>46</sup> Fe	D: other $\beta^+ p$ 14Po05=66(4)% 01Gi01=36(20)%; $\beta^+ 2p$ , 1 event in 14Po05							
<sup>47</sup> P	29710# 800#		2# ms	1/2 <sup>+</sup> #				$\beta^- ?$ ; $\beta^- n=0.4\#$ ; $\beta^- 2n=0.03\#$
<sup>47</sup> S	7370# 500#		20# ms (>200 ns)	3/2 <sup>-</sup> #	07	89Gu03 I	1989	$\beta^- ?$ ; $\beta^- n=10\#$ ; $\beta^- 2n=10\#$
<sup>47</sup> Cl	-9780# 400#		101 ms 5	3/2 <sup>+</sup> #	07		1989	$\beta^- =100$ ; $\beta^- n<3$ ; $\beta^- 2n=0.3\#$
<sup>47</sup> Ar	-25366.3 1.1		1.23 s 0.03	(3/2) <sup>-</sup>	07		1985	$\beta^- =100$ ; $\beta^- n<0.2$
<sup>47</sup> K	-35712.0 1.4		17.50 s 0.24	1/2 <sup>+</sup>	07	14Kr04 J	1964	$\beta^- =100$
<sup>47</sup> Ca	-42344.4 2.2		4.536 d 0.003	7/2 <sup>-</sup>	07		1951	$\beta^- =100$
<sup>47</sup> Sc	-44336.6 1.9		3.3492 d 0.0006	7/2 <sup>-</sup>	07		1945	$\beta^- =100$
<sup>47</sup> Sc <sup>m</sup>	-43569.8 1.9	766.83 0.09	272 ns 8	(3/2) <sup>+</sup>	07		1968	IT=100
<sup>47</sup> Ti	-44937.36 0.12		STABLE	5/2 <sup>-</sup>	07		1934	IS=7.44 2
<sup>47</sup> Ti <sup>i</sup>	-37588.4 0.7	7349.0 0.7		7/2 <sup>-</sup> T=5/2	07			
<sup>47</sup> V	-42006.62 0.17		32.6 m 0.3	3/2 <sup>-</sup>	07		1942	$\beta^+ =100$
<sup>47</sup> V <sup>i</sup>	-37856.27 0.20	4150.35 0.11		5/2(-)T=3/2	07			IT=100
<sup>47</sup> Cr	-34563 6		500 ms 15	3/2 <sup>-</sup>	07		1972	$\beta^+ =100$
<sup>47</sup> Cr <sup>j</sup>	-29803# 21#	4760# 20#		5/2 <sup>-</sup> #T=5/2				
<sup>47</sup> Mn	-22570 30		88.0 ms 1.3	5/2 <sup>-</sup> #	07	07Do17 TD	1987	$\beta^+ =100$ ; $\beta^+ p<1.7$
<sup>47</sup> Mn <sup>i</sup>	-15191 17	7380 40 RQ		7/2 <sup>-</sup> #T=5/2	07		2001	p=100
<sup>47</sup> Fe	-6870# 500#		21.9 ms 0.2	7/2 <sup>-</sup> #	07	07Do17 TD	1992	$\beta^+ =100$ ; $\beta^+ p=88.4 9$
<sup>47</sup> Fe <sup>m</sup>	-6100# 510#	770# 100#		3/2 <sup>+</sup> #		Mirror I		IT?
<sup>47</sup> Co	10370# 600#			7/2 <sup>-</sup> #	07	Mirror I		p?
<sup>48</sup> S	12760# 600#		10# ms (>200 ns)	0 <sup>+</sup>	06		1990	$\beta^- ?$ ; $\beta^- n=80\#$ ; $\beta^- 2n=10\#$
<sup>48</sup> Cl	-4280# 500#		100# ms (>200 ns)		06	89Gu03 I	1989	$\beta^- ?$ ; $\beta^- n=60\#$ ; $\beta^- 2n=40\#$
<sup>48</sup> Ar	-22280 310		415 ms 15	0 <sup>+</sup>	10	12We08 TD	2004	$\beta^- =100$ ; $\beta^- n=38 6$
<sup>48</sup> K	-32284.5 0.8		6.8 s 0.2	1 <sup>-</sup>	06	14Pa45 J	1972	$\beta^- =100$ ; $\beta^- n=1.14 15$
<sup>48</sup> Ca	-44224.63 0.10		45 Ey 6	0 <sup>+</sup>	06	15Ba11 T	1938	IS=0.187 21; $2\beta^- =75 +25-38$ ; $\beta^- ?$
<sup>48</sup> Sc	-44504 5		43.67 h 0.09	6 <sup>+</sup>	06		1937	$\beta^- =100$
<sup>48</sup> Ti	-48492.71 0.11		STABLE	0 <sup>+</sup>	06		1923	IS=73.72 3
<sup>48</sup> Ti <sup>i</sup>	-37767 6	10726 6		(6 <sup>+</sup> )T=3	06			
<sup>48</sup> V	-44477.7 1.0		15.9735 d 0.0025	4 <sup>+</sup>	06		1937	$\beta^+ =100$
<sup>48</sup> V <sup>i</sup>	-41458.84 0.24	3018.9 0.9 RQ		(0 <sup>+</sup> )T=2	06			IT=100
<sup>48</sup> Cr	-42822 7		21.56 h 0.03	0 <sup>+</sup>	06		1952	$\beta^+ =100$
<sup>48</sup> Cr <sup>i</sup>	-37029 7	5792.77 0.24		4 <sup>+</sup> T=1	06		1987	IT=100
<sup>48</sup> Cr <sup>j</sup>	-34062 15	8760 17 RQ		0 <sup>+</sup> frg.T=2	06			
<sup>48</sup> Mn	-29296 7		158.1 ms 2.2	4 <sup>+</sup>	06		1987	$\beta^+ =100$ ; $\beta^+ p=0.28 4$ ; $\beta^+ \alpha=6e-4$
<sup>48</sup> Mn <sup>i</sup>	-26260 7	3036.7 0.9		0 <sup>+</sup> T=2	06	MMC12 J		p=100
<sup>48</sup> Fe	-18000# 400#		45.3 ms 0.6	0 <sup>+</sup>	06	07Do17 TD	1987	$\beta^+ =100$ ; $\beta^+ p=15.3 5$

... A-group is continued on next page ...



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>48</sup> Co	1500#	500#						p ?	
<sup>48</sup> Ni	16790#	500#	2.8 ms	0.8	0 <sup>+</sup>	06 11Po09	TD 2000	2p=70 20; β <sup>+</sup> =30 20; β <sup>+</sup> p ?	*
* <sup>48</sup> Ar	T : average 12We08=381(35) 412(19) 04Gr20=475(40)								
* <sup>48</sup> Ca	T : symmetrized from 15Ba11=44(+6-5)								
* <sup>48</sup> Cr <sup>i</sup>	E : strongest frg; other: 10(15)keV lower								
* <sup>48</sup> Fe	D : average 07Do17=15.9(6)% 16Or03=14.4(7)%; other 96Fa09>3.6(1.1)%								
* <sup>48</sup> Fe	T : other 16Or03=51(3) 96Fa09=44(7)								
* <sup>48</sup> Ni	T : average 05Do20=2.1(+2.1-0.7) 14Po05=11Po09=2.1(+1.4-0.4)								
<sup>49</sup> S	21090#	670#		<200 ns	3/2 <sup>-</sup> #	08 90Le03	I	n ?; β <sup>-</sup> ?; β <sup>-</sup> n=4#; β <sup>-</sup> 2n=30#	*
<sup>49</sup> Cl	940#	600#		50# ms (>200 ns)	3/2 <sup>+</sup> #	08 89Gu03	I	1989	β <sup>-</sup> ?; β <sup>-</sup> n=70#; β <sup>-</sup> 2n=20#
<sup>49</sup> Ar	-17190#	400#		236 ms	8	08 12We08	TD	1989	β <sup>-</sup> =100; β <sup>-</sup> n=29 6; β <sup>-</sup> 2n=0.3#
<sup>49</sup> K	-29611.5	0.8		1.26 s	0.05	11 14Pa45	J	1972	β <sup>-</sup> =100; β <sup>-</sup> n=86 9
<sup>49</sup> Ca	-41299.77	0.20		8.718 m	0.006			1950	β <sup>-</sup> =100
<sup>49</sup> Sc	-46561.3	2.7		57.18 m	0.13			1940	β <sup>-</sup> =100
<sup>49</sup> Ti	-48563.79	0.11		STABLE				1934	IS=5.41 2
<sup>49</sup> V	-47961.9	0.8		330 d	15			1940	ε=100
<sup>49</sup> V <sup>i</sup>	-41530	4	6432	4	RQ				
<sup>49</sup> Cr	-45333.1	2.2		42.3 m	0.1			1942	β <sup>+</sup> =100
<sup>49</sup> Cr <sup>i</sup>	-40569	5	4764	5				1969	IT=100
<sup>49</sup> Mn	-37620.6	2.3		382 ms	7			1970	β <sup>+</sup> =100
<sup>49</sup> Mn <sup>i</sup>	-32804	18	4817	18	p				p=100
<sup>49</sup> Fe	-24751	24		64.7 ms	0.3			1970	β <sup>+</sup> =100; β <sup>+</sup> p=56.7 4
<sup>49</sup> Co	-9880#	500#			<35 ns				p ?
<sup>49</sup> Ni	8200#	600#		7.5 ms	1.0			1996	β <sup>+</sup> =100; β <sup>+</sup> p=83 13
* <sup>49</sup> S	I : statistics precludes any conclusion, say authors								
* <sup>49</sup> Cr <sup>i</sup>	E : strongest component surrounded by several weak l=3 lines								
* <sup>49</sup> Cr <sup>i</sup>	E : 85Fu03 cannot confirm IAS identity and frgs								
<sup>50</sup> Cl	7740#	600#		20# ms (>620 ns)		10 09Ta24	I	2009	β <sup>-</sup> ?; β <sup>-</sup> n=70#; β <sup>-</sup> 2n=30#
<sup>50</sup> Ar	-13330#	500#		106 ms	6			1989	β <sup>-</sup> =100; β <sup>-</sup> n=37 7; β <sup>-</sup> 2n=0.1#
<sup>50</sup> K	-25728	8		472 ms	4	10 14Pa45	J	1972	β <sup>-</sup> =100; β <sup>-</sup> n=29 3; β <sup>-</sup> 2n=10#
<sup>50</sup> K <sup>m</sup>	-25557	8	171.4	0.4		10 FGK127	J	1999	IT=100
<sup>50</sup> Ca	-39589.2	1.6		13.9 s	0.6			1964	β <sup>-</sup> =100
<sup>50</sup> Sc	-44547	15		102.5 s	0.5			1959	β <sup>+</sup> =100
<sup>50</sup> Sc <sup>m</sup>	-44290	15	256.895	0.010				1963	IT>97.5; β <sup>-</sup> <2.5
<sup>50</sup> Ti	-51431.66	0.12		STABLE				1934	IS=5.18 2
<sup>50</sup> V	-49224.0	0.4		150 Py	40			1949	IS=0.250 4; β <sup>+</sup> =83 11; β <sup>-</sup> =17 11
<sup>50</sup> V <sup>i</sup>	-44410.42	0.29	4813.6	0.5	RQ				
<sup>50</sup> Cr	-50262.1	0.4		STABLE	(>1.3 Ey)			1930	IS=4.345 13; 2β <sup>+</sup> ?
<sup>50</sup> Cr <sup>i</sup>	-41836	7	8426	7	RQ				
<sup>50</sup> Cr <sup>j</sup>	-37039	6	13223	6	RQ				
<sup>50</sup> Mn	-42627.6	0.4		283.19 ms	0.10			1952	β <sup>+</sup> =100
<sup>50</sup> Mn <sup>m</sup>	-42402.3	0.4	225.31	0.07	MD			1962	β <sup>+</sup> =100
<sup>50</sup> Fe	-34476	8		152.1 ms	0.6			1977	β <sup>+</sup> =100; β <sup>+</sup> p≈0
<sup>50</sup> Fe <sup>i</sup>	-26000	10	8477	13	RQ				
<sup>50</sup> Co	-17630#	400#		38.8 ms	0.2			1987	β <sup>+</sup> =100; β <sup>+</sup> p=70.5 7; β <sup>+</sup> 2p ?
<sup>50</sup> Co <sup>i</sup>	-12746	15	4880#	400#	2p				p=100
<sup>50</sup> Ni	-4120#	500#		18.5 ms	1.2			1994	β <sup>+</sup> =100; β <sup>+</sup> p=86.7 39; β <sup>+</sup> 2p ?
* <sup>50</sup> K <sup>m</sup>	E : also 12Ka36=172.4(0.5) J : E2 to ground-state								
* <sup>50</sup> K <sup>m</sup>	T : others recent 12Ka36=138(+50-41) 09Cr03<500 ns; discovered in 99Le68								
* <sup>50</sup> V	T : symmetrized from 140(+40-30)								
* <sup>50</sup> Cr	T : 03Bi05>1.3Ey 85No03>0.18Ey								
* <sup>50</sup> Mn	T : also 13Su07=288(7)								
* <sup>50</sup> Ni	T : other 03Ma34=12(+3-2) D : other 03Ma34=70(20)%								
<sup>51</sup> Cl	14290#	700#		2# ms (>200 ns)	3/2 <sup>+</sup> #	06		1990	β <sup>-</sup> ?; β <sup>-</sup> n=40#; β <sup>-</sup> 2n=20#
<sup>51</sup> Ar	-6690#	600#		60# ms (>200 ns)	3/2 <sup>-</sup> #	06 89Gu03	I	1989	β <sup>-</sup> ?; β <sup>-</sup> n=40#; β <sup>-</sup> 2n=10#
<sup>51</sup> K	-22516	13		365 ms	5	06 14Pa45	J	1983	β <sup>-</sup> =100; β <sup>-</sup> n=65 6; β <sup>-</sup> 2n=4#
<sup>51</sup> Ca	-36332.3	0.5		10.0 s	0.8			1980	β <sup>-</sup> =100; β <sup>-</sup> n=3#
<sup>51</sup> Sc	-43229	20		12.4 s	0.1			1966	β <sup>-</sup> =100; β <sup>-</sup> n=0#
<sup>51</sup> Ti	-49732.8	0.5		5.76 m	0.01			1947	β <sup>-</sup> =100
<sup>51</sup> V	-52203.8	0.4		STABLE				1924	IS=99.750 4
<sup>51</sup> Cr	-51451.4	0.4		27.7010 d	0.0011			1940	ε=100
<sup>51</sup> Cr <sup>i</sup>	-44838	5	6613	5	RQ				
... A-group is continued on next page ...									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...													
<sup>51</sup> Mn	-48243.9	0.5			46.2	m	0.1	5/2 <sup>-</sup>	06 15Ba49 J	1938	$\beta^+=100$		
<sup>51</sup> Mn <sup>i</sup>	-43793.3	1.6	4450.6	1.5	RQ			7/2 <sup>-</sup> T=3/2	06		IT=100	*	
<sup>51</sup> Fe	-40203	9			305.4	ms	2.3	5/2 <sup>-</sup>	06 15Sh16 T	1972	$\beta^+=100$	*	
<sup>51</sup> Co	-27340	50			68.8	ms	1.9	7/2 <sup>-</sup> #	06 07Do17 TD	1987	$\beta^+=100; \beta^+p<3.8$		
<sup>51</sup> Co <sup>i</sup>	-20674	18	6670	50	p			7/2 <sup>-</sup> #T=5/2	07Do17 D		p=100		
<sup>51</sup> Ni	-11900#	500#			23.8	ms	0.2	7/2 <sup>-</sup> #	06 07Do17 TD	1987	$\beta^+=100; \beta^+p=87.2\ 8; \beta^+2p=0.5\ 2$	*	
* <sup>51</sup> K	D : average 06Pe16=63(8)% 83La23=68(10)%; other 82Ca04=47(5)%											**	
* <sup>51</sup> Mn <sup>i</sup>	E : NDS916 gives 4450.0(0.6) may be based on mis-interpretation of 86Di01											**	
* <sup>51</sup> Fe	T : average 15Sh16=308(5) 13Su07=301(4) 87Ha.B=305(5) 84Ay01=310(5)											**	
* <sup>51</sup> Ni	D : $\beta^+2p$ from 12Au08											**	
<sup>52</sup> Ar	-1280#	600#			10#	ms	(>620 ns)	0 <sup>+</sup>	15		2009	$\beta^- ?; \beta^-n=30#; \beta^-2n=7#$	
<sup>52</sup> K	-17140	30			110	ms	4	2 <sup>-</sup> #	15 06Pe16 T	1983	$\beta^-=100; \beta^-n=74\ 9; \beta^-2n=2.3\ 3$	*	
<sup>52</sup> Ca	-34266.3	0.7			4.6	s	0.3	0 <sup>+</sup>	15 83La23 D	1985	$\beta^-=100; \beta^-n<2$		
<sup>52</sup> Sc	-40440	80			8.2	s	0.2	3 <sup>(+)</sup>	15		1980	$\beta^-=100; \beta^-n=4#$	
<sup>52</sup> Ti	-49470	7			1.7	m	0.1	0 <sup>+</sup>	15		1966	$\beta^-=100$	
<sup>52</sup> V	-51443.8	0.4			3.743	m	0.005	3 <sup>+</sup>	15		1934	$\beta^-=100$	
<sup>52</sup> Cr	-55419.2	0.3			STABLE			0 <sup>+</sup>	15		1923	IS=83.789 18	
<sup>52</sup> Cr <sup>i</sup>	-44154.3	0.5	11264.9	0.4				3 <sup>+</sup> T=3	15			IT=100	
<sup>52</sup> Mn	-50707.3	1.8			5.591	d	0.003	6 <sup>+</sup>	15		1938	$\beta^+=100$	
<sup>52</sup> Mn <sup>m</sup>	-50329.6	1.8	377.749	0.005	21.1	m	0.2	2 <sup>+</sup>	15		1937	$\beta^+=98.22\ 5; IT=1.78\ 5$	*
<sup>52</sup> Mn <sup>i</sup>	-47785	5	2922	5	RQ			0 <sup>+</sup> T=2	15			IT=100	
<sup>52</sup> Fe	-48330	5			8.275	h	0.008	0 <sup>+</sup>	15		1948	$\beta^+=100$	*
<sup>52</sup> Fe <sup>m</sup>	-41372	5	6958.0	0.4	45.9	s	0.6	12 <sup>+</sup>	15		1979	$\beta^+\approx 100; IT=0.021\ 5$	
<sup>52</sup> Fe <sup>i</sup>	-42676	5	5654.5	0.4				6 <sup>+</sup> T=1	15			IT=100	
<sup>52</sup> Fe <sup>j</sup>	-39776	6	8555	8	RQ			0 <sup>+</sup> frg.T=2	15				*
<sup>52</sup> Co	-34361	8			111.1	ms	2.3	(6 <sup>+</sup> )	15 16Or08 T	1987	$\beta^+=100; \beta^+p ?$	*	
<sup>52</sup> Co <sup>m</sup>	-33974	10	387	13	MD			2 <sup>+</sup> #	16Or08 T	2016	$\beta^+=100; IT ?; \beta^+p ?$		
<sup>52</sup> Co <sup>i</sup>	-31426	10	2935	13				0 <sup>+</sup> T=2	16Or03 D	2016	IT=75 23; p=?		
<sup>52</sup> Ni	-22330#	400#			41.8	ms	1.0	0 <sup>+</sup>	15 16Or03 TD	1987	$\beta^+=100; \beta^+p=31.1\ 5$	*	
<sup>52</sup> Cu	-2280#	600#						3 <sup>+</sup> #	Mirror I		p ?		
* <sup>52</sup> K	T : average 06Pe16=118(6) 85Hu03=110(30) 83La23=105(5)											**	
* <sup>52</sup> Mn <sup>m</sup>	T : other: 95Ir01=22.7(3.0) for q=25 <sup>+</sup> (bare ion)											**	
* <sup>52</sup> Fe	T : other: 95Ir01=12.5(+1.5-1.2) for q=26 <sup>+</sup> (bare ion)											**	
* <sup>52</sup> Fe <sup>j</sup>	E : probably fragmented, unresolved doublet separated by $\approx 4$ keV											**	
* <sup>52</sup> Co	T : average 16Or08=112(3) 15Sh16=112(4) 13Su07=103(7) other: 97Ha04=104(11)											**	
* <sup>52</sup> Ni	T : average 16Or03=42.8(3) 07Do17=40.8(2); other 94Fa06=38(5)											**	
* <sup>52</sup> Ni	D : other 07Do17=31.4(15) 94Fa06=17.0(14)											**	
<sup>53</sup> Ar	6790#	700#			3#	ms	(>620 ns)	5/2 <sup>-</sup> #	11 09Ta24 I	2009	$\beta^- ?; \beta^-n=20#; \beta^-2n=30#$		
<sup>53</sup> K	-12300	110			30	ms	5	(3/2 <sup>+</sup> )	09 06Pe16 JD	1983	$\beta^-=100; \beta^-n=64\ 11; \beta^-2n\approx 10\ 5$		
<sup>53</sup> Ca	-29390	40			461	ms	90	3/2 <sup>-</sup> #	14		1983	$\beta^-=100; \beta^-n=40\ 10$	
<sup>53</sup> Sc	-38910	90			2.4	s	0.6	(7/2 <sup>-</sup> )	14 10Cr02 TJ	1980	$\beta^-=100; \beta^-n=0.2#$		
<sup>53</sup> Ti	-46830	100			32.7	s	0.9	(3/2 <sup>-</sup> )	09		1977	$\beta^-=100$	
<sup>53</sup> V	-51851	3			1.543	m	0.014	7/2 <sup>-</sup>	09		1960	$\beta^-=100$	
<sup>53</sup> Cr	-55287.0	0.3			STABLE			3/2 <sup>-</sup>	09		1930	IS=9.501 17	
<sup>53</sup> Mn	-54690.1	0.5			3.7	My	0.4	7/2 <sup>-</sup>	09 15Ba49 J	1955	$\epsilon=100$	*	
<sup>53</sup> Mn <sup>i</sup>	-47717	4	6974	4	RQ			3/2 <sup>-</sup> T=5/2	09		1976		
<sup>53</sup> Fe	-50947.5	1.7			8.51	m	0.02	7/2 <sup>-</sup>	09		1938	$\beta^+=100$	*
<sup>53</sup> Fe <sup>m</sup>	-47907.1	1.7	3040.4	0.3	2.54	m	0.02	19/2 <sup>-</sup>	09		1967	IT=100	
<sup>53</sup> Fe <sup>i</sup>	-46698	3	4250	3				7/2 <sup>-</sup> T=3/2	09				
<sup>53</sup> Co	-42659.4	1.7			242	ms	8	7/2 <sup>-</sup> #	09 02Lo13 T	1970	$\beta^+=100$	*	
<sup>53</sup> Co <sup>m</sup>	-39485.2	1.9	3174.2	0.9	MD			(19/2 <sup>-</sup> )	09		1970	$\beta^+\approx 98.5; p\approx 1.5$	*
<sup>53</sup> Co <sup>i</sup>	-38334.4	2.6	4325.0	2.0				(7/2 <sup>-</sup> )T=3/2	09 16Su10 ED	1976	IT $\approx 100; p<0.9\ 3$		
<sup>53</sup> Ni	-29631	25			55.2	ms	0.7	(7/2 <sup>-</sup> )T=3/2	13 16Su10 D	1976	$\beta^+=100; \beta^+p=22.7\ 7$	*	
<sup>53</sup> Cu	-13270#	500#					<130 ns	3/2 <sup>-</sup> #	13			p ?	
* <sup>53</sup> Mn	T : 3.74(0.04) My as given in ENSDF2009 is typo											**	
* <sup>53</sup> Fe	T : other: 95Ir01=8.5(0.3) for q=26 <sup>+</sup> (bare ion)											**	
* <sup>53</sup> Co	T : average 02Lo13=240(9) 89Ho13=240(20) 73Ko10=262(25)											**	
* <sup>53</sup> Co	T : 13Su07=230(17) for which state ?											**	
* <sup>53</sup> Co <sup>m</sup>	D : p $\approx 1.5$ from ENSDF'90											**	
* <sup>53</sup> Ni	D : average 16Su10=22(1) 07Do17=23.4(1.0); other: 76Vi02 $\approx 45$											**	

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)		Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)				
$^{54}\text{K}$	-5000#	600#			10 ms	5	2 <sup>-</sup> #	14		1983	$\beta^-$ =100; $\beta^-$ n=1#; $\beta^-$ 2n=30#			
$^{54}\text{Ca}$	-25160	50			90 ms	6	0 <sup>+</sup>	14	08Ma01	TD	1997	$\beta^-$ =100; $\beta^-$ n=7#; $\beta^-$ 2n=0#	*	
$^{54}\text{Sc}$	-33890	270			526 ms	15	(3 <sup>+</sup> )	14			1990	$\beta^-$ =100; $\beta^-$ n=16 9		
$^{54}\text{Sc}^m$	-33780	270	110.5	0.3	2.77 $\mu$ s	0.02	(5 <sup>+</sup> , 4 <sup>+</sup> )	14	10Cr02	J	1998	IT=100		
$^{54}\text{Ti}$	-45620	80			2.1 s	1.0	0 <sup>+</sup>	14			1980	$\beta^-$ =100		
$^{54}\text{V}$	-49893	15			49.8 s	0.5	3 <sup>+</sup>	14			1970	$\beta^-$ =100		
$^{54}\text{V}^m$	-49785	15	108.0	1.0	900 ns	500	(5 <sup>+</sup> )	14			1998	IT=100		
$^{54}\text{Cr}$	-56934.8	0.4			STABLE		0 <sup>+</sup>	14			1930	IS=2.365 7		
$^{54}\text{Mn}$	-55557.6	1.1			312.20 d	0.20	3 <sup>+</sup>	14			1938	$\epsilon$ =100; $\beta^-$ =0.93e-4; $e^+$ =1.28e-7 25	*	
$^{54}\text{Mn}^i$	-49411.5	2.8	6146.2	3.0	RQ		0 <sup>+</sup> T=3							
$^{54}\text{Fe}$	-56254.5	0.4			STABLE		0 <sup>+</sup>	14			1923	IS=5.845 35; 2 $\beta^+$ ?		
$^{54}\text{Fe}^m$	-49727.4	1.2	6527.1	1.1	364 ns	7	10 <sup>+</sup>	14			1983	IT=100		
$^{54}\text{Fe}^j$	-41386	20	14868	20	RQ		0 <sup>+</sup> T=3	14						
$^{54}\text{Co}$	-48010.0	0.4			193.28 ms	0.07	0 <sup>+</sup> T=1	14			1952	$\beta^+$ =100		
$^{54}\text{Co}^m$	-47812.4	0.4	197.57	0.10	MD		7 <sup>+</sup> T=0	14			1962	$\beta^+$ =100		
$^{54}\text{Ni}$	-39278	5			114.2 ms	0.3	0 <sup>+</sup>	14			1977	$\beta^+$ =100; $\beta^+$ p ?		
$^{54}\text{Ni}^m$	-32821	5	6457.4	0.9	152 ns	4	10 <sup>+</sup>	14	08Ru09	JD	2008	IT=64 2; p=36 2		
$^{54}\text{Cu}$	-21410#	400#			<75 ns		3 <sup>+</sup> #	14				p ?		
$^{54}\text{Zn}$	-6270#	400#			1.8 ms	0.5	0 <sup>+</sup>	14	11As08	TD	2005	2p=87 7	*	
* $^{54}\text{Ca}$	T : average 10Cr02=107(14) 08Ma01=86(7)											**		
* $^{54}\text{Mn}$	D : e <sup>+</sup> average 98Wu01=1.20(0.26)e-7% 97Za07=2.2(0.9)e-7%											**		
* $^{54}\text{Zn}$	T : symmetrized from 11As08=1.59(+0.60-0.35); other 05B115=3.2(+1.8-0.8)											**		
* $^{54}\text{Zn}$	D : averaged from 11As08=92(+6-13)% 05B115=87(+10-17)%											**		
$^{55}\text{K}$	710#	700#			3# ms	(>620 ns)	3/2 <sup>+</sup> #	09	09Ta24	I	2009	$\beta^-$ ?; $\beta^-$ n=40#; $\beta^-$ 2n=1#		
$^{55}\text{Ca}$	-18350#	300#			22 ms	2	5/2 <sup>-</sup> #	09			1997	$\beta^-$ =100; $\beta^-$ n=1#; $\beta^-$ 2n=0.4#		
$^{55}\text{Sc}$	-30160	450			96 ms	2	(7/2) <sup>-</sup>	08	10Cr02	TJD	1990	$\beta^-$ =100; $\beta^-$ n=17 7; $\beta^-$ 2n=0#	*	
$^{55}\text{Ti}$	-41670	160			1.3 s	0.1	(1/2) <sup>-</sup>	10			1980	$\beta^-$ =100; $\beta^-$ n=0#		
$^{55}\text{V}$	-49140	100			6.54 s	0.15	7/2 <sup>-</sup> #	08			1977	$\beta^-$ =100		
$^{55}\text{Cr}$	-55109.7	0.4			3.497 m	0.003	3/2 <sup>-</sup>	08			1952	$\beta^-$ =100		
$^{55}\text{Mn}$	-57712.4	0.3			STABLE		5/2 <sup>-</sup>	08	15Ba49	J	1923	IS=100.		
$^{55}\text{Fe}$	-57481.3	0.3			2.744 y	0.009	3/2 <sup>-</sup>	09			1939	$\epsilon$ =100		
$^{55}\text{Fe}^i$	-49848	6	7633	6	RQ		5/2 <sup>-</sup> T=5/2	09						
$^{55}\text{Co}$	-54029.9	0.4			17.53 h	0.03	7/2 <sup>-</sup>	09			1938	$\beta^+$ =100		
$^{55}\text{Co}^j$	-49308.5	0.4	4721.44	0.10			3/2 <sup>-</sup> frg.T=3/2	09			1981	IT=100	*	
$^{55}\text{Ni}$	-45335.8	0.7			204.7 ms	1.7	7/2 <sup>-</sup>	08	02Lo13	T	1972	$\beta^+$ =100	*	
$^{55}\text{Ni}^i$	-40736.8	1.2	4599	1			7/2 <sup>-</sup> frg.T=3/2	13	Tr09	E			*	
$^{55}\text{Cu}$	-31640	160			57 ms	3	3/2 <sup>-</sup> #	08	13Tr09	T	1987	$\beta^+$ =100; $\beta^+$ p=15.0 43	*	
$^{55}\text{Zn}$	-14570#	400#			19.8 ms	1.3	5/2 <sup>-</sup> #	08	07Do17	TD	2001	$\beta^+$ =100; $\beta^+$ p=91.0 51		
* $^{55}\text{Sc}$	T : others 04Li75=115(15) 02Sh43=103(7) 98So03=120(40)											**		
* $^{55}\text{Co}^j$	E : strongest frg (spectr. factor 0.45); other 26.69(0.15) higher (sf=0.37)											**		
* $^{55}\text{Ni}$	T : average 02Lo13=196(5) 99Re06=204(3) 87Ha.A=212.1(3.8) 84Ay01=208(5)											**		
* $^{55}\text{Ni}$	T : and 77Ho25=189(5) 76Ed.A=219(6); 97Wo06=204(3) superseded by 99Re06											**		
* $^{55}\text{Ni}$	J : spectroscopy factor information in 14Sa46											**		
* $^{55}\text{Ni}^i$	E : strongest frg (total strength 2.0); other 20keV lower (tot.str 0.8)											**		
* $^{55}\text{Cu}$	T : 07Do17 27(8)ms poor statistics D : from 07Do17											**		
$^{56}\text{K}$	7930#	800#			1# ms	(>620 ns)	2 <sup>-</sup> #	11	09Ta24	I	2009	$\beta^-$ ?; $\beta^-$ n=50#; $\beta^-$ 2n=40#		
$^{56}\text{Ca}$	-13900#	400#			11 ms	2	0 <sup>+</sup>	11			1997	$\beta^-$ =100; $\beta^-$ n=5#; $\beta^-$ 2n=0.2#		
$^{56}\text{Sc}$	-24850	590			26 ms	6	(1 <sup>+</sup> )	11	10Cr02	J	1997	$\beta^-$ =100; $\beta^-$ n=10#; $\beta^-$ 2n=0.5#		
$^{56}\text{Sc}^m$	-24850#	600#	0#	100#	75 ms	6	(6 <sup>+</sup> , 5 <sup>+</sup> )	11	10Cr02	J	2004	$\beta^-$ =100; $\beta^-$ n>14 2; $\beta^-$ 2n=0.5#		
$^{56}\text{Sc}^n$	-24080	590	774.9	0.3	290 ns	30	(4 <sup>+</sup> )	11			2004	IT=100	*	
$^{56}\text{Ti}$	-39320	120			200 ms	5	0 <sup>+</sup>	11	98Am04	D	1980	$\beta^-$ =100; $\beta^-$ n=0.1#		
$^{56}\text{V}$	-46150	180			216 ms	4	(1 <sup>+</sup> )	11	98Am04	D	1980	$\beta^-$ =100; $\beta^-$ n=0#		
$^{56}\text{Cr}$	-55285.0	0.6			5.94 m	0.10	0 <sup>+</sup>	11	60Dr03	D	1960	$\beta^-$ =100		
$^{56}\text{Mn}$	-56911.5	0.3			2.5789 h	0.0001	3 <sup>+</sup>	11			1934	$\beta^-$ =100		
$^{56}\text{Fe}$	-60607.1	0.3			STABLE		0 <sup>+</sup>	11			1923	IS=91.754 36		
$^{56}\text{Fe}^i$	-49103.4	0.4	11503.7	0.3			3 <sup>+</sup> T=3	11						
$^{56}\text{Co}$	-56040.4	0.5			77.236 d	0.026	4 <sup>+</sup>	11			1941	$\beta^+$ =100		
$^{56}\text{Co}^j$	-52448	9	3593	9	RQ		(0 <sup>+</sup> ) frg.T=2	11						
$^{56}\text{Ni}$	-53907.5	0.4			6.075 d	0.010	0 <sup>+</sup>	11			1952	$\beta^+$ =100	*	
$^{56}\text{Ni}^i$	-47475.6	0.8	6431.9	0.7			4 <sup>+</sup> T=1	11						
$^{56}\text{Ni}^j$	-43964	4	9944	4	RQ		0 <sup>+</sup> frg.T=2	11						*
$^{56}\text{Cu}$	-38643	15			93 ms	3	(4 <sup>+</sup> )	11	01Bo54	TJD	1987	$\beta^+$ =100; $\beta^+$ p=0.40 12		
$^{56}\text{Cu}^i$	-35099	10	3544	18	p		T=2	16	Or03	D	2007	IT=56 6; p=46 6		

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>56</sup> Zn	-25390#	400#	32.9 ms	0.8	0 <sup>+</sup>	11 14Or04	TD 2001	$\beta^+=100; \beta^+p=88.0$	23 *
<sup>56</sup> Zn <sup>i</sup>	-21530#	650#	3860#	510#	3 <sup>+</sup> #T=3			p ?	
<sup>56</sup> Ga	-3390#	500#			3 <sup>+</sup> #			p ?	
* <sup>56</sup> Sc <sup>n</sup>	T : other 12Ka36=350(+260-120)								
* <sup>56</sup> Co <sup>j</sup>	E : strongest frg (cross section 115); other 70(9) keV lower (xs=55)								
* <sup>56</sup> Ni <sup>j</sup>	E : strongest frg; others 68(6) and 98(6) keV higher								
* <sup>56</sup> Zn	T : other 07Do17=30.0(1.7)								
* <sup>56</sup> Zn	D : average 14Or04=88.5(26) 07Do17=86.0(49)								
<sup>57</sup> Ca	-6870#	400#	5# ms	(>620 ns)	5/2 <sup>-</sup> #	10 09Ta24	I 2009	$\beta^- ?; \beta^-n=20\#; \beta^-2n=2\#$	
<sup>57</sup> Sc	-21000	1300	22 ms	2	7/2 <sup>-</sup> #	10 10Cr02	T 1997	$\beta^- =100; \beta^-n=30\#; \beta^-2n=1\#$	*
<sup>57</sup> Ti	-33920	260	95 ms	8	5/2 <sup>-</sup> #	10 99So20	T 1985	$\beta^- =100; \beta^-n=0.04\#$	*
<sup>57</sup> V	-44410	80	350 ms	10	(7/2 <sup>-</sup> )	10 03Ma02	T 1980	$\beta^- =100; \beta^-n=0.4\#$	*
<sup>57</sup> Cr	-52524.7	1.1	21.1 s	1.0	(3/2 <sup>-</sup> )	10	1978	$\beta^- =100$	
<sup>57</sup> Mn	-57486.3	1.5	85.4 s	1.8	5/2 <sup>-</sup>	98 15Ba49	J 1954	$\beta^- =100$	
<sup>57</sup> Fe	-60181.8	0.3	STABLE		1/2 <sup>-</sup>	98	1935	IS=2.119 10	
<sup>57</sup> Co	-59345.6	0.5	271.70 d	0.10	7/2 <sup>-</sup>	98 14Un01	T 1941	$\epsilon=100$	*
<sup>57</sup> Co <sup>j</sup>	-52092.3	0.4	7253.3	0.6	RQ	1/2 <sup>-</sup> T=5/2	MMC120	J	
<sup>57</sup> Ni	-56083.8	0.6	35.60 h	0.06	3/2 <sup>-</sup>	98	1938	$\beta^+=100$	
<sup>57</sup> Ni <sup>i</sup>	-50845.0	0.9	5238.8	0.7	7/2 <sup>-</sup> frg.T=3/2	98			*
<sup>57</sup> Cu	-47308.9	0.5	196.3 ms	0.7	3/2 <sup>-</sup>	98	1976	$\beta^+=100$	
<sup>57</sup> Cu <sup>i</sup>	-42010	25	5299	25	p	7/2 <sup>-</sup> T=3/2			
<sup>57</sup> Zn	-32550#	200#	38 ms	4	7/2 <sup>-</sup> #	98 02Lo13	T 1976	$\beta^+=100; \beta^+p\approx 65$	*
<sup>57</sup> Ga	-15010#	400#			1/2 <sup>-</sup> #			p ?	
* <sup>57</sup> Sc	T : other 03So21=13(4)								
* <sup>57</sup> Ti	T : average 05Li53=98(5) 99So20=67(25) 96Do23=56(20)								
* <sup>57</sup> Ti	T : 98Am04=180(30) conflicting, not used								
* <sup>57</sup> V	J : 98So03 proposed 3/2 <sup>-</sup> , supported in 03Ma02; same group 05Li53 favors 7/2 <sup>-</sup>								
* <sup>57</sup> Co	T : average 14Un01=271.87(0.44) (supersedes 92Un01=272.11(0.26)),								
* <sup>57</sup> Co	T : 12Da06=271.82(0.17) 97Ma75=271.68(0.27) 83Wa26=271.84(0.27)								
* <sup>57</sup> Co	T : 81Va11=270.90(0.27) 80Ho17=271.77(0.27) 72La14=271.23(0.21)								
* <sup>57</sup> Co	T : 65An07=271.65(0.13); original unc of 97Ma75=0.09 83Wa26=0.04								
* <sup>57</sup> Co	T : 81Va11=0.09 80Ho17=0.05 increased to 0.1% by evaluator								
* <sup>57</sup> Ni <sup>i</sup>	E : strongest frg; 79Ik04 others 98(7)keV lower(5.5%) 128(7)keV higher(10.0%)								
* <sup>57</sup> Ni <sup>i</sup>	E : strongest frg; 78Na11 others 104(5)keV lower, 129(5)keV higher								
* <sup>57</sup> Zn	T : average 02Lo13=37(5) 76Vi02=40(10)								
<sup>58</sup> Ca	-1920#	500#	3# ms	(>620 ns)	0 <sup>+</sup>	10	2009	$\beta^- ?; \beta^-n=2\#; \beta^-2n=4\#$	
<sup>58</sup> Sc	-14880#	400#	12 ms	5	3 <sup>+</sup> #	10	1997	$\beta^- =100; \beta^-n=20\#; \beta^-2n=1\#$	*
<sup>58</sup> Ti	-31110#	200#	55 ms	6	0 <sup>+</sup>	14 11Da08	T 1992	$\beta^- =100; \beta^-n=1\#$	*
<sup>58</sup> V	-40400	90	191 ms	10	(1 <sup>+</sup> )	10	1980	$\beta^- =100; \beta^-n=0.8\#$	
<sup>58</sup> Cr	-51991.8	1.5	7.0 s	0.3	0 <sup>+</sup>	10	1980	$\beta^- =100$	
<sup>58</sup> Mn	-55827.6	2.7	3.0 s	0.1	1 <sup>+</sup>	10	1961	$\beta^- =100$	
<sup>58</sup> Mn <sup>m</sup>	-55755.8	2.7	71.77	0.05	65.4 s	0.5	1961	$\beta^- =?; IT=20\#$	
<sup>58</sup> Fe	-62155.1	0.3	STABLE		0 <sup>+</sup>	10	1935	IS=0.282 4	
<sup>58</sup> Co	-59847.2	1.2	70.86 d	0.06	2 <sup>+</sup>	10	1941	$\beta^+=100$	
<sup>58</sup> Co <sup>m</sup>	-59822.3	1.2	24.95	0.06	9.10 h	0.09	1950	IT=100	
<sup>58</sup> Co <sup>n</sup>	-59794.1	1.2	53.15	0.07	10.5 $\mu$ s	0.3	1964	IT=100	
<sup>58</sup> Co <sup>i</sup>	-54095	8	5752	8	RQ	0 <sup>+</sup> frg.T=3	10		*
<sup>58</sup> Ni	-60228.7	0.4	STABLE	(>700 Ey)	0 <sup>+</sup>	10	1921	IS=68.077 19; 2 $\beta^+$ ?	*
<sup>58</sup> Ni <sup>i</sup>	-51400	40	8830	40	RQ	2 <sup>+</sup> T=2	10		
<sup>58</sup> Ni <sup>j</sup>	-45690	7	14539	7	RQ	0 <sup>+</sup> T=3	10	MMC12	J
<sup>58</sup> Cu	-51667.7	0.6	3.204 s	0.007	1 <sup>+</sup> T=0	10	1952	$\beta^+=100$	
<sup>58</sup> Cu <sup>i</sup>	-51464.7	0.6	202.99	0.24	0 <sup>+</sup> T=1	10			
<sup>58</sup> Zn	-42300	50	86.7 ms	2.4	0 <sup>+</sup>	14	1986	$\beta^+=100; \beta^+p<3$	
<sup>58</sup> Ga	-23540#	300#			2 <sup>+</sup> #	Mirror	I	p ?	
<sup>58</sup> Ga <sup>m</sup>	-23510#	320#	30#	100#	*	5 <sup>+</sup> #	Mirror	I	p ?
<sup>58</sup> Ge	-7080#	500#			0 <sup>+</sup>	Mirror	I	2p ?	
* <sup>58</sup> Ti	T : average 11Da08=57(10) 03So21=59(9) 99So20=47(10)								
* <sup>58</sup> Co <sup>j</sup>	E : strongest fragment (cross section 98); other 20(8) keV lower (xs=90)								
* <sup>58</sup> Ni	T : >400 Ey to 2 <sup>+</sup> level of <sup>58</sup> Fe, >700 Ey to ground-state								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>59</sup> Sc	-10300#	400#		10# ms (>620 ns)	7/2 <sup>-</sup> #	09	09Ta24 I	2009	$\beta^-$ ?; $\beta^-$ -n=50#; $\beta^-$ -2n=1#
<sup>59</sup> Ti	-25510#	200#		28.5 ms 1.9	5/2 <sup>-</sup> #	02	11Da08 T	1997	$\beta^-$ =100; $\beta^-$ -n=0.3#; $\beta^-$ -2n=0.01# *
<sup>59</sup> Ti <sup>m</sup>	-25400#	200#	109.0 0.5	590 ns 50	(1/2 <sup>-</sup> )		12Ka36 ETJ	2012	IT=100 *
<sup>59</sup> V	-37830	160		95 ms 6	(5/2 <sup>-</sup> )	02	05Li53 TJ	1985	$\beta^-$ =100; $\beta^-$ -n=6# *
<sup>59</sup> Cr	-48090	220		1050 ms 90	(1/2 <sup>-</sup> )	02	05Li53 TJ	1980	$\beta^-$ =100 *
<sup>59</sup> Cr <sup>m</sup>	-47590	220	503.0 1.7	96 $\mu$ s 20	(9/2 <sup>+</sup> )	02		1998	IT=100
<sup>59</sup> Mn	-55525.3	2.3		4.59 s 0.05	5/2 <sup>-</sup>	02	15Ba49 J	1976	$\beta^-$ =100
<sup>59</sup> Fe	-60664.8	0.4		44.495 d 0.009	3/2 <sup>-</sup>	02		1938	$\beta^-$ =100
<sup>59</sup> Co	-62229.7	0.4		STABLE	7/2 <sup>-</sup>	02		1923	IS=100.
<sup>59</sup> Ni	-61156.7	0.4		81 ky 5	3/2 <sup>-</sup>	02	94Ru19 T	1951	$\beta^+$ =100 *
<sup>59</sup> Ni <sup>i</sup>	-53814.8	2.1	7341.9 2.1 RQ		7/2 <sup>-</sup> frg, T=5/2				
<sup>59</sup> Cu	-56358.3	0.5		81.5 s 0.5	3/2 <sup>-</sup>	02		1947	$\beta^+$ =100 *
<sup>59</sup> Cu <sup>i</sup>	-52472.8	2.2	3885.5 2.1		3/2 <sup>-</sup> frg, T=3/2	02			IT=100 *
<sup>59</sup> Zn	-47215.6	0.8		182.0 ms 1.8	3/2 <sup>-</sup>	15		1981	$\beta^+$ =100; $\beta^+$ -p=0.10 3
<sup>59</sup> Ga	-33760#	170#		<43 ns	3/2 <sup>-</sup> #	15			p? *
<sup>59</sup> Ge	-15870#	400#		8# ms (>620 ns)	7/2 <sup>-</sup> #	15	15Ci06 IT	2015	$\beta^+$ ? *
<sup>59</sup> Ti	T: average 11Da08=27.5(2.5) 03So21=30(3); other 99So20=58(17) **								
<sup>59</sup> Ti <sup>m</sup>	T: symmetrized from 587(+57-51) **								
<sup>59</sup> V	T: average 05Li53=97(2) 99So20=75(7) (supersedes 98So03=70(40)) **								
<sup>59</sup> V	T: 98Am04=130(20) conflicting, not used **								
<sup>59</sup> Cr	T: others 96Do23=460(50), 88Bo06=600(300), 85Bo49=1000(400) **								
<sup>59</sup> Ni	T: average 94Ru19=108(13) 94Ru19(meteorite)=120(22) 81Ni08=76(5) **								
<sup>59</sup> Ni <sup>i</sup>	E: strongest frg(100%); 3 others 40.1(0.3)keV higher (0.140%), 17.7(0.3)keV **								
<sup>59</sup> Ni <sup>i</sup>	E: higher (0.122%) and 36.3(0.2)keV lower (0.110%) **								
<sup>59</sup> Cu <sup>i</sup>	E: 76Ga19 strongest fragment (sp.factor 0.6); other 21(6) (sf 0.4) higher **								
<sup>60</sup> Sc	-4050#	500#		3# ms (>620 ns)	3+#		09Ta24 I	2009	$\beta^-$ ?; $\beta^-$ -n=0.4#; $\beta^-$ -2n=50#
<sup>60</sup> Ti	-22330#	300#		22.2 ms 1.6	0 <sup>+</sup>	14	11Da08 T	1997	$\beta^-$ =100; $\beta^-$ -n=2#; $\beta^-$ -2n=0# *
<sup>60</sup> V	-33240	220		122 ms 18	3+#	13		1985	$\beta^-$ =100; $\beta^-$ -n=5#; $\beta^-$ -2n=0.03#
<sup>60</sup> V <sup>m</sup>	-33240#	270#	0# 150# *	40 ms 15	1+#	13		1999	$\beta^-$ =?; IT?; $\beta^-$ -n=5#; $\beta^-$ -2n=0.03#
<sup>60</sup> V <sup>n</sup>	-33040	220	203.7 0.7	230 ns 24	(4 <sup>+</sup> )	13	12Ka36 ET	1999	IT=100 *
<sup>60</sup> Cr	-46670	190		490 ms 10	0 <sup>+</sup>	13		1980	$\beta^-$ =100; $\beta^-$ -n=0#
<sup>60</sup> Mn	-52967.9	2.3		280 ms 20	1 <sup>+</sup>	13		1978	$\beta^-$ =100
<sup>60</sup> Mn <sup>m</sup>	-52696.0	2.3	271.90 0.10	1.77 s 0.02	4 <sup>+</sup>	13		1978	$\beta^-$ =88.5 8; IT=11.5 8 *
<sup>60</sup> Fe	-61413	3		2.62 My 0.04	0 <sup>+</sup>	13		1957	$\beta^-$ =100 *
<sup>60</sup> Co	-61650.3	0.4		5.2712 y 0.0004	5 <sup>+</sup>	13		1941	$\beta^-$ =100
<sup>60</sup> Co <sup>m</sup>	-61591.7	0.4	58.59 0.01	10.467 m 0.006	2 <sup>+</sup>	13		1963	IT $\approx$ 100; $\beta^-$ =0.25 3
<sup>60</sup> Ni	-64473.1	0.4		STABLE	0 <sup>+</sup>	13		1921	IS=26.223 15
<sup>60</sup> Ni <sup>i</sup>	-53347	4	11126 4 RQ		5 <sup>+</sup> T=3				
<sup>60</sup> Cu	-58345.1	1.6		23.7 m 0.4	2 <sup>+</sup>	13		1947	$\beta^+$ =100
<sup>60</sup> Cu <sup>i</sup>	-55804	5	2541 5 RQ		(0 <sup>+</sup> )T=2	13			IT=100
<sup>60</sup> Zn	-54174.3	0.6		2.38 m 0.05	0 <sup>+</sup>	13		1955	$\beta^+$ =100
<sup>60</sup> Zn <sup>i</sup>	-49322.1	0.9	4852.2 0.7		(2 <sup>+</sup> )T=1	13			IT=100
<sup>60</sup> Zn <sup>j</sup>	-46807	24	7367 24 RQ		0 <sup>+</sup> T=2	13			
<sup>60</sup> Ga	-39590#	200#		70 ms 10	(2 <sup>+</sup> )	13	01Ma96 TJ	1995	$\beta^+$ =100; $\beta^+$ -p=1.6 7; $\beta^+$ - $\alpha$ <0.023 20 *
<sup>60</sup> Ga <sup>i</sup>	-37050#	210#	2540# 50#						
<sup>60</sup> Ge	-27090#	300#		30# ms (>110 ns)	0 <sup>+</sup>	13		2005	$\beta^+$ ?; $\beta^+$ -p?
<sup>60</sup> As	-5470#	400#			5 <sup>+</sup> #		Mirror I		p?
<sup>60</sup> As <sup>m</sup>	-5410#	400#	60# 20#		2 <sup>+</sup> #		Mirror I		p?
<sup>60</sup> Ti	T: average 11Da08=22.4(2.5) 03So21=22(2) **								
<sup>60</sup> V <sup>n</sup>	E: 12Ka36=99.7(0.5) and 104.0(0.5) $\gamma$ rays in cascade to ground-state **								
<sup>60</sup> V <sup>n</sup>	T: symmetrized from 12Ka36=229(+25-23); others 10Da06=320(90) 99Da.A=320(90) **								
<sup>60</sup> Mn <sup>m</sup>	I: also an isomer T=1.0(+0.3-0.2) $\mu$ s decay by 114 keV $\gamma$ -rays to ground-state or <sup>60</sup> Mn <sup>m</sup> **								
<sup>60</sup> Fe	T: 15Wa06=2.50(0.12) confirms 09Ru08=2.62(0.04), rules out 84Ku28=1.49(27) **								
<sup>60</sup> Fe	T: and 57Ro54=0.3 **								
<sup>60</sup> Ga	T: average 02Lo13=70(13) 01Ma96=70(15) **								
<sup>61</sup> Sc	930#	600#		2# ms (>620 ns)	7/2 <sup>-</sup> #	15		2009	$\beta^-$ ?; $\beta^-$ -n=60#; $\beta^-$ -2n=1#
<sup>61</sup> Ti	-16350#	400#		15 ms 4	1/2 <sup>-</sup> #	15		1997	$\beta^-$ =100; $\beta^-$ -n=1#; $\beta^-$ -2n=1#
<sup>61</sup> V	-30510	890		48.2 ms 0.8	(3/2 <sup>-</sup> , 5/2 <sup>-</sup> )	15		1992	$\beta^-$ =100; $\beta^-$ -n>10; $\beta^-$ -2n=0.01#
<sup>61</sup> Cr	-42480	100		243 ms 9	(5/2 <sup>-</sup> )	15	09Cr02 T	1985	$\beta^-$ =100; $\beta^-$ -n=0.6#
<sup>61</sup> Mn	-51742.1	2.3		709 ms 8	5/2 <sup>-</sup>	15	15Ba49 J	1980	$\beta^-$ =100; $\beta^-$ -n=0.6# *
<sup>61</sup> Fe	-58920.5	2.6		5.98 m 0.06	(3/2 <sup>-</sup> )	15		1957	$\beta^-$ =100
<sup>61</sup> Fe <sup>m</sup>	-58058.8	2.6	861.67 0.11	238 ns 5	9/2 <sup>+</sup>	15		1998	IT=100
<sup>61</sup> Co	-62898.1	0.8		1.649 h 0.005	7/2 <sup>-</sup>	15		1947	$\beta^-$ =100
<sup>61</sup> Ni	-64221.9	0.4		STABLE	3/2 <sup>-</sup>	15		1934	IS=1.1399 13
<sup>61</sup> Cu	-61984.1	1.0		3.339 h 0.008	3/2 <sup>-</sup>	15	10Vi07 J	1937	$\beta^+$ =100 *
<sup>61</sup> Cu <sup>i</sup>	-55610	7	6374 7 RQ		3/2 <sup>-</sup> frg, T=5/2				
<sup>61</sup> Zn	-56349	16		89.1 s 0.2	3/2 <sup>-</sup>	15		1955	$\beta^+$ =100
<sup>61</sup> Zn <sup>i</sup>	-53190#	100#	3160# 100#		3/2 <sup>-</sup> #T=3/2				
<sup>61</sup> Zn <sup>j</sup>	-46360	70	9990 70		3/2 <sup>-</sup> -T=5/2	15			

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>61</sup> Ga	-47130	40	167 ms 3	3/2 <sup>-</sup>	15		1987	$\beta^+$ =100; $\beta^+$ p<0.25
<sup>61</sup> Ga <sup>m</sup>	-47040#	110#	90# 100#	1/2 <sup>-</sup> #		Mirror I		
<sup>61</sup> Ga <sup>i</sup>	-43780	30	3360 50	(3/2 <sup>-</sup> )T=3/2	15		1987	p=100
<sup>61</sup> Ge	-33360#	300#	44 ms 6	3/2 <sup>-</sup> #	15		1987	$\beta^+$ =100; $\beta^+$ p>62
<sup>61</sup> As	-16900#	300#		3/2 <sup>-</sup> #		Mirror I		p ?
* <sup>61</sup> Mn	D : $\beta^-$ n has been observed by 99Ha05; 13Ra17 quotes $\beta^-$ n=0.6(0.1)% unpubl.							
* <sup>61</sup> Cu	J : directly measured in 10Vi07							
* <sup>61</sup> Cu <sup>i</sup>	E : strongest frg (xs=55); other 18(7) keV higher (xs=35)							
<sup>62</sup> Ti	-12500#	400#	10# ms (>620 ns)	0 <sup>+</sup>	12		2009	$\beta^-$ ?; $\beta^-$ n=4#; $\beta^-$ 2n=0.1#
<sup>62</sup> V	-25480#	300#	33.6 ms 2.3	3 <sup>+</sup> #	12		1997	$\beta^-$ =100; $\beta^-$ n=20#; $\beta^-$ 2n=0.5#
<sup>62</sup> Cr	-40890	150	206 ms 12	0 <sup>+</sup>	12		1985	$\beta^-$ =100; $\beta^-$ n=1#
<sup>62</sup> Mn	-48524	7	92 ms 13	1 <sup>+</sup>	12	15He28 J	1983	$\beta^-$ =100; $\beta^-$ n=0.03#
<sup>62</sup> Mn <sup>m</sup>	-48181.0	2.6	343 6	4 <sup>+</sup>	12	15He28 J	1983	$\beta^-$ =100; $\beta^-$ n=0.03#; IT ?
<sup>62</sup> Fe	-58878.0	2.8	671 ms 5	0 <sup>+</sup>	12		1975	$\beta^-$ =100
<sup>62</sup> Co	-61424	19	1.54 m 0.10	(2) <sup>+</sup>	12		1949	$\beta^-$ =100
<sup>62</sup> Co <sup>m</sup>	-61402	20	22 5	(5) <sup>+</sup>	12		1957	$\beta^-$ >99; IT<1
<sup>62</sup> Ni	-66746.3	0.4	STABLE	0 <sup>+</sup>	12		1934	IS=3.6346 40
<sup>62</sup> Cu	-62787.4	0.6	9.67 m 0.03	1 <sup>+</sup>	12	10Vi07 J	1936	$\beta^+$ =100
<sup>62</sup> Cu <sup>i</sup>	-58174	6	4614 6	(0) <sup>+</sup> T=3	12			
<sup>62</sup> Zn	-61168.0	0.6	9.193 h 0.015	0 <sup>+</sup>	12		1948	$\beta^+$ =100
<sup>62</sup> Ga	-51986.9	0.6	116.121 ms 0.021	0 <sup>+</sup> T=1	12		1978	$\beta^+$ =100
<sup>62</sup> Ga <sup>j</sup>	-51415.7	0.6	571.2 0.1	1 <sup>(+)</sup> T=2	12	98Vi06 EJ	1998	IT=100
<sup>62</sup> Ge	-41740#	140#	129 ms 35	0 <sup>+</sup>	12		1991	$\beta^+$ =100; $\beta^+$ p ?
<sup>62</sup> As	-24320#	300#		1 <sup>+</sup> #				p=100#
* <sup>62</sup> Mn	D : $\beta^-$ n 99So20≈0 99Ha05>0							
* <sup>62</sup> Cu	J : directly measured in 10Vi07							
* <sup>62</sup> Cu <sup>i</sup>	E : ENSDF=4628(10)							
* <sup>62</sup> Ga	T : also 13Da16=116.15(0.13) no weight							
* <sup>62</sup> As	D : most probably p-unstable from estimated $S_p$ =-1860#(420#) keV							
<sup>63</sup> Ti	-5750#	500#	3# ms (>620 ns)	1/2 <sup>-</sup> #	09	09Ta24 I	2009	$\beta^-$ ?; $\beta^-$ n=7#; $\beta^-$ 2n=4#
<sup>63</sup> V	-21890#	400#	19.6 ms 0.9	(3/2 <sup>-</sup> ,5/2 <sup>-</sup> )	09	14Su07 TJ	1997	$\beta^-$ =100; $\beta^-$ n>35; $\beta^-$ 2n=0.2#
<sup>63</sup> Cr	-36010	360	129 ms 2	1/2 <sup>-</sup> #	09		1992	$\beta^-$ =100; $\beta^-$ n=1#
<sup>63</sup> Mn	-46887	4	275 ms 4	5/2 <sup>-</sup>	09	15Ba49 J	1985	$\beta^-$ =100; $\beta^-$ n=0.2#
<sup>63</sup> Fe	-55636	4	6.1 s 0.6	(5/2 <sup>-</sup> )	09		1980	$\beta^-$ =100
<sup>63</sup> Co	-61851	19	26.9 s 0.4	7/2 <sup>-</sup>	09	94It.A T	1960	$\beta^-$ =100
<sup>63</sup> Ni	-65512.8	0.4	101.2 y 1.5	1/2 <sup>-</sup>	09		1951	$\beta^-$ =100
<sup>63</sup> Ni <sup>m</sup>	-65425.7	0.4	87.15 0.11	5/2 <sup>-</sup>	09		1978	IT=100
<sup>63</sup> Cu	-65579.8	0.4	STABLE	3/2 <sup>-</sup>	09	10Vi07 J	1923	IS=69.15 15
<sup>63</sup> Zn	-62213.4	1.6	38.47 m 0.05	3/2 <sup>-</sup>	09		1937	$\beta^+$ =100
<sup>63</sup> Zn <sup>i</sup>	-56723	6	5490 6	3/2 <sup>-</sup> T=5/2	09			
<sup>63</sup> Ga	-56547.1	1.3	32.4 s 0.5	3/2 <sup>(-)</sup>	09	12Pr11 J	1965	$\beta^+$ =100
<sup>63</sup> Ge	-46920	40	142 ms 8	3/2 <sup>-</sup> #	09	02Lo13 TD	1991	$\beta^+$ =100; $\beta^+$ p ?
<sup>63</sup> As	-33500#	200#	<43 ns	3/2 <sup>-</sup> #	09			p=100#
* <sup>63</sup> V	T : average 14Su07=20(1) 11Da08=19.2(2.4) 03So02=17(3)							
* <sup>63</sup> Cr	T : other 11Da08=128(8)							
* <sup>63</sup> Mn	D : $\beta^-$ n has been observed by 99Ha05 but not quantified							
* <sup>63</sup> Co	T : average 94It.A=26.41(0.27) 72Jo08=27.5(0.3) 69Wa15=26(1)							
* <sup>63</sup> Cu	J : directly measured in 10Vi07							
* <sup>63</sup> Ge	T : average 02Lo13=150(9) 93Wi03=95(+23-20)							
* <sup>63</sup> As	D : most probably p-unstable from estimated $S_p$ =-980#(240#) keV							
<sup>64</sup> Ti	-1030#	600#	4# ms (>620 ns)	0 <sup>+</sup>	13		2013	$\beta^-$ ?; $\beta^-$ n=90#; $\beta^-$ 2n=2#
<sup>64</sup> V	-16320#	400#	15 ms 2	(1,2)	14		1997	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=4#
<sup>64</sup> V <sup>m</sup>	-16240#	400#	81.0 0.7	< 1 $\mu$ s			2014	IT≈100
<sup>64</sup> Cr	-33480	440	43 ms 1	0 <sup>+</sup>	14		1992	$\beta^-$ =100; $\beta^-$ n=2#
<sup>64</sup> Mn	-42989	4	88.8 ms 2.4	1 <sup>+</sup>	07	11Da08 T	1985	$\beta^-$ =100; $\beta^-$ n=33.2
<sup>64</sup> Mn <sup>m</sup>	-42815	4	174.1 0.5	4 <sup>+</sup>	07	10Da06 E	1998	IT=100
<sup>64</sup> Fe	-54970	5	2.0 s 0.2	0 <sup>+</sup>	07		1980	$\beta^-$ =100
<sup>64</sup> Co	-59792	20	300 ms 30	1 <sup>+</sup>	07		1969	$\beta^-$ =100
<sup>64</sup> Co <sup>m</sup>	-59686	4	107 20	5 <sup>+</sup> #		08Bi05 E	2008	$\beta^-$ ?; IT ?
<sup>64</sup> Ni	-67098.9	0.5	STABLE	0 <sup>+</sup>	07		1935	IS=0.9255 19
<sup>64</sup> Cu	-65424.5	0.4	12.7004 h 0.0020	1 <sup>+</sup>	07	12Be04 TD	1936	$\beta^+$ =61.52 26; $\beta^+$ =38.48 26
<sup>64</sup> Cu <sup>i</sup>	-58599	6	6826 6	0 <sup>+</sup> frg.T=4	07	71Be29 E		
<sup>64</sup> Zn	-66004.0	0.6	STABLE (>8.9Ey)	0 <sup>+</sup>	07		1922	IS=49.17 75; 2 $\beta^+$ ?
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>64</sup> Ga	-58832.8	1.4	2.627 m	0.012	0(+ <sup>#</sup> )	07	1953	$\beta^+=100$
<sup>64</sup> Ga <sup>m</sup>	-58790.0	1.4 42.85 0.08	21.9 $\mu$ s	0.7	(2 <sup>+</sup> )	07	1999	IT=100
<sup>64</sup> Ga <sup>i</sup>	-56925.8	2.5 1907.0 2.2	RQ		(0 <sup>+</sup> )T=2	07		
<sup>64</sup> Ge	-54315	4	63.7 s	2.5	0 <sup>+</sup>	07	1972	$\beta^+=100$
<sup>64</sup> As	-39530#	200#	40 ms	30	0 <sup>+</sup> #	07	1995	$\beta^+=100; \beta^+p?$
<sup>64</sup> Se	-26700#	500#	30# ms	(>180 ns)	0 <sup>+</sup>	07	2005	$\beta^+?; \beta^+p?$
* <sup>64</sup> Mn	T: average 11Da08=90(9) 02So.A=91(4) 99So20=85(5) 99Ha05=89(4)							
* <sup>64</sup> Mn	J: 15He28=1 <sup>+</sup>							
* <sup>64</sup> Mn <sup>m</sup>	T: average 11Li50=400(40) 05Ga.B=500(50)							
* <sup>64</sup> Cu	J: directly measured in 10Vi07							
* <sup>64</sup> Cu <sup>i</sup>	E: strongest fragment (xs=100); other 16 keV lower (xs=37)							
* <sup>64</sup> As	T: symmetrized from 18(+43-7)							
<sup>65</sup> V	-11780#	500#	10# ms	(>620 ns)	5/2 <sup>-</sup> #	10 09Ta24	I 2009	$\beta^-?; \beta^-n=40\#; \beta^-2n=1\#$
<sup>65</sup> Cr	-28220#	300#	27.5 ms	2.1	1/2 <sup>-</sup> #	10 11Da08	T 1997	$\beta^-=100; \beta^-n=5\#; \beta^-2n=0.3\#$
<sup>65</sup> Mn	-40967	4	91.9 ms	0.7	(5/2 <sup>-</sup> )	10 13O106	TJ 1985	$\beta^-=100; \beta^-n=7\#$
<sup>65</sup> Fe	-51218	5	810 ms	50	(1/2 <sup>-</sup> )	10 13O106	D 1980	$\beta^-=100; \beta^-n=7.9$ 12
<sup>65</sup> Fe <sup>m</sup>	-50824	5 393.7 0.2	1.12 s	0.15	(9/2 <sup>+</sup> )	10 13O106	E 2008	$\beta^-?$
<sup>65</sup> Fe <sup>n</sup>	-50820	5 397.6 0.2	420 ns	13	(5/2 <sup>+</sup> )	10 13O106	EJ 1998	IT=100
<sup>65</sup> Co	-59185.2	2.1	1.16 s	0.03	(7/2 <sup>-</sup> )	10	1978	$\beta^-=100$
<sup>65</sup> Ni	-65125.7	0.5	2.5175 h	0.0005	5/2 <sup>-</sup>	10	1946	$\beta^-=100$
<sup>65</sup> Ni <sup>m</sup>	-65062.3	0.5 63.37 0.05	69 $\mu$ s	3	1/2 <sup>-</sup>	10	1978	IT=100
<sup>65</sup> Cu	-67263.7	0.6	STABLE		3/2 <sup>-</sup>	10 10Vi07	J 1923	IS=30.85 15
<sup>65</sup> Zn	-65912.0	0.6	243.93 d	0.09	5/2 <sup>-</sup>	10	1939	$\beta^+=100$
<sup>65</sup> Zn <sup>m</sup>	-65858.1	0.6 53.928 0.010	1.6 $\mu$ s	0.6	1/2 <sup>-</sup>	10 FGK149	J	IT=100
<sup>65</sup> Ga	-62657.5	0.8	15.2 m	0.2	3/2 <sup>-</sup>	10	1938	$\beta^+=100$
<sup>65</sup> Ge	-56478.2	2.2	30.9 s	0.5	3/2 <sup>-</sup>	10	1972	$\beta^+=100; \beta^+p=0.011$ 3
<sup>65</sup> As	-46940	80	170 ms	30	3/2 <sup>-</sup> #	10 02Lo13	T 1991	$\beta^+=100; \beta^+p?$
<sup>65</sup> As <sup>i</sup>	-43451	17 3490 90	p		(3/2 <sup>-</sup> )T=3/2	10 11Ro47	J 1993	p=100
<sup>65</sup> Se	-33020#	300#	33 ms	4	3/2 <sup>-</sup> #	10 11Ro47	T 1993	$\beta^+=100; \beta^+p=?$
* <sup>65</sup> Cr	T: average 11Da08=28(3) 03So21=27(3)							
* <sup>65</sup> Mn	T: average 13O106=91.9(0.9) 03So21=92(1)							
* <sup>65</sup> Mn	T: other recent 11Da08=84(8), outweighed, not used							
* <sup>65</sup> Mn	D: $\beta^-n$ has been observed by 99Ha05 but not quantified							
* <sup>65</sup> Fe	J: 09Pa16=(1/2 <sup>-</sup> )							
* <sup>65</sup> Fe <sup>n</sup>	E: also 10Da06=396.8 uncertainty not given T: 10Da06=420(13)							
* <sup>65</sup> Cu	J: directly measured in 10Vi07							
* <sup>65</sup> Zn <sup>m</sup>	J: E2 to ground-state (5/2 <sup>-</sup> ) and M1 from 3/2 <sup>-</sup>							
* <sup>65</sup> As	T: average 02Lo13=126(16) 95Mo26=190(11) with Birge ratio B=3.3							
* <sup>65</sup> As <sup>i</sup>	J: IAS studied in 93Ba12 and 11Ro47							
<sup>66</sup> V	-5610#	500#	5# ms	(>620 ns)		10 09Ta24	I 2009	$\beta^-?; \beta^-n=20\#; \beta^-2n=40\#$
<sup>66</sup> Cr	-24720#	400#	23.8 ms	1.8	0 <sup>+</sup>	15 11Li50	T 1997	$\beta^-=100; \beta^-n=7\#; \beta^-2n=0\#$
<sup>66</sup> Mn	-36750	11	64.2 ms	0.8	(1 <sup>+</sup> )	10 11Pa.A	TD 1992	$\beta^-=100; \beta^-n=8.4$ 9; $\beta^-2n=0.2\#$
<sup>66</sup> Mn <sup>m</sup>	-36286	11 464.5 0.4	780 $\mu$ s	40	(5 <sup>-</sup> )	11Li50	ETJ 2005	IT $\approx$ 100; $\beta^-?$
<sup>66</sup> Fe	-50068	4	351 ms	6	0 <sup>+</sup>	10 12Li02	T 1985	$\beta^-=100; \beta^-n=0\#$
<sup>66</sup> Co	-56409	14	194 ms	17	(1 <sup>+</sup> )	10 12Li02	J 1985	$\beta^-=100; \beta^-n=0\#$
<sup>66</sup> Co <sup>m</sup>	-56234	14 175.1 0.3	1.21 $\mu$ s	0.01	(3 <sup>+</sup> )	10 12Li02	EJ 1998	IT=100
<sup>66</sup> Co <sup>n</sup>	-55767	15 642 5	> 100 $\mu$ s		(8 <sup>-</sup> )	10 98Gr14	E 1998	IT=100
<sup>66</sup> Ni	-66006.3	1.4	54.6 h	0.3	0 <sup>+</sup>	10	1948	$\beta^-=100$
<sup>66</sup> Cu	-66258.3	0.7	5.120 m	0.014	1 <sup>+</sup>	10 10Vi07	J 1937	$\beta^-=100$
<sup>66</sup> Cu <sup>m</sup>	-65104.1	1.6 1154.2 1.4	600 ns	17	(6 <sup>-</sup> )	10 11Lo01	T 1972	IT=100
<sup>66</sup> Zn	-68899.2	0.7	STABLE		0 <sup>+</sup>	10	1922	IS=27.73 98
<sup>66</sup> Ga	-63723.7	1.1	9.304 h	0.008	0 <sup>+</sup>	10 10Se16	T 1937	$\beta^+=100$
<sup>66</sup> Ga <sup>i</sup>	-59874	6 3850 6	RQ		0 <sup>+</sup> T=3			
<sup>66</sup> Ge	-61607.0	2.4	2.26 h	0.05	0 <sup>+</sup>	10	1950	$\beta^+=100$
<sup>66</sup> As	-52025	6	95.77 ms	0.23	0 <sup>+</sup> T=1	10 MMC156	J 1978	$\beta^+=100$
<sup>66</sup> As <sup>m</sup>	-50668	6 1356.63 0.17	1.14 $\mu$ s	0.04	5 <sup>+</sup>	10 13Ru10	TJ 1995	IT=100
<sup>66</sup> As <sup>n</sup>	-49001	6 3023.8 0.3	7.98 $\mu$ s	0.26	9 <sup>+</sup>	10 13Ru10	TJ 1998	IT=100
<sup>66</sup> Se	-41660#	200#	33 ms	12	0 <sup>+</sup>	10 02Lo13	TD 1993	$\beta^+=100; \beta^+p?$
* <sup>66</sup> Cr	T: average 11Li50=24(2) 11Da08=23(4); other 05Ga01=10(6) outweighed							
* <sup>66</sup> Mn	J: 11Li50=(1 <sup>+</sup> ) due to large ground-state feeding from <sup>66</sup> Cr							
* <sup>66</sup> Mn <sup>m</sup>	E: other 05Ga.B=294 + 170 keV T: other 05Ga.B=750(250)							
* <sup>66</sup> Cu	J: directly measured in 10Vi07							
* <sup>66</sup> Cu <sup>m</sup>	T: average 11Lo01=601(30) 72Bi16=600(20)							
* <sup>66</sup> Ga	T: other 12Gy01=9.312(0.032) not used; ENSDF=9.49(0.03)							
* <sup>66</sup> As	J: 0 <sup>+</sup> since super-allowed $\beta^-$ decay; see also 98Gr12							
* <sup>66</sup> As <sup>m</sup>	T: average 13Ru10=1.15(0.04) 01Gr07=1.1(0.1)							
* <sup>66</sup> As <sup>n</sup>	T: average 13Ru10=7.9(0.3) 01Gr07=8.2(0.5)							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{67}\text{V}$	-650#	600#	2# ms (>620 ns)	5/2 <sup>-</sup> #	13		2013	$\beta^-$ ?; $\beta^-_{n=60\#}$ ; $\beta^-_{2n=3\#}$
$^{67}\text{Cr}$	-18680#	400#	10# ms (>300 ns)	1/2 <sup>-</sup> #	05	97Be70 I	1997	$\beta^-$ ?; $\beta^-_{n=10\#}$ ; $\beta^-_{2n=1\#}$
$^{67}\text{Mn}$	-33460#	300#	46.7 ms 2.3	5/2 <sup>-</sup> #	05	11Da08 TD	1997	$\beta^-_{n=100}$ ; $\beta^-_{n=10\ 5}$ ; $\beta^-_{2n=0.01\#}$ *
$^{67}\text{Fe}$	-45610	270	394 ms 9	(1/2 <sup>-</sup> )	05	02So.A TD	1985	$\beta^-_{n=100}$ ; $\beta^-_{n=1\#}$ *
$^{67}\text{Fe}^m$	-45210	270	64 $\mu\text{s}$ 17	(5/2 <sup>+</sup> , 7/2 <sup>+</sup> )	05	11Da08 EJ	1998	IT=100 *
$^{67}\text{Fe}^n$	-45160#	290#	450# 100#	75 $\mu\text{s}$ 21	(9/2 <sup>+</sup> )	08B105 TJ	2008	IT=100
$^{67}\text{Co}$	-55322	6	329 ms 28	(7/2 <sup>-</sup> )	05	08Pa33 TJ	1985	$\beta^-_{n=100}$ ; $\beta^-_{n=0.04\#}$
$^{67}\text{Co}^m$	-54830	6	491.6 1.0	496 ms 33	(1/2 <sup>-</sup> )	09Pa16 E	2008	IT>80; $\beta^-$ ? *
$^{67}\text{Ni}$	-63742.7	2.9	21 s 1	1/2 <sup>-</sup>	05	00Ri14 J	1978	$\beta^-_{n=100}$
$^{67}\text{Ni}^m$	-62736.1	2.9	1006.6 0.2	13.34 $\mu\text{s}$ 0.19	9/2 <sup>+</sup>	05 14Di08 ETJ	1998	IT=100 *
$^{67}\text{Cu}$	-67319.5	0.9	61.83 h 0.12	3/2 <sup>-</sup>	05		1948	$\beta^-_{n=100}$
$^{67}\text{Zn}$	-67880.3	0.8	STABLE	5/2 <sup>-</sup>	05		1928	IS=4.04 16
$^{67}\text{Zn}^m$	-67787.0	0.8	93.312 0.005	9.19 $\mu\text{s}$ 0.06	1/2 <sup>-</sup>	05 15Ch57 T	1972	IT=100 *
$^{67}\text{Zn}^n$	-67275.8	0.8	604.48 0.05	333 ns 14	9/2 <sup>+</sup>	05	1973	IT=100
$^{67}\text{Ga}$	-66879.0	1.2	3.2617 d 0.0005	3/2 <sup>-</sup>	05		1938	$\epsilon=100$
$^{67}\text{Ge}$	-62658	5	18.9 m 0.3	1/2 <sup>-</sup>	05		1950	$\beta^+_{n=100}$
$^{67}\text{Ge}^m$	-62640	5	18.20 0.05	13.7 $\mu\text{s}$ 0.9	5/2 <sup>-</sup>	05	1978	IT=100
$^{67}\text{Ge}^n$	-61906	5	751.70 0.06	109.1 ns 3.8	9/2 <sup>+</sup>	05 00Ch07 T	1973	IT=100 *
$^{67}\text{As}$	-56587.2	0.4	42.5 s 1.2	(5/2 <sup>-</sup> )	05		1980	$\beta^+_{n=100}$
$^{67}\text{Se}$	-46580	70	133 ms 11	5/2 <sup>-</sup> #	05	95B123 T	1991	$\beta^+_{n=100}$ ; $\beta^+_{p=0.5\ 1}$ *
$^{67}\text{Br}$	-32790#	400#		1/2 <sup>-</sup> #				p ?
* $^{67}\text{Mn}$	T : average 11Da08=51(4) 03So21=47(4) 99Ha05=42(4) **							
* $^{67}\text{Fe}$	T : others recent 11Da08=304(81) 08Pa33=416(29), outweighed, not used **							
* $^{67}\text{Fe}^m$	T : average 03Sa02=75(21) 98Gr14=43(30), same authors, different experiment **							
* $^{67}\text{Fe}^n$	E : less than 30 keV above 387.7 level **							
* $^{67}\text{Co}^m$	E : 09Pa16=491.55(0.11) $\gamma$ ray; 08Pa33=491.6(1.0) D : from 08Pa33 **							
* $^{67}\text{Ni}^m$	T : average 14Di08=13.7(0.6) 98Gr14=13.3(0.2); other 02Ge16=13(1) **							
* $^{67}\text{Zn}^m$	T : unweighed average 15Ch57=9.37(0.04) 98At04=9.34(0.20) 96Hw03=9.01(0.03) **							
* $^{67}\text{Zn}^n$	T : 75Ro25=9.1(0.4) 73Le18=9.20(0.07) 72Le37=9.15(0.05) **							
* $^{67}\text{Ge}^n$	T : average 00Ch07=101(3) 79A104=110.9(1.4); Birge ratio B=2.99 **							
* $^{67}\text{Se}$	T : average 02Lo13=136(12) 94Ba50=107(35) **							
* $^{67}\text{Se}$	T : values from 95B123 for $^{67}\text{Se}=60(+17-11)$ and $^{71}\text{Kr}$ questioned in 97Oi01 **							
$^{68}\text{Cr}$	-14800#	500#	5# ms (>620 ns)	0 <sup>+</sup>	12	09Ta24 I	2009	$\beta^-$ ?; $\beta^-_{n=10\#}$ ; $\beta^-_{2n=0.1\#}$
$^{68}\text{Mn}$	-28380#	400#	33.7 ms 1.5	> 3	12	15Be32 T	1995	$\beta^-_{n=100}$ ; $\beta^-_{n=10\#}$ ; $\beta^-_{2n=2\#}$ *
$^{68}\text{Fe}$	-43490	370	188 ms 4	0 <sup>+</sup>	12		1985	$\beta^-_{n=100}$ ; $\beta^-_{n>0}$
$^{68}\text{Co}$	-51930	190	200 ms 20	(7 <sup>-</sup> )	12		1985	$\beta^-_{n=100}$ ; $\beta^-_{n=1\#}$
$^{68}\text{Co}^m$	-51780#	240#	150# 150#	1.6 s 0.3	(1 <sup>+</sup> )	12	1998	$\beta^-_{n=100}$
$^{68}\text{Co}^n$	-51740#	240#	195# 150#	101 ns 10	(0, 1)	12 10Da06 T	2010	IT=100 *
$^{68}\text{Ni}$	-63463.8	3.0	29 s 2	0 <sup>+</sup>	12		1977	$\beta^-_{n=100}$
$^{68}\text{Ni}^m$	-61860	3	1603.52 0.27	270 ns 5	0 <sup>+</sup>	12 15F101 E		4.IT=100 *
$^{68}\text{Ni}^n$	-60615	3	2849.1 0.3	850 $\mu\text{s}$ 30	5 <sup>-</sup>	12 15Wi02 T	1995	IT=100 *
$^{68}\text{Cu}$	-65567.0	1.6	30.9 s 0.6	30.9 s 0.6	1 <sup>+</sup>	12 10Vi07 J	1953	$\beta^-_{n=100}$
$^{68}\text{Cu}^m$	-64845.7	1.6	721.26 0.08	3.75 m 0.05	6 <sup>-</sup>	12 10Vi07 J	1969	IT=86 2; $\beta^-_{n=14\ 2}$ *
$^{68}\text{Zn}$	-70007.1	0.8	STABLE	0 <sup>+</sup>	12		1922	IS=18.45 63
$^{68}\text{Ga}$	-67086.0	1.4	67.845 m 0.018	67.845 m 0.018	1 <sup>+</sup>	12 14Ga09 T	1937	$\beta^+_{n=100}$ *
$^{68}\text{Ge}$	-66978.8	1.9	270.93 d 0.13	270.93 d 0.13	0 <sup>+</sup>	12	1948	$\epsilon=100$
$^{68}\text{As}$	-58894.5	1.8	151.6 s 0.8	151.6 s 0.8	3 <sup>+</sup>	12	1971	$\beta^+_{n=100}$
$^{68}\text{As}^m$	-58469.4	1.8	425.1 0.2	111 ns 20	1 <sup>+</sup>	12	1994	IT=100 *
$^{68}\text{Se}$	-54189.4	0.5	35.5 s 0.7	35.5 s 0.7	0 <sup>+</sup>	12	1990	$\beta^+_{n=100}$
$^{68}\text{Br}$	-38790#	260#		<1.5 $\mu\text{s}$	3 <sup>+</sup> #	12 95B106 I		p ?
* $^{68}\text{Mn}$	T : average 15Be32=38.3(3.6) 35.2(2.0) 11Da08=29(4) 03So21=28(8) 99Ha05=28(4) **							
* $^{68}\text{Mn}$	D : $\beta^-_{n}$ has been observed by 99Ha05 but not quantified **							
* $^{68}\text{Co}^n$	J : 12Li02 strong feeding in $\beta^-$ of $^{68}\text{Fe}$ (0 <sup>+</sup> ) **							
* $^{68}\text{Ni}^m$	E : average 15F101=1603.6(0.6) 13Re18=1603.5(0.3) **							
* $^{68}\text{Ni}^n$	T : average 15Wi02=840(40) 95Br10=860(50) **							
* $^{68}\text{Cu}$	J : directly measured in 10Vi07 **							
* $^{68}\text{Cu}^m$	J : directly measured in 10Vi07 **							
* $^{68}\text{Ga}$	T : also 12Lu14=67.87(0.10); discrepant 83Iw02=67.629(0.24) **							
* $^{68}\text{As}^m$	T : symmetrized from 94Ba50=107(+23-16) **							



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>69</sup> Cr	-8580# 500#		2# ms (>620 ns)	7/2 <sup>+</sup> #	14		2013	$\beta^-$ ?; $\beta^-n=20\#$ ; $\beta^-2n=6\#$
<sup>69</sup> Mn	-24770# 400#		22.1 ms 1.6	5/2 <sup>-</sup> #	14	15Be32 TD	1995	$\beta^-n=100$ ; $\beta^-n=50$ 20; $\beta^-2n=0.4\#$ *
<sup>69</sup> Fe	-39030# 400#		108.2 ms 4.5	1/2 <sup>-</sup> #	14	13Ma87 T	1992	$\beta^-n=100$ ; $\beta^-n=7\#$ ; $\beta^-2n=0\#$ *
<sup>69</sup> Co	-50280 140		180 ms 20	7/2 <sup>-</sup> #	14	15Li33 T	1985	$\beta^-n=100$ ; $\beta^-n=1\#$
<sup>69</sup> Co <sup>m</sup>	-49780# 240#	500# 200#	750 ms 250	1/2 <sup>-</sup> #		15Li33 TD	2015	$\beta^-n=100$
<sup>69</sup> Ni	-59979 4		11.4 s 0.3	(9/2 <sup>+</sup> )	14		1984	$\beta^-n=100$
<sup>69</sup> Ni <sup>m</sup>	-59658 4	321 2	3.5 s 0.4	(1/2 <sup>-</sup> )	14		1998	$\beta^-n\approx 100$ ; IT<0.01 *
<sup>69</sup> Ni <sup>n</sup>	-57279 4	2700.0 1.0	439 ns 3	(17/2 <sup>-</sup> )	14		1998	IT=100
<sup>69</sup> Cu	-65736.2 1.4		2.85 m 0.15	3/2 <sup>-</sup>	14	10Vi07 J	1966	$\beta^-n=100$ *
<sup>69</sup> Cu <sup>m</sup>	-62994.2 1.6	2742.0 0.7	357 ns 2	(13/2 <sup>+</sup> )	14		1997	IT=100
<sup>69</sup> Zn	-68417.8 0.8		56.4 m 0.9	1/2 <sup>-</sup>	14		1937	$\beta^-n=100$
<sup>69</sup> Zn <sup>m</sup>	-67979.2 0.8	438.636 0.018	13.756 h 0.018	9/2 <sup>+</sup>	14		1970	IT $\approx$ 100; $\beta^-n=0.033$ 3
<sup>69</sup> Ga	-69327.8 1.2		STABLE	3/2 <sup>-</sup>	14		1923	IS=60.108 9
<sup>69</sup> Ge	-67100.7 1.3		39.05 h 0.10	5/2 <sup>-</sup>	14		1938	$\beta^+n=100$
<sup>69</sup> Ge <sup>m</sup>	-67013.9 1.3	86.76 0.02	5.1 $\mu$ s 0.2	1/2 <sup>-</sup>	14		1978	IT=100
<sup>69</sup> Ge <sup>n</sup>	-66702.8 1.3	397.94 0.02	2.81 $\mu$ s 0.05	9/2 <sup>+</sup>	14		1978	IT=100
<sup>69</sup> As	-63110 30		15.2 m 0.2	5/2 <sup>-</sup>	14		1955	$\beta^+n=100$
<sup>69</sup> Se	-56434.7 1.5		27.4 s 0.2	1/2 <sup>-</sup>	14		1974	$\beta^+n=100$ ; $\beta^+p=0.045$ 10
<sup>69</sup> Se <sup>m</sup>	-56395.9 1.5	38.85 0.22	2.0 $\mu$ s 0.2	5/2 <sup>-</sup>	14		1988	IT=100
<sup>69</sup> Se <sup>n</sup>	-55860.7 1.6	574.0 0.4	955 ns 16	9/2 <sup>+</sup>	14	00Ch07 T	1988	IT=100 *
<sup>69</sup> Br	-46260 40		* < 24 ns	(5/2 <sup>-</sup> )	15		1988	p=100
<sup>69</sup> Br <sup>m</sup>	-46220# 110#	40# 100#	*	5/2 <sup>-</sup> #		Mirror I		
<sup>69</sup> Br <sup>n</sup>	-45690# 110#	570# 100#		9/2 <sup>+</sup> #		Mirror I		
<sup>69</sup> Br <sup>i</sup>	-42771 19	3490 50	p	(5/2 <sup>-</sup> )T=3/2	14	11Ro47 I	2011	p=100
<sup>69</sup> Kr	-32440# 400#		28 ms 1	(5/2 <sup>-</sup> )	15	14De41 D	1995	$\beta^+n=100$ ; $\beta^+p=55$ 7 *
* <sup>69</sup> Mn	T: average 15Be32=24.1(2.6) 25.8(2.8) 11Da08=18(4) 99Ha05=14(4) **							
* <sup>69</sup> Fe	T: average 13Ma87=102(10) 11Da08=110(6) 03So21=109(9) **							
* <sup>69</sup> Ni <sup>m</sup>	E: 9/2 <sup>+</sup> level in isotones: <sup>73</sup> Ge=-66 <sup>71</sup> Zn=157(1) <sup>69</sup> Ni=-321(2) exhibits **							
* <sup>69</sup> Ni <sup>m</sup>	E: unusually strong variations **							
* <sup>69</sup> Cu	J: directly measured in 10Vi07 **							
* <sup>69</sup> Se <sup>n</sup>	T: average 00Ch07=950(21) 95Po01=960(23) **							
* <sup>69</sup> Kr	T: 14De41=28(1) 11Ro47=27(3) 97Xu01=32(10) D: $\beta^+p=52.5(6.5) + 2.4(0.5)$ **							
<sup>70</sup> Cr	-4480# 600#		1# ms (>620 ns)	0 <sup>+</sup>	13		2013	$\beta^-$ ?; $\beta^-n=40\#$ ; $\beta^-2n=2\#$
<sup>70</sup> Mn	-19500# 500#		19.9 ms 1.7		09	15Be32 TD	2009	$\beta^-n=100$ ; $\beta^-n=20\#$ ; $\beta^-2n=7\#$
<sup>70</sup> Fe	-36510# 400#		63.0 ms 4.5	0 <sup>+</sup>	04	13Ma87 T	1997	$\beta^-n=100$ ; $\beta^-n=10\#$ *
<sup>70</sup> Co	-46630# 300#		* 112 ms 7	(6 <sup>-</sup> , 7 <sup>-</sup> )	16		1985	$\beta^-n=100$ ; $\beta^-n=3\#$ ; $\beta^-2n=0\#$
<sup>70</sup> Co <sup>m</sup>	-46430# 360#	200# 200#	* 470 ms 50	3 <sup>+</sup> #	16		1998	$\beta^-n\approx 100$ ; IT ?; $\beta^-n=3\#$ ; $\beta^-2n=0\#$
<sup>70</sup> Ni	-59213.9 2.1		6.0 s 0.3	0 <sup>+</sup>	04		1987	$\beta^-n=100$
<sup>70</sup> Ni <sup>m</sup>	-56353.9 2.9	2860 2	232 ns 1	(8 <sup>+</sup> )	04		1997	IT=100
<sup>70</sup> Cu	-62976.4 1.1		44.5 s 0.2	6 <sup>-</sup>	04	10Vi07 J	1971	$\beta^-n=100$
<sup>70</sup> Cu <sup>m</sup>	-62875.3 1.1	101.1 0.3	33 s 2	3 <sup>-</sup>	04	10Vi07 J	2002	$\beta^-n=52$ 9; IT=48 9 *
<sup>70</sup> Cu <sup>n</sup>	-62733.8 1.2	242.6 0.5	6.6 s 0.2	1 <sup>+</sup>	04	10Vi07 J	1971	$\beta^-n=93.2$ 9; IT=6.8 9 *
<sup>70</sup> Zn	-69564.7 1.9		STABLE	0 <sup>+</sup>	04		1922	IS=0.61 10; 2 $\beta^-$ ? *
<sup>70</sup> Ga	-68910.1 1.2		21.14 m 0.03	1 <sup>+</sup>	04		1937	$\beta^-n\approx 100$ ; $\epsilon=0.41$ 6
<sup>70</sup> Ge	-70561.9 0.8		STABLE	0 <sup>+</sup>	04		1923	IS=20.57 27
<sup>70</sup> As	-64340 50		52.6 m 0.3	4 <sup>+</sup>	04		1950	$\beta^+n=100$
<sup>70</sup> As <sup>m</sup>	-64310 50	32.008 0.002	96 $\mu$ s 3	2 <sup>+</sup>	04		1979	IT=100
<sup>70</sup> Se	-61929.9 1.6		41.1 m 0.3	0 <sup>+</sup>	04		1950	$\beta^+n=100$
<sup>70</sup> Br	-51426 15		79.1 ms 0.8	0 <sup>+</sup> T=1	04		1978	$\beta^+n=100$ ; $\beta^+p$ ?
<sup>70</sup> Br <sup>m</sup>	-49134 15	2292.3 0.8	2.2 s 0.2	9 <sup>+</sup>	04		1981	$\beta^+n$ ?; IT ?; $\beta^+p$ ?
<sup>70</sup> Kr	-41100# 200#		52 ms 17	0 <sup>+</sup>	04		1995	$\beta^+n=100$ ; $\beta^+p<1.3$
* <sup>70</sup> Fe	T: average 13Ma87=61(5) 11Da08=71(10); other 03So21=94(17) not used **							
* <sup>70</sup> Cu	J: directly measured in 10Vi07 **							
* <sup>70</sup> Cu <sup>m</sup>	J: directly measured in 10Vi07 **							
* <sup>70</sup> Cu <sup>n</sup>	J: directly measured in 10Vi07 **							
* <sup>70</sup> Zn	T: 03Ki08 : 0v- $\beta\beta$ >13 Py **							
<sup>71</sup> Mn	-15570# 500#		5# ms (>400 ns)	5/2 <sup>-</sup> #	10	10Oh02 I	2010	$\beta^-$ ?; $\beta^-n=30\#$ ; $\beta^-2n=3\#$
<sup>71</sup> Fe	-31430# 400#		33.7 ms 3.8	7/2 <sup>+</sup> #	10	13Ma87 T	1997	$\beta^-n=100$ ; $\beta^-n=10\#$ ; $\beta^-2n=0.3\#$ *
<sup>71</sup> Co	-44370 470		80 ms 3	(7/2 <sup>-</sup> )	10	12Ra10 TJD	1992	$\beta^-n=100$ ; $\beta^-n=3$ 1 *
<sup>71</sup> Ni	-55406.2 2.2		2.56 s 0.03	(9/2 <sup>+</sup> )	10		1987	$\beta^-n=100$
<sup>71</sup> Ni <sup>m</sup>	-55406.0 2.3	499 5	2.3 s 0.3	(1/2 <sup>-</sup> )	10		2009	$\beta^-n=100$
<sup>71</sup> Cu	-62711.1 1.5		19.4 s 1.4	3/2 <sup>-</sup>	10	10Vi07 J	1983	$\beta^-n=100$ *
<sup>71</sup> Cu <sup>m</sup>	-59955.4 1.6	2755.7 0.6	271 ns 13	(19/2 <sup>-</sup> )	10	98Gr14 TJ	1998	IT=100 *
<sup>71</sup> Zn	-67328.8 2.7		2.45 m 0.10	1/2 <sup>-</sup>	10		1955	$\beta^-n=100$
<sup>71</sup> Zn <sup>m</sup>	-67171.1 2.4	157.7 1.3	MD 4.125 h 0.007	9/2 <sup>+</sup>	10	12Re05 T	1958	$\beta^-n\approx 100$ ; IT<0.05
<sup>71</sup> Ga	-70139.1 0.8		STABLE	3/2 <sup>-</sup>	10		1923	IS=39.892 9

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>71</sup> Ge	-69906.5	0.8	11.43 d 0.03	1/2 <sup>-</sup>	10		1941	$\epsilon=100$
<sup>71</sup> Ge <sup>m</sup>	-69708.1	0.8	198.354	0.014	20.41 ms 0.18		1959	IT=100
<sup>71</sup> As	-67893	4			65.30 h 0.07		1939	$\beta^+=100$
<sup>71</sup> Se	-63146.5	2.8			4.74 m 0.05		1957	$\beta^+=100$
<sup>71</sup> Se <sup>m</sup>	-63097.7	2.8	48.79	0.05	5.6 $\mu$ s 0.7		1982	IT=100
<sup>71</sup> Se <sup>n</sup>	-62886.0	2.8	260.48	0.10	19.0 $\mu$ s 0.5		1982	IT=100
<sup>71</sup> Br	-56502	5			21.4 s 0.6		1981	$\beta^+=100$
<sup>71</sup> Kr	-46330	130			100 ms 3		1981	$\beta^+=100; \beta^+p=2.17$
<sup>71</sup> Rb	-32060#	400#			*			p ?
<sup>71</sup> Rb <sup>m</sup>	-32010#	410#	50#	100#	*			
<sup>71</sup> Rb <sup>n</sup>	-31800#	410#	260#	100#				
* <sup>71</sup> Fe	T : average 13Ma87=42(6) 11Da08=28(5)							
* <sup>71</sup> Co	D : taking into account 12Ra10<2.7(0.9)% and 05Ma95>3(1)% of same group							
* <sup>71</sup> Cu	T : average 99Pr10=19(3) 83Ru06=19.5(1.6)							
* <sup>71</sup> Cu	J : directly measured in 10Vi07							
* <sup>71</sup> Cu <sup>m</sup>	T : average 98Is11=250(30) 98Gr14=275(14)							
<sup>72</sup> Mn	-9900#	600#			2# ms (>620 ns)		13	2013
<sup>72</sup> Fe	-28430#	500#			19 ms 4		0 <sup>+</sup>	10 13Ma87 TD 1997
<sup>72</sup> Co	-40200#	400#			52.5 ms 0.3		(6 <sup>-</sup> , 7 <sup>-</sup> )	10 16Mo07 T 1992
<sup>72</sup> Co <sup>m</sup>	-40000#	450#	200#	200#	*		(0 <sup>+</sup> , 1 <sup>+</sup> )	10 16Mo07 TJ 2016
<sup>72</sup> Ni	-54226.1	2.2			1.57 s 0.05		0 <sup>+</sup>	10 1987
<sup>72</sup> Cu	-59783.0	1.4			6.63 s 0.03		2 <sup>-</sup>	10 10Vi07 J 1983
<sup>72</sup> Cu <sup>m</sup>	-59512.7	1.7	270.3	1.0	1.76 $\mu$ s 0.03		(6 <sup>-</sup> )	10 1998
<sup>72</sup> Zn	-68145.5	2.1			46.5 h 0.1		0 <sup>+</sup>	10 1951
<sup>72</sup> Ga	-68588.3	0.8			14.025 h 0.010		3 <sup>-</sup>	10 12Kr07 T 1939
<sup>72</sup> Ga <sup>m</sup>	-68468.6	0.8	119.66	0.05	39.68 ms 0.13		(0 <sup>+</sup> )T=1	10 1968
<sup>72</sup> Ge	-72585.90	0.08			STABLE		0 <sup>+</sup>	10 1923
<sup>72</sup> Ge <sup>m</sup>	-71894.47	0.09	691.43	0.04	444.2 ns 0.8		0 <sup>+</sup>	10 1984
<sup>72</sup> As	-68230	4			26.0 h 0.1		2 <sup>-</sup>	10 1939
<sup>72</sup> Se	-67868.2	2.0			8.40 d 0.08		0 <sup>+</sup>	10 1948
<sup>72</sup> Br	-59061.7	1.0			78.6 s 2.4		1 <sup>+</sup>	10 1970
<sup>72</sup> Br <sup>m</sup>	-58960.9	1.0	100.76	0.15	10.6 s 0.3		(3 <sup>-</sup> )	10 1980
<sup>72</sup> Kr	-53941	8			17.16 s 0.18		0 <sup>+</sup>	10 03Pi03 T 1973
<sup>72</sup> Rb	-38330#	500#			<1.5 $\mu$ s		1 <sup>+</sup> #	10 95Bi06 I
<sup>72</sup> Rb <sup>m</sup>	-38230#	510#	100#	100#	*		3 <sup>-</sup> #	
* <sup>72</sup> Co	T : others 14Xu07=52.8(1.6) 14Ra20=55(4) 05Ma59=59(2) 03Sa40=62(3)							
* <sup>72</sup> Co	J : feeding of the 6 <sup>+</sup> level in <sup>72</sup> Ni and shell model D : 05Ma95 $\beta^-n>62$							
* <sup>72</sup> Cu	J : directly measured in 10Vi07							
* <sup>72</sup> Cu <sup>m</sup>	D : no $\beta^-$ decay observed in 05Th.A							
* <sup>72</sup> Kr	T : average 03Pi03=17.1(0.2) 73Da22=17.4(0.4)							
<sup>73</sup> Fe	-22900#	500#			12.9 ms 1.6		7/2 <sup>+</sup> #	16 2010
<sup>73</sup> Co	-37420#	400#			40.7 ms 1.3		7/2 <sup>-</sup> #	16 12Ra10 D 1995
<sup>73</sup> Ni	-50108.2	2.4			840 ms 30		(9/2 <sup>+</sup> )	16 1987
<sup>73</sup> Cu	-58987.4	1.9			4.2 s 0.3		3/2 <sup>-</sup>	04 10Vi07 J 1983
<sup>73</sup> Zn	-65593.4	1.9			23.5 s 1.0		(1/2 <sup>-</sup> )	04 1972
<sup>73</sup> Zn <sup>m</sup>	-65397.9	1.9	195.5	0.2	13.0 ms 0.2		(5/2 <sup>+</sup> )	04 1985
<sup>73</sup> Zn <sup>n</sup>	-65355.8	2.8	237.6	2.0	EU		(9/2 <sup>+</sup> )	04 1998
<sup>73</sup> Ga	-69699.3	1.7			4.86 h 0.03		1/2 <sup>-</sup>	04 11Ch16 J 1949
<sup>73</sup> Ge	-71297.52	0.06			STABLE		9/2 <sup>+</sup>	04 1933
<sup>73</sup> Ge <sup>m</sup>	-71284.24	0.06	13.2845	0.0015	2.92 $\mu$ s 0.03		5/2 <sup>+</sup>	04 1975
<sup>73</sup> Ge <sup>n</sup>	-71230.79	0.06	66.726	0.009	499 ms 11		1/2 <sup>-</sup>	04 1957
<sup>73</sup> As	-70953	4			80.30 d 0.06		3/2 <sup>-</sup>	04 1948
<sup>73</sup> As <sup>m</sup>	-70525	4	427.906	0.021	5.7 $\mu$ s 0.2		9/2 <sup>+</sup>	04 1956
<sup>73</sup> Se	-68227	7			7.15 h 0.08		9/2 <sup>+</sup>	04 1948
<sup>73</sup> Se <sup>m</sup>	-68201	7	25.71	0.04	39.8 m 1.3		3/2 <sup>-</sup>	04 1960
<sup>73</sup> Br	-63647	7			3.4 m 0.2		1/2 <sup>-</sup>	04 1970
<sup>73</sup> Kr	-56552	7			27.3 s 1.0		3/2 <sup>-</sup>	04 1972
<sup>73</sup> Kr <sup>m</sup>	-56118	7	433.66	0.12	107 ns 10		(9/2 <sup>+</sup> )	04 1993
<sup>73</sup> Rb	-46080#	200#			<30 ns		3/2 <sup>-</sup> #	04 96Pf01 I
<sup>73</sup> Rb <sup>m</sup>	-45650#	220#	430#	100#			9/2 <sup>+</sup> #	Mirror I
<sup>73</sup> Rb <sup>i</sup>	-42850	40	3230#	200#	p		1/2 <sup>-</sup> T=3/2	93Ba61 JD 1993
<sup>73</sup> Sr	-31950#	400#			> 25 ms		1/2 <sup>-</sup> #	04 1993
* <sup>73</sup> Co	D : taking into account 12Ra10<22(8)% 10Ho12<7.9% 05Ma95>9(4)%							
* <sup>73</sup> Cu	J : directly measured in 10Vi07							
* <sup>73</sup> Zn <sup>n</sup>	E : if 42.1 keV $\gamma$ feeds <sup>73</sup> Zn <sup>m</sup> , EU: see discussion in ENSDF'04							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{74}\text{Fe}$	-19590#	600#	2# ms (>400 ns)	$0^+$	10	10Oh02 I	2010	$\beta^- ?$ ; $\beta^-_{n=30\#}$ ; $\beta^-_{2n=2\#}$
$^{74}\text{Co}$	-32820#	500#	31.3 ms 1.3		06	05Ma95 TD	1995	$\beta^-_{=100}$ ; $\beta^-_{n>26\ 9}$ ; $\beta^-_{2n=1\#}$ *
$^{74}\text{Ni}$	-48460#	200#	507.7 ms 4.6	$0^+$	06	14Xu07 T	1987	$\beta^-_{=100}$ ; $\beta^-_{n=5\#}$
$^{74}\text{Cu}$	-56006	6	1.63 s 0.05	$2^-$	06	10Vi07 J	1987	$\beta^-_{=100}$ ; $\beta^-_{n=40\#}$ *
$^{74}\text{Zn}$	-65756.7	2.5	95.6 s 1.2	$0^+$	06		1972	$\beta^-_{=100}$
$^{74}\text{Ga}$	-68049.6	3.0	8.12 m 0.12	$(3^-)$	06	13Ma15 J	1956	$\beta^-_{=100}$
$^{74}\text{Ga}^m$	-67990	3	59.571 0.014	$(0)^{(+\#)}$	06		1974	$\text{IT}=75\ 25$ ; $\beta^- ?$
$^{74}\text{Ge}$	-73422.442	0.013	STABLE	$0^+$	06		1923	$\text{IS}=36.50\ 20$
$^{74}\text{As}$	-70860.1	1.7	17.77 d 0.02	$2^-$	06		1938	$\beta^+_{=66\ 2}$ ; $\beta^-_{=34\ 2}$
$^{74}\text{Se}$	-72213.201	0.015	STABLE (>15 Ey)	$0^+$	06	15Je02 T	1922	$\text{IS}=0.89\ 4$ ; $2\beta^+ ?$
$^{74}\text{Br}$	-65288	6	25.4 m 0.3	$(0^-)$	06		1952	$\beta^+_{=100}$
$^{74}\text{Br}^m$	-65274	6	13.58 0.21	$4^{(+\#)}$	06		1953	$\beta^+_{=100}$
$^{74}\text{Kr}$	-62331.8	2.0	46 m 2	$0^+$	06		1960	$\beta^+_{=100}$
$^{74}\text{Kr}^i$	-61790	30	540 30			98Gr.A E	1998	$\text{IT}=100$ *
$^{74}\text{Rb}$	-51916	3	64.776 ms 0.030	$0^+T=1$	06		1977	$\beta^+_{=100}$ ; $\beta^+_{p ?}$
$^{74}\text{Sr}$	-40830#	100#	27 ms 8	$0^+$	15		1995	$\beta^+_{=100}$ ; $\beta^+_{p ?}$
* $^{74}\text{Co}$	T : average $14\text{Xu}07=31.6(1.5)$ $05\text{Ma}95=30(3)$ **							
* $^{74}\text{Co}$	T : others recent $11\text{Da}08=19(7)$ $10\text{Ho}12=34(+6-9)$ D : $\beta^-_{n=18(15)\%}$ in $10\text{Ho}12$ **							
* $^{74}\text{Cu}$	D : $\beta^-_{n}$ has been observed by $91\text{Kr}15$ but not quantified **							
* $^{74}\text{Kr}^i$	E : $E(g)<85$ to $2^+$ level at $455.61(0.10)$ keV **							
$^{75}\text{Fe}$	-13640#	600#	2# ms (>620 ns)	$9/2^+\#$	13		2013	$\beta^- ?$ ; $\beta^-_{n=80\#}$ ; $\beta^-_{2n=20\#}$
$^{75}\text{Co}$	-29650#	500#	26.5 ms 1.2	$7/2^-\#$	13	14Xu07 T	1995	$\beta^-_{=100}$ ; $\beta^-_{n<16}$ ; $\beta^-_{2n=0.5\#}$
$^{75}\text{Ni}$	-44030#	300#	331.6 ms 3.2	$7/2^+\#$	13	14Xu07 T	1992	$\beta^-_{=100}$ ; $\beta^-_{n=10.0\ 28}$
$^{75}\text{Cu}$	-54471.3	2.3	1.224 s 0.003	$5/2^-$	13	10Vi07 J	1985	$\beta^-_{=100}$ ; $\beta^-_{n=3.5\ 6}$
$^{75}\text{Cu}^m$	-54409.6	2.3	310 ns 8	$(1/2^-, 3/2^-)$	13		2010	$\text{IT}=100$
$^{75}\text{Cu}^n$	-54405.1	2.3	149 ns 5	$(3/2^-, 1/2^-)$	13	16Pe14 EJT	2010	$\text{IT}=100$
$^{75}\text{Zn}$	-62558.9	2.0	10.2 s 0.2	$(7/2^+)$	13		1974	$\beta^-_{=100}$
$^{75}\text{Zn}^m$	-62432.0	2.0	126.94 0.09	$(1/2^-)$	13		2011	$\beta^- ?$ ; $\text{IT} ?$
$^{75}\text{Ga}$	-68464.6	2.4	126 s 2	$3/2^-$	13		1960	$\beta^-_{=100}$
$^{75}\text{Ge}$	-71856.96	0.05	82.78 m 0.04	$1/2^-$	13		1939	$\beta^-_{=100}$
$^{75}\text{Ge}^m$	-71717.27	0.06	139.69 0.03	$7/2^+$	13		1952	$\text{IT}\approx 100$ ; $\beta^-_{=0.030\ 6}$
$^{75}\text{Ge}^n$	-71664.77	0.08	192.19 0.06	$5/2^+$	13		1982	$\text{IT}=100$
$^{75}\text{As}$	-73034.2	0.9	STABLE	$3/2^-$	13		1920	$\text{IS}=100$ .
$^{75}\text{As}^m$	-72730.3	0.9	303.9243 0.0008	$9/2^+$	13		1957	$\text{IT}=100$
$^{75}\text{Se}$	-72169.48	0.07	119.78 d 0.05	$5/2^+$	13		1947	$\epsilon=100$
$^{75}\text{Br}$	-69107	4	96.7 m 1.3	$3/2^-$	13		1948	$\beta^+_{=100}$
$^{75}\text{Kr}$	-64324	8	4.60 m 0.07	$5/2^+$	13		1960	$\beta^+_{=100}$
$^{75}\text{Rb}$	-57218.7	1.2	19.0 s 1.2	$3/2^{(-)}$	13		1975	$\beta^+_{=100}$
$^{75}\text{Sr}$	-46620	220	88 ms 3	$(3/2^-)$	13		1991	$\beta^+_{=100}$ ; $\beta^+_{p}=5.2\ 9$
$^{75}\text{Y}$	-31820#	300#	100# $\mu\text{s}$	$5/2^+\#$				$\beta^+ ?$ ; $\beta^+_{p ?}$ ; $p ?$
$^{76}\text{Co}$	-24510#	600#	& 23 ms 6	$(8^-)$	14	14Xu07 TD	2010	$\beta^-_{=100}$ ; $\beta^-_{n=4\#}$ ; $\beta^-_{2n=6\#}$ *
$^{76}\text{Co}^m$	-24410#	610#	100# 100#	$(1^-)$		15So23 TJD	2015	$\beta^-_{=100}$
$^{76}\text{Co}^n$	-23770#	610#	740# 100#	$(3^+)$		15So23 TJD	2015	$\text{IT}=100$ *
$^{76}\text{Ni}$	-41630#	400#	234.6 ms 2.7	$0^+$	07	14Xu07 T	1995	$\beta^-_{=100}$ ; $\beta^-_{n=14\ 3.6}$
$^{76}\text{Ni}^m$	-39210#	400#	2418.7 1.0	$(8^+)$	07	15So23 T	2005	$\text{IT}=100$ *
$^{76}\text{Cu}$	-50976	7	* 637.7 ms 5.5	$(3, 4)$	95	09Wi03 D	1987	$\beta^-_{=100}$ ; $\beta^-_{n=7.2\ 5}$ *
$^{76}\text{Cu}^m$	-50980#	200#	0# 200#	$(1, 3)$	95	90Wi12 J	1990	$\beta^-_{=100}$
$^{76}\text{Zn}$	-62303.0	1.5	5.7 s 0.3	$0^+$	95		1974	$\beta^-_{=100}$
$^{76}\text{Ga}$	-66296.6	2.0	32.6 s 0.6	$2^-$	95	11Ma45 J	1961	$\beta^-_{=100}$
$^{76}\text{Ge}$	-73212.889	0.018	1.66 Zy 0.13	$0^+$	95	15Ba11 T	1933	$\text{IS}=7.73\ 12$ ; $2\beta^-_{=100}$ *
$^{76}\text{As}$	-72291.4	0.9	1.0778 d 0.0020	$2^-$	95		1934	$\beta^-_{\approx 100}$ ; $\epsilon<0.02$
$^{76}\text{As}^m$	-72247.0	0.9	44.425 0.001	$(1)^+$	95		1966	$\text{IT}=100$
$^{76}\text{Se}$	-75251.950	0.016	STABLE	$0^+$	95		1922	$\text{IS}=9.37\ 29$
$^{76}\text{Br}$	-70289	9	16.2 h 0.2	$1^-$	95		1952	$\beta^+_{=100}$
$^{76}\text{Br}^m$	-70186	9	102.58 0.03	$(4)^+$	95		1979	$\text{IT}>99.4$ ; $\beta^+<0.6$
$^{76}\text{Kr}$	-69014	4	14.8 h 0.1	$0^+$	95		1954	$\beta^+_{=100}$
$^{76}\text{Rb}$	-60479.1	0.9	36.5 s 0.6	$1^{(-)}$	95		1969	$\beta^+_{=100}$ ; $\beta^+_{\alpha}=3.8e-7\ 10$
$^{76}\text{Rb}^m$	-60162.2	0.9	316.93 0.08	$(4^+)$	95	00Ch07 T	1986	$\text{IT}=100$

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
$^{76}\text{Sr}$	-54250	30	7.89 s	0.07	0 <sup>+</sup>	11	1990	$\beta^+=100$ ; $\beta^+p=3.4e-5$ 8
$^{76}\text{Y}$	-38480#	300#	120# $\mu\text{s}$ (>200 ns)		1 <sup>-</sup> #	07	01Ki13	I 2001 $\beta^+ ?$ ; p ?; $\beta^+p ?$
$^{76}\text{Co}$	T: symmetrized from 14Xu07=21.7(+6.5-4.9)		J: 15So23=(8 <sup>-</sup> )					
$^{76}\text{Co}^n$	E: 15So23=638.4(0.8) above (1 <sup>-</sup> ) 76Com							
$^{76}\text{Co}^m$	T: symmetrized from 15So23=2.96(+0.29-0.25)							
$^{76}\text{Ni}^m$	E: 12Ka36=142.7(0.5), 355.5(0.5), 929.9(0.5) and 990.6(0.5) $\gamma$ rays in							
$^{76}\text{Ni}^m$	E: cascade to ground-state =2418.7(1.0); other 05Ma59=2420(4)							
$^{76}\text{Ni}^m$	T: others 14Ra20=636(90) 12Ka36=409(+58-50) 05Ma59=590(+180-110)							
$^{76}\text{Cu}$	T: average 10Ho12=599(18) 05Va19=653(24) 91Kr15=641(6)							
$^{76}\text{Cu}$	J: from 05Va19 and 90Wi12							
$^{76}\text{Ge}$	T: symmetrized from 15Ba11=1.65(+0.14-0.12)							
$^{76}\text{Ge}$	T: 13Ag11: 0v- $\beta\beta$ >30Yy (90% C.L.) combined GERDA+HDM+IGEX results;							
$^{76}\text{Ge}$	T: claim for 0v- $\beta\beta$ 01K113=15 Yy 04K103=11.2 Yy not trusted. See also							
$^{76}\text{Ge}$	T: 02Aa.A and 02Zd02							
$^{76}\text{Y}$	I: also 00We.A >170 ns same group							
$^{77}\text{Co}$	-21020#	600#	15 ms	6	7/2 <sup>-</sup> #	14	2014	$\beta^-=100$ ; $\beta^-n=90\#$ ; $\beta^-2n=5\#$
$^{77}\text{Ni}$	-36800#	500#	158.9 ms	4.2	9/2 <sup>+</sup> #	12	14Xu07	T 1995 $\beta^-=100$ ; $\beta^-n=30$ 24; $\beta^-2n=0\#$
$^{77}\text{Cu}$	-48620#	150#	467.9 ms	2.1	5/2 <sup>-</sup>	12	1987	$\beta^-=100$ ; $\beta^-n=30.3$ 20
$^{77}\text{Zn}$	-58789.2	2.0	2.08 s	0.05	(7/2 <sup>+</sup> )	12	1977	$\beta^-=100$
$^{77}\text{Zn}^m$	-58016.8	2.0	772.440	0.015	1.05 s	0.10	(1/2 <sup>-</sup> )	12 09Pa35 J 1986 $\beta^-=66$ 7; IT=34 7
$^{77}\text{Ga}$	-65992.3	2.4	13.2 s	0.2	3/2 <sup>(-)</sup>	12	1968	$\beta^-=100$
$^{77}\text{Ge}$	-71212.86	0.05	11.211 h	0.003	7/2 <sup>+</sup>	12	1939	$\beta^-=100$
$^{77}\text{Ge}^m$	-71053.15	0.08	159.71	0.06	53.7 s	0.6	1/2 <sup>-</sup>	12 1947 $\beta^-=81$ 2; IT=19 2
$^{77}\text{As}$	-73916.3	1.7	38.79 h	0.05	3/2 <sup>-</sup>	12	1951	$\beta^-=100$
$^{77}\text{As}^m$	-73440.8	1.7	475.48	0.04	114.0 $\mu\text{s}$	2.5	9/2 <sup>+</sup>	12 1957 IT=100
$^{77}\text{Se}$	-74599.49	0.06	STABLE		1/2 <sup>-</sup>	12	1922	IS=7.63 16
$^{77}\text{Se}^m$	-74437.57	0.06	161.9223	0.0010	17.36 s	0.05	7/2 <sup>+</sup>	12 1947 IT=100
$^{77}\text{Br}$	-73234.8	2.8	57.04 h	0.12	3/2 <sup>-</sup>	12	1948	$\beta^+=100$
$^{77}\text{Br}^m$	-73128.9	2.8	105.86	0.08	4.28 m	0.10	9/2 <sup>+</sup>	12 1961 IT=100
$^{77}\text{Kr}$	-70169.4	2.0	74.4 m	0.6	5/2 <sup>+</sup>	12	1948	$\beta^+=100$
$^{77}\text{Kr}^m$	-70102.9	2.0	66.50	0.05	118 ns	12	3/2 <sup>-</sup>	12 1975 IT=100
$^{77}\text{Rb}$	-64830.5	1.3	3.78 m	0.04	3/2 <sup>-</sup>	12	1972	$\beta^+=100$
$^{77}\text{Sr}$	-57803	8	9.0 s	0.2	5/2 <sup>+</sup>	12	13Ma15	J 1976 $\beta^+=100$ ; $\beta^+p=0.08$ 3
$^{77}\text{Y}$	-46440#	200#	63 ms	17	5/2 <sup>+</sup> #	12	00We.A	D 1999 $\beta^+=?$ ; $\beta^+p ?$ ; p<10
$^{77}\text{Zr}$	-32040#	400#	100# $\mu\text{s}$		3/2 <sup>-</sup> #			$\beta^+ ?$ ; $\beta^+p ?$ ; p ?
$^{77}\text{Co}$	T: symmetrized from 14Xu07=13.0(+7.2-4.3)							
$^{77}\text{Ni}$	D: from 10Ho12							
$^{77}\text{Y}$	D: limit for p is from 00We.A		T: symmetrized from 01Ki13=57(+22-12)					
$^{78}\text{Ni}$	-33890#	600#	122.2 ms	5.1	0 <sup>+</sup>	09	14Xu07	T 1995 $\beta^-=100$ ; $\beta^-n=50\#$ ; $\beta^-2n=0\#$
$^{78}\text{Cu}$	-44500	500	330.7 ms	2.0	(5 <sup>-</sup> )	09	14Xu07	T 1991 $\beta^-=100$ ; $\beta^-n=50.6$ 45; $\beta^-2n=0.2\#$
$^{78}\text{Zn}$	-57483.2	1.9	1.47 s	0.15	0 <sup>+</sup>	09	1977	$\beta^-=100$ ; $\beta^-n=0\#$
$^{78}\text{Zn}^m$	-54807.9	2.1	2675.3	1.0	320 ns	6	(8 <sup>+</sup> )	09 12Ka36 ET 1998 IT=100
$^{78}\text{Ga}$	-63706.0	1.9	5.09 s	0.05	2 <sup>-</sup>	09	11Ma45	J 1972 $\beta^-=100$
$^{78}\text{Ga}^m$	-63207.1	2.0	498.9	0.5	110 ns	3	09	10Da06 ET 2010 IT=100
$^{78}\text{Ge}$	-71862	4	88.0 m	1.0	0 <sup>+</sup>	09	1953	$\beta^-=100$
$^{78}\text{As}$	-72817	10	90.7 m	0.2	2 <sup>-</sup>	09	1937	$\beta^-=100$
$^{78}\text{Se}$	-77025.94	0.18	STABLE		0 <sup>+</sup>	09	1922	IS=23.77 28
$^{78}\text{Br}$	-73452	4	6.45 m	0.04	1 <sup>+</sup>	09	1937	$\beta^+\approx 100$ ; $\beta^- < 0.01$
$^{78}\text{Br}^m$	-73271	4	180.89	0.13	119.4 $\mu\text{s}$	1.0	(4 <sup>+</sup> )	09 1958 IT=100
$^{78}\text{Kr}$	-74178.3	0.3	STABLE		(>110 Ey)		0 <sup>+</sup>	09 94Sa31 T 1920 IS=0.355 3; 2 $\beta^+ ?$
$^{78}\text{Rb}$	-66935	3	17.66 m	0.03	0 <sup>(+)</sup>	09	1968	$\beta^+=100$
$^{78}\text{Rb}^m$	-66888	3	46.84	0.14	910 ns	40	(1 <sup>-</sup> )	09 1996 IT=100
$^{78}\text{Rb}^n$	-66824	3	111.19	0.22	5.74 m	0.03	4 <sup>(-)</sup>	09 1968 $\beta^+=91$ 2; IT=9 2
$^{78}\text{Rb}^x$	-66861	12	74	12	R = 2.0 0.5		spmix	
$^{78}\text{Sr}$	-63174	7	156.1 s	2.7	0 <sup>+</sup>	09	11Pe29	T 1982 $\beta^+=100$
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...										
<sup>78</sup> Y	-52170#	300#		*	54 ms	5	(0 <sup>+</sup> )	09 01Ga24 TJ	1992	$\beta^+=100; \beta^+p?$
<sup>78</sup> Y <sup>m</sup>	-52170#	580#	0#	500#	5.8 s	0.6	(5 <sup>+</sup> )	09	1998	$\beta^+=100; \beta^+p?$
<sup>78</sup> Zr	-40850#	400#			50# ms	(>200 ns)	0 <sup>+</sup>	09 01Ki13 I	2001	$\beta^+?; \beta^+p?$
* <sup>78</sup> Cu	D: $\beta^-n$ average 10Ho12=44.0(5.4)% 09Wi03=65(8)%									
* <sup>78</sup> Cu	J: from 12Ko29; other 11Ko36 (6 <sup>-</sup> )									
* <sup>78</sup> Zn <sup>m</sup>	E: 12Ka36=145.7(0.5), 730.0(0.5), 890.5(0.5) and 909.1(0.5) $\gamma$ rays in									
* <sup>78</sup> Zn <sup>m</sup>	E: cascade to ground-state =2675.3(1.0)									
* <sup>78</sup> Zn <sup>m</sup>	T: average 12Ka36=320(+9-8) 00Da07=319(9)									
* <sup>78</sup> Ga <sup>m</sup>	E: this is level 559.6(0.7) <500 ns in ENSDF'09									
* <sup>78</sup> Kr	T: limit given here is for the K-e <sup>+</sup> decay (theoretically faster)									
* <sup>78</sup> Sr	T: average 11Pe29=155(3) 97Mu02=168(12) 92Gr09=159(8)									
* <sup>78</sup> Y	T: average 01Ga24=50(8) 01Ki13=55(+9-6)									
* <sup>78</sup> Zr	I: also 00We.A>170 ns same group									
<sup>79</sup> Ni	-27570#	600#			44 ms	8	5/2 <sup>+</sup> #	16	2010	$\beta^-=100; \beta^-n=60\#; \beta^-2n=40\#$
<sup>79</sup> Cu	-41740#	300#			241.0 ms	3.2	5/2 <sup>-</sup> #	16	1991	$\beta^-=100; \beta^-n=66\ 12; \beta^-2n=0\#$
<sup>79</sup> Zn	-53432.3	2.2			746 ms	42	9/2 <sup>+</sup>	16	1981	$\beta^-=100; \beta^-n=1.7\ 5$
<sup>79</sup> Zn <sup>m</sup>	-52330	150	1100	150	> 200 ms		1/2 <sup>+</sup>	16	2015	IT=?; $\beta^-?$
<sup>79</sup> Ga	-62547.7	1.9			2.848 s	0.003	3/2 <sup>(-)</sup>	16	1974	$\beta^-=100; \beta^-n=0.089\ 19$
<sup>79</sup> Ge	-69530	40			18.98 s	0.03	(1/2) <sup>-</sup>	16	1970	$\beta^-=100$
<sup>79</sup> Ge <sup>m</sup>	-69340	40	185.95	0.04	39.0 s	1.0	7/2 <sup>+</sup> #	16	1970	$\beta^-=96\ 1; IT=4\ 1$
<sup>79</sup> As	-73636	5			9.01 m	0.15	3/2 <sup>-</sup>	16	1950	$\beta^-=100$
<sup>79</sup> As <sup>m</sup>	-72863	5	772.81	0.06	1.21 $\mu$ s	0.01	(9/2) <sup>+</sup>	16	98Gr14 T	IT=100
<sup>79</sup> Se	-75917.46	0.22			327 ky	28	7/2 <sup>+</sup>	16	1950	$\beta^-=100$
<sup>79</sup> Se <sup>m</sup>	-75821.69	0.22	95.77	0.03	3.92 m	0.01	1/2 <sup>-</sup>	16	1950	IT $\approx$ 100; $\beta^-=0.056\ 11$
<sup>79</sup> Br	-76068.0	1.0			STABLE		3/2 <sup>-</sup>	16	1920	IS=50.69 7
<sup>79</sup> Br <sup>m</sup>	-75860.4	1.0	207.61	0.09	4.85 s	0.04	9/2 <sup>+</sup>	16	1954	IT=100
<sup>79</sup> Kr	-74442	3			35.04 h	0.10	1/2 <sup>-</sup>	16	1948	$\beta^+=100$
<sup>79</sup> Kr <sup>m</sup>	-74312	3	129.77	0.05	50 s	3	7/2 <sup>+</sup>	16	1940	IT=100
<sup>79</sup> Rb	-70803.0	2.1			22.9 m	0.5	5/2 <sup>+</sup>	16	1957	$\beta^+=100$
<sup>79</sup> Sr	-65477	8			2.25 m	0.10	3/2 <sup>(-)</sup>	16	1972	$\beta^+=100$
<sup>79</sup> Y	-57820	80			14.8 s	0.6	5/2 <sup>+</sup> #	16	1992	$\beta^+=100$
<sup>79</sup> Zr	-46770#	300#			56 ms	30	5/2 <sup>+</sup> #	16	1999	$\beta^+=100; \beta^+p?$
<sup>79</sup> Nb	-31650#	500#					9/2 <sup>+</sup> #			$p?; \beta^+?; \beta^+p?$
<sup>80</sup> Ni	-22630#	700#			30 ms	22	0 <sup>+</sup>	14	2014	$\beta^-=100; \beta^-n=60\#; \beta^-2n=40\#$
<sup>80</sup> Cu	-36200#	400#			113.3 ms	6.4		14 14Xu07 T	1995	$\beta^-?; \beta^-n=40\#; \beta^-2n=20\#$
<sup>80</sup> Zn	-51648.6	2.6			562.2 ms	3.0	0 <sup>+</sup>	14 14Xu07 T	1981	$\beta^-=100; \beta^-n=1.0\ 5$
<sup>80</sup> Ga	-59223.7	2.9			1.9 s	0.1	6 <sup>(-)</sup>	14 13Ve03 TJ	1974	$\beta^-=100; \beta^-n=0.86\ 7$
<sup>80</sup> Ga <sup>m</sup>	-59201.3	2.9	22.45	0.10	1.3 s	0.2	3 <sup>(-)</sup>	14 13Ve03 TJ	2011	$\beta^-?; \beta^-n=1\#; IT?$
<sup>80</sup> Ge	-69535.3	2.1			29.5 s	0.4	0 <sup>+</sup>	05	1972	$\beta^-=100$
<sup>80</sup> As	-72214	3			15.2 s	0.2	1 <sup>+</sup>	05	1954	$\beta^-=100$
<sup>80</sup> Se	-77759.5	1.0			STABLE		0 <sup>+</sup>	05	1922	IS=49.61 41; 2 $\beta^-?$
<sup>80</sup> Br	-75889.0	1.0			17.68 m	0.02	1 <sup>+</sup>	05	1937	$\beta^-=91.7\ 2; \beta^+=8.3\ 2$
<sup>80</sup> Br <sup>m</sup>	-75803.2	1.0	85.843	0.004	4.4205 h	0.0008	5 <sup>-</sup>	05	1937	IT=100
<sup>80</sup> Kr	-77893.3	0.7			STABLE		0 <sup>+</sup>	05	1920	IS=2.286 10
<sup>80</sup> Rb	-72175.5	1.9			33.4 s	0.7	1 <sup>+</sup>	05 93Al03 T	1961	$\beta^+=100$
<sup>80</sup> Rb <sup>m</sup>	-71681.6	2.0	493.9	0.5	1.63 $\mu$ s	0.04	(6 <sup>+</sup> )	05 92Do10 E	1980	IT=100
<sup>80</sup> Sr	-70311	3			106.3 m	1.5	0 <sup>+</sup>	05	1961	$\beta^+=100$
<sup>80</sup> Y	-61148	6			30.1 s	0.5	4 <sup>-</sup>	05	1981	$\beta^+=100$
<sup>80</sup> Y <sup>m</sup>	-60920	6	228.5	0.1	4.8 s	0.3	1 <sup>-</sup>	05 01No07 J	1998	IT=81 2; $\beta^+=19\ 2$
<sup>80</sup> Y <sup>n</sup>	-60835	6	312.6	0.9	4.7 $\mu$ s	0.3	(2 <sup>+</sup> )	05	1997	IT=100
<sup>80</sup> Zr	-54360#	300#			4.6 s	0.6	0 <sup>+</sup>	05 01Ki13 T	1987	$\beta^+=100$
<sup>80</sup> Nb	-38420#	400#					4 <sup>-</sup> #			$p?; \beta^+?; \beta^+p?$
* <sup>80</sup> Ni	T: symmetrized from 14Xu07=23.9(+26.0-17.2)									
* <sup>80</sup> Y <sup>m</sup>	J: 228.5 M3 $\gamma$ ray to 4 <sup>-</sup> level									
* <sup>80</sup> Zr	T: average 01Ki13=5.3(+1.1-0.9) 00Re03=4.1(+0.8-0.6)									
<sup>81</sup> Cu	-31420#	500#			73.2 ms	6.8	5/2 <sup>-</sup> #	10 14Xu07 TD	2010	$\beta^-=100; \beta^-n=70\#; \beta^-2n=30\#$
<sup>81</sup> Zn	-46200	5			303.2 ms	2.6	(5/2 <sup>+</sup> )	08 14Xu07 T	1991	$\beta^-=100; \beta^-n=9.1\ 24; \beta^-2n=0\#$
<sup>81</sup> Ga	-57628	3			1.217 s	0.005	5/2 <sup>+</sup>	08 11Ch16 J	1976	$\beta^-=100; \beta^-n=11.9\ 7$
<sup>81</sup> Ge	-66291.7	2.1			8 s	2	9/2 <sup>+</sup> #	08	1972	$\beta^-=100$
<sup>81</sup> Ge <sup>m</sup>	-65612.6	2.1	679.14	0.04	8 s	2	(1/2 <sup>+</sup> )	08	1981	$\beta^-\approx 100; IT<1$
<sup>81</sup> As	-72533.3	2.6			33.3 s	0.8	3/2 <sup>-</sup>	08	1960	$\beta^-=100$
<sup>81</sup> Se	-76389.0	1.0			18.45 m	0.12	1/2 <sup>-</sup>	08	1948	$\beta^-=100$
<sup>81</sup> Se <sup>m</sup>	-76286.0	1.0	103.00	0.06	57.28 m	0.02	7/2 <sup>+</sup>	08	1971	IT $\approx$ 100; $\beta^-=0.051\ 14$
... A-group is continued on next page ...										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>81</sup> Br	-77977.0	1.0	STABLE	3/2 <sup>-</sup>	08		1920	IS=49.31 7	
<sup>81</sup> Br <sup>m</sup>	-77440.8	1.0	536.20 0.09	34.6 μs	2.8		1967	IT=100	
<sup>81</sup> Kr	-77696.2	1.1		229 ky	11		1950	ε=100	
<sup>81</sup> Kr <sup>m</sup>	-77505.6	1.1	190.64 0.04	13.10 s	0.03		1940	IT≈100; ε=0.0025 4	
<sup>81</sup> Rb	-75457	5		4.572 h	0.004		1949	β <sup>+</sup> =100	
<sup>81</sup> Rb <sup>m</sup>	-75371	5	86.31 0.07	30.5 m	0.3		1956	IT=97.6 6; β <sup>+</sup> =2.4 6	
<sup>81</sup> Sr	-71528	3		22.3 m	0.4		1952	β <sup>+</sup> =100	
<sup>81</sup> Sr <sup>m</sup>	-71449	3	79.23 0.04	390 ns	50		1983	IT=100	
<sup>81</sup> Sr <sup>n</sup>	-71439	3	89.05 0.07	6.4 μs	0.5		1989	IT ?	
<sup>81</sup> Y	-65713	5		70.4 s	1.0		1981	β <sup>+</sup> =100	
<sup>81</sup> Zr	-57460	90		5.5 s	0.4		1997	β <sup>+</sup> =100; β <sup>+</sup> p=0.12 2	
<sup>81</sup> Nb	-46360#	400#		< 44 ns			00We.A I	p ?; β <sup>+</sup> ?; β <sup>+</sup> p ?	
<sup>81</sup> Mo	-31750#	500#		1# ms (>400 ns)			13Su23 I	β <sup>+</sup> ?; β <sup>+</sup> p ?	
* <sup>81</sup> Zn	D : β <sup>-</sup> n average 12Ma37=12(4) 91Kr15=7.5(3.0)%; other 10Ho12=30(13)% not used								
* <sup>81</sup> Ge	T : derived from 7.6(0.6), for mixture of ground-state and isomer with almost same half-life								
* <sup>81</sup> Nb	I : also 99Ja02<80 01Ki13<200 ns T : estimated half-life for β <sup>+</sup> : 100# ms								
<sup>82</sup> Cu	-25320#	600#		50# ms (>400 ns)		10 100h02 I	2010	β <sup>-</sup> ?; β <sup>-</sup> n=30#; β <sup>-</sup> 2n=60#	
<sup>82</sup> Zn	-42314	3		177.9 ms	2.5	0 <sup>+</sup>	12 16A110 D	1997	β <sup>-</sup> =100; β <sup>-</sup> n=69 7; β <sup>-</sup> 2n=0#
<sup>82</sup> Ga	-52930.7	2.4		599 ms	2	(2)	03 12Ch51 J	1976	β <sup>-</sup> =100; β <sup>-</sup> n=21.3 13; β <sup>-</sup> 2n=0#
<sup>82</sup> Ga <sup>m</sup>	-52789.7	2.5	141.0 0.5	93.5 ns	6.7	(4 <sup>-</sup> )	16A110 TJ	2009	IT=100
<sup>82</sup> Ge	-65415.1	2.2		4.56 s	0.26	0 <sup>+</sup>	11	1972	β <sup>-</sup> =100
<sup>82</sup> As	-70105	4		19.1 s	0.5	(2 <sup>-</sup> )	03 04Ga44 J	1968	β <sup>-</sup> =100
<sup>82</sup> As <sup>m</sup>	-69973	4	132.1 0.2	13.6 s	0.4	(5 <sup>-</sup> )	03 14Mi16 E	1970	β <sup>-</sup> =100
<sup>82</sup> Se	-77593.9	0.5		92 Ey	7	0 <sup>+</sup>	03 15Ba11 T	1922	IS=8.73 22; 2β <sup>-</sup> =100
<sup>82</sup> Br	-77498.7	1.0		35.282 h	0.007	5 <sup>-</sup>	03	1937	β <sup>-</sup> =100
<sup>82</sup> Br <sup>m</sup>	-77452.8	1.0	45.9492 0.0010	6.13 m	0.05	2 <sup>-</sup>	03	1965	IT=97.6 3; β <sup>-</sup> =2.4 3
<sup>82</sup> Kr	-80591.785	0.005		STABLE		0 <sup>+</sup>	03	1920	IS=11.593 31
<sup>82</sup> Rb	-76188	3		1.273 m	0.002	1 <sup>+</sup>	03	1949	β <sup>+</sup> =100
<sup>82</sup> Rb <sup>m</sup>	-76118.8	2.6	69.0 1.5	6.472 h	0.006	5 <sup>-</sup>	03	1957	β <sup>+</sup> ≈100; IT<0.33
<sup>82</sup> Sr	-76010	6		25.36 d	0.03	0 <sup>+</sup>	03 87Ho06 T	1952	ε=100
<sup>82</sup> Y	-68064	5		8.30 s	0.20	1 <sup>+</sup>	03	1980	β <sup>+</sup> =100
<sup>82</sup> Y <sup>m</sup>	-67661	5	402.63 0.14	258 ns	22	4 <sup>-</sup>	03 94Mu02 T	1994	IT=100
<sup>82</sup> Y <sup>n</sup>	-67557	5	507.50 0.13	147 ns	7	6 <sup>+</sup>	03	1994	IT=100
<sup>82</sup> Zr	-63631	11		32 s	5	0 <sup>+</sup>	03	1982	β <sup>+</sup> =100
<sup>82</sup> Nb	-52090#	300#		50 ms	5	(0 <sup>+</sup> )	08	1992	β <sup>+</sup> =100; β <sup>+</sup> p ?
<sup>82</sup> Nb <sup>m</sup>	-50910#	300#	1180 1	92 ns	17	(5 <sup>+</sup> )	08 08Ga04 ETJ	2008	IT=100
<sup>82</sup> Mo	-40370#	400#		30# ms (>400 ns)		0 <sup>+</sup>	15 13Su23 I	2013	β <sup>+</sup> ?; β <sup>+</sup> p ?
* <sup>82</sup> Zn	T : 14Xu07=177.9(2.5); others 16A110=155(26) 12Ma37=228(10)								
* <sup>82</sup> Ga	D : average 93Ru01=31.1(4.4)% 86Wa17=19.8(1.7)% 80Lu04=21.4(2.2)%								
* <sup>82</sup> Ga <sup>m</sup>	T : average 16A110=89(9) 12Ka36=98(+10-9); other 09Fo05<500 ns								
* <sup>82</sup> As <sup>m</sup>	E : 15Et01=131.6(1.5) 14Mi16=132.1(0.2)								
* <sup>82</sup> Sr	T : average 87Ho06=25.36(0.03) 87Ju02=25.342(0.053)								
* <sup>82</sup> Y <sup>m</sup>	T : average 94Mu02=220(50) 93Wo04=268(25)								
<sup>83</sup> Zn	-36290#	300#		119 ms	16	3/2 <sup>+</sup> #	15 16A110 T	1997	β <sup>-</sup> =100; β <sup>-</sup> n=10#; β <sup>-</sup> 2n=3#
<sup>83</sup> Ga	-49257.1	2.6		308.1 ms	1.0	5/2 <sup>-</sup> #	15	1976	β <sup>-</sup> =100; β <sup>-</sup> n=62.8 25; β <sup>-</sup> 2n=0#
<sup>83</sup> Ga <sup>m</sup>	-49059.8	2.6	197.3 0.5	120 ns	5		16A110 ETD	2016	IT=100
<sup>83</sup> Ge	-60976.4	2.4		1.85 s	0.06	(5/2 <sup>+</sup> )	15	1972	β <sup>-</sup> =100; β <sup>-</sup> n=0.1#
<sup>83</sup> As	-69669.3	2.8		13.4 s	0.4	5/2 <sup>-</sup> #	15	1968	β <sup>-</sup> =100
<sup>83</sup> Se	-75341	3		22.25 m	0.04	9/2 <sup>+</sup>	15 15Kr02 T	1937	β <sup>-</sup> =100
<sup>83</sup> Se <sup>m</sup>	-75112	3	228.92 0.07	70.1 s	0.4	1/2 <sup>-</sup>	15	1969	β <sup>-</sup> =100
<sup>83</sup> Br	-79014	4		2.374 h	0.004	3/2 <sup>-</sup>	15	1937	β <sup>-</sup> =100
<sup>83</sup> Br <sup>m</sup>	-75945	4	3069.2 0.4	729 ns	77	(19/2 <sup>-</sup> )	15 11Ru.A T	1989	IT=100
<sup>83</sup> Kr	-79990.633	0.009		STABLE		9/2 <sup>+</sup>	15	1920	IS=11.500 19
<sup>83</sup> Kr <sup>m</sup>	-79981.228	0.009	9.4053 0.0008	156.8 ns	0.5	7/2 <sup>+</sup>	15	1963	IT=100
<sup>83</sup> Kr <sup>n</sup>	-79949.076	0.009	41.5575 0.0007	1.830 h	0.013	1/2 <sup>-</sup>	15 10Li13 T	1971	IT=100
<sup>83</sup> Rb	-79070.6	2.3		86.2 d	0.1	5/2 <sup>-</sup>	15	1950	ε=100
<sup>83</sup> Rb <sup>m</sup>	-79028.5	2.3	42.0780 0.0020	7.8 ms	0.7	9/2 <sup>+</sup>	15 68Et01 T	1968	IT=100
<sup>83</sup> Sr	-76798	7		32.41 h	0.03	7/2 <sup>+</sup>	15	1952	β <sup>+</sup> =100
<sup>83</sup> Sr <sup>m</sup>	-76539	7	259.15 0.09	4.95 s	0.12	1/2 <sup>-</sup>	15	1972	IT=100
<sup>83</sup> Y	-72206	19		7.08 m	0.08	(9/2 <sup>+</sup> )	15 92Bu10 J	1962	β <sup>+</sup> =100
<sup>83</sup> Y <sup>m</sup>	-72144	19	62.04 0.10	2.85 m	0.02	(3/2 <sup>-</sup> )	15	1972	β <sup>+</sup> =60 5; IT=40 5
<sup>83</sup> Zr	-65912	6		42 s	2	1/2 <sup>-</sup> #	15	1974	β <sup>+</sup> =100; β <sup>+</sup> p=?
<sup>83</sup> Zr <sup>m</sup>	-65859	6	52.72 0.05	530 ns	120	(5/2 <sup>-</sup> )	15	1988	IT=100
<sup>83</sup> Zr <sup>n</sup>	-65835	6	77.04 0.07	1.8 μs	0.1	(7/2 <sup>+</sup> )	15	1988	IT=100
<sup>83</sup> Nb	-57560	150		3.9 s	0.2	(9/2 <sup>+</sup> )	15 GAu15b J	1988	β <sup>+</sup> =100
... A-group is continued on next page ...									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>83</sup> Mo	-46340#	400#	23 ms	19		15 01Ki13	TD	1999	$\beta^+=100; \beta^+p?$	*	
<sup>83</sup> Tc	-31320#	500#							$p?; \beta^+?; \beta^+p?$	**	
* <sup>83</sup> Zn	T : average 16Al10=122(28) 12Ma37=117(20)										
* <sup>83</sup> Br <sup>m</sup>	T : average 11Ru.A=862(148) 97Is13=700(100) 89Wi01=600(200)										
* <sup>83</sup> Kr <sup>n</sup>	T : average 10Li13=1.82(0.02) 09Ka30=1.85(0.03) 71Ru17=1.83(0.02)										
* <sup>83</sup> Nb	J : ENSDF=(5/2 <sup>+</sup> , 7/2 <sup>+</sup> , 9/2 <sup>+</sup> ); TNN trends in N=41 isotopes suggests (9/2 <sup>+</sup> )										
* <sup>83</sup> Mo	T : symmetrized from 01Ki13=6(+30-3)										
<sup>84</sup> Zn	-31930#	400#	50# ms	(>400 ns)	0 <sup>+</sup>	10 100h02	I	2010	$\beta^-?; \beta^-n=40\#; \beta^-2n=4\#$		
<sup>84</sup> Ga	-44090#	200#	85 ms	10	0 <sup>-</sup>	09 16Ma50	D	1991	$\beta^-n=100; \beta^-n=40?; \beta^-2n=2\#$	*	
<sup>84</sup> Ge	-58148	3	951 ms	9	0 <sup>+</sup>	09 13Ma22	T	1972	$\beta^-n=100; \beta^-n=10.7?; 6$	**	
<sup>84</sup> As	-65854	3	4.02 s	0.03	(3 <sup>-</sup> )	09 93Ru01	TD	1968	$\beta^-n=100; \beta^-n=0.28?; 4$	*	
<sup>84</sup> As <sup>m</sup>	-65850#	100#	0#	100#	*	650 ms	150	09	1974	$\beta^-n=100$	*
<sup>84</sup> Se	-75947.7	2.0	3.26 m	0.10	0 <sup>+</sup>	09		1960	$\beta^-n=100$		
<sup>84</sup> Br	-77783	26	31.76 m	0.08	2 <sup>-</sup>	09		1943	$\beta^-n=100$		
<sup>84</sup> Br <sup>m</sup>	-77470	100	310	100	BD	6.0 m	0.2	09	1957	$\beta^-n=100$	
<sup>84</sup> Br <sup>n</sup>	-77375	26	408.2	0.4		< 140 ns		09	1970	IT=100	
<sup>84</sup> Kr	-82439.335	0.004	STABLE		0 <sup>+</sup>	09		1920	IS=56.987 15		
<sup>84</sup> Kr <sup>m</sup>	-79203.27	0.18	3236.07	0.18		1.83 $\mu$ s	0.04	09	1982	IT=100	
<sup>84</sup> Rb	-79759.0	2.2				32.82 d	0.07	09	1947	$\beta^+=96.1?; 20; \beta^-n=3.9?; 20$	
<sup>84</sup> Rb <sup>m</sup>	-79295.4	2.2	463.59	0.08		20.26 m	0.04	09	1940	IT $\approx$ 100; $\beta^+<0.0012$	
<sup>84</sup> Sr	-80649.6	1.2	STABLE		0 <sup>+</sup>	09		1936	IS=0.56 1; $2\beta^+?$		
<sup>84</sup> Y	-73894	4	39.5 m	0.8	(6 <sup>+</sup> )	09		1962	$\beta^+=100$		
<sup>84</sup> Y <sup>m</sup>	-73827	4	67.0	0.2		4.6 s	0.2	09	1976	$\beta^+=100$	
<sup>84</sup> Y <sup>n</sup>	-73684	4	210.42	0.16		292 ns	10	09	2005	IT=100	
<sup>84</sup> Zr	-71422	5	25.8 m	0.5	0 <sup>+</sup>	09		1977	$\beta^+=100$		
<sup>84</sup> Nb	-61219	13	9.8 s	0.9	(1 <sup>+</sup> )	09 09St04	J	1977	$\beta^+=100$		
<sup>84</sup> Nb <sup>m</sup>	-61171	13	48	1		176 ns	46	09	2009	IT=100	
<sup>84</sup> Nb <sup>n</sup>	-60881	13	337.7	0.4		92 ns	5	09	2000	IT=100	
<sup>84</sup> Mo	-54170#	300#	2.3 s	0.3	0 <sup>+</sup>	09		1991	$\beta^+=100; \beta^+p?$		
<sup>84</sup> Tc	-37700#	400#			1 <sup>+</sup> #				$p?; \beta^+?; \beta^+p?$	**	
* <sup>84</sup> Ga	D : $\beta^-n$ others 10Wi03=74(14)% 91Kr15=70(15)%										
* <sup>84</sup> Ga	I : a $\beta^-$ decaying isomer was identified in 09Le26 and adopted in ENSDF'2009,										
* <sup>84</sup> Ga	I : questioned in 10Wi03										
* <sup>84</sup> Ge	T : average 13Ma22=942(17) 93Ru01=947(11) 91Kr15=984(23)										
* <sup>84</sup> Ge	D : average 93Ru01=10.8(0.6)% 91Kr15=9.5(2.0)%										
* <sup>84</sup> As	J : 16Ko24 proposed (2 <sup>-</sup> )										
* <sup>84</sup> As <sup>m</sup>	I : identification discussed in ENSDF2009										
<sup>85</sup> Zn	-25230#	500#	50# ms	(>400 ns)	5/2 <sup>+</sup> #	14 100h02	I	2010	$\beta^-?; \beta^-n=30\#; \beta^-2n=7\#$		
<sup>85</sup> Ga	-39850#	300#	92.2 ms	3.5	(5/2 <sup>-</sup> )	14 13Mi19	T	1997	$\beta^-n=100; \beta^-n>35; \beta^-2n=6\#$	*	
<sup>85</sup> Ge	-53123	4	494 ms	8	(3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	14 13Ma22	T	1991	$\beta^-n=100; \beta^-n=16.5?; 23; \beta^-2n=0\#$	*	
<sup>85</sup> As	-63189	3	2.021 s	0.012	(5/2 <sup>-</sup> )	14 12Ku06	J	1967	$\beta^-n=100; \beta^-n=62.9?; 20$		
<sup>85</sup> Se	-72413.6	2.6	32.9 s	0.3	(5/2 <sup>+</sup> )	14		1960	$\beta^-n=100$		
<sup>85</sup> Br	-78575	3	2.90 m	0.06	3/2 <sup>-</sup>	14		1943	$\beta^-n=100$		
<sup>85</sup> Kr	-81480.3	2.0	10.739 y	0.014	9/2 <sup>+</sup>	14		1940	$\beta^-n=100$		
<sup>85</sup> Kr <sup>m</sup>	-81175.4	2.0	304.871	0.020		4.480 h	0.008	14	1937	$\beta^-n=78.8?; 5; IT=21.2?; 5$	
<sup>85</sup> Kr <sup>n</sup>	-79488.5	2.0	1991.8	0.2		1.82 $\mu$ s	0.05	14	1989	IT=100	
<sup>85</sup> Rb	-82167.331	0.005	STABLE		5/2 <sup>-</sup>	14		1921	IS=72.17 2		
<sup>85</sup> Rb <sup>m</sup>	-81653.325	0.005	514.0065	0.0022		1.015 $\mu$ s	0.001	14	1964	IT=100	
<sup>85</sup> Sr	-81103.3	2.8	64.849 d	0.007	9/2 <sup>+</sup>	14		1940	$\epsilon=100$		
<sup>85</sup> Sr <sup>m</sup>	-80864.5	2.8	238.79	0.05		67.63 m	0.04	14	1940	IT=86.6 4; $\beta^+=13.4?; 4$	
<sup>85</sup> Y	-77842	19	2.68 h	0.05	(1/2 <sup>-</sup> )	14		1952	$\beta^+=100$		
<sup>85</sup> Y <sup>m</sup>	-77822	19	19.68	0.17		4.86 h	0.20	14	1952	$\beta^+\approx 100; IT<0.002$	
<sup>85</sup> Y <sup>n</sup>	-77576	19	266.18	0.10		178 ns	7	14	1977	IT=100	
<sup>85</sup> Zr	-73175	6	7.86 m	0.04	(7/2 <sup>+</sup> )	14		1963	$\beta^+=100$		
<sup>85</sup> Zr <sup>m</sup>	-72883	6	292.2	0.3		10.9 s	0.3	14	1976	IT<100; $\beta^+>0$	
<sup>85</sup> Nb	-66280	4	20.5 s	0.7	9/2 <sup>+</sup> #	14		1988	$\beta^+=100$	*	
<sup>85</sup> Nb <sup>m</sup>	-66130#	80#	150#	80#		3.3 s	0.9	14	1988	IT=?; $\beta^+?$	*
<sup>85</sup> Mo	-57510	16	3.2 s	0.2	(1/2 <sup>+</sup> )	14 05Xu04	J	1992	$\beta^+=100; \beta^+p=0.14?; 2$		
<sup>85</sup> Tc	-45850#	400#	<110 ns		1/2 <sup>-</sup> #	14 00We.A	I			$p?$	*
<sup>85</sup> Ru	-30950#	500#	1# ms	(>400 ns)	3/2 <sup>-</sup> #	15 13Su23	I	2013	$\beta^+?; \beta^+p?; p?$	**	
* <sup>85</sup> Ga	T : average 13Mi19=92(4) 12Ma37=93(7)										
* <sup>85</sup> Ge	J : (3/2 <sup>+</sup> , 5/2 <sup>+</sup> ) from 13Ko31										
* <sup>85</sup> Nb	T : average 05Ka39=17(2) 88Ku14=20.9(0.7)										
* <sup>85</sup> Nb <sup>m</sup>	E : 05Ka39 > 69 keV										
* <sup>85</sup> Tc	I : also 99Ja02<100 ns T : estimated half-life for $\beta^+$ decay: 100#ms										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>86</sup> Ga	-34080# 400#		47 ms 18			15 13Mi19 TD	1997	$\beta^- = 100; \beta^- n = 60 10; \beta^- 2n = 20 10$ *
<sup>86</sup> Ge	-49400 440		226 ms 21	0 <sup>+</sup>		15	1994	$\beta^- = 100; \beta^- n = 45 15$
<sup>86</sup> As	-58962 3		945 ms 8	(1 <sup>-</sup> , 2 <sup>-</sup> )		15 15Ma61 J	1973	$\beta^- = 100; \beta^- n = 35.5 6; \beta^- 2n = 0.02\#$
<sup>86</sup> Se	-70503.2 2.5		14.3 s 0.3	0 <sup>+</sup>		15	1973	$\beta^- = 100; \beta^- n = 0\#$
<sup>86</sup> Se <sup>m</sup>	-68131.2 2.7	2372.0 1.0	620 ms 240			15Ma61 ET	2015	IT≈100 *
<sup>86</sup> Br	-75632 3		55.1 s 0.4	(1 <sup>-</sup> )		15	1962	$\beta^- = 100$
<sup>86</sup> Kr	-83265.666 0.004		STABLE			15	1920	IS=17.279 41; 2 $\beta^-$ ?
<sup>86</sup> Rb	-82746.99 0.20		18.642 d 0.018	2 <sup>-</sup>		15	1941	$\beta^- \approx 100; \epsilon = 0.0052 5$
<sup>86</sup> Rb <sup>m</sup>	-82190.94 0.27	556.05 0.18	1.017 m 0.003	6 <sup>-</sup>		15	1951	IT≈100; $\beta^- < 0.3$
<sup>86</sup> Sr	-84523.089 0.005		STABLE	0 <sup>+</sup>		15	1931	IS=9.86 1
<sup>86</sup> Sr <sup>m</sup>	-81567.00 0.12	2956.09 0.12	455 ns 7	8 <sup>+</sup>		15	1971	IT=100
<sup>86</sup> Y	-79283 14		14.74 h 0.02	4 <sup>-</sup>		15	1951	$\beta^+ = 100$
<sup>86</sup> Y <sup>m</sup>	-79065 14	218.21 0.09	47.4 m 0.4	(8 <sup>+</sup> )		15	1962	IT=99.31 4; $\beta^+ = 0.69 4$
<sup>86</sup> Y <sup>n</sup>	-78981 14	302.18 0.09	125.3 ns 5.5	6 <sup>+</sup>		15 10Ru07 J	2000	IT=100 *
<sup>86</sup> Zr	-77969 4		16.5 h 0.1	0 <sup>+</sup>		15	1951	$\beta^+ = 100$
<sup>86</sup> Nb	-69134 5		88 s 1	(6 <sup>+</sup> )		15	1974	$\beta^+ = 100$
<sup>86</sup> Nb <sup>m</sup>	-68880# 160#	250# 160#	56.3 s 8.3	high		15 94Sh07 TJD	1994	$\beta^+ = 100$ *
<sup>86</sup> Mo	-64110 4		19.1 s 0.3	0 <sup>+</sup>		15	1991	$\beta^+ = 100$
<sup>86</sup> Tc	-51570# 300#		55 ms 7	(0 <sup>+</sup> )		15	1992	$\beta^+ = 100; \beta^+ p ?$
<sup>86</sup> Tc <sup>m</sup>	-50050# 300#	1524 10	1.10 $\mu$ s 0.12	(6 <sup>+</sup> )		15 08Ga04 T	2000	IT=100 *
<sup>86</sup> Ru	-39770# 400#		50# ms (>400 ns)	0 <sup>+</sup>		15 13Su23 I	2013	$\beta^+ ?; \beta^+ p ?$
* <sup>86</sup> Ga	T : symmetrized from 13Mi19=43(+21-15) **							
* <sup>86</sup> Se <sup>m</sup>	E : error estimated by evaluator **							
* <sup>86</sup> Y <sup>n</sup>	T : average 10Ru07=127(14) 00Io02=125(6) **							
* <sup>86</sup> Nb <sup>m</sup>	I : existence considered as uncertain in ENSDF'15; needs confirmation **							
* <sup>86</sup> Tc <sup>m</sup>	T : average 08Ga04=1.10(0.14) 00Ch07=1.11(0.21) E : unc. estimated by GAu **							
<sup>87</sup> Ga	-29250# 500#		10# ms (>400 ns)	5/2 <sup>-</sup> #		15 10Oh02 I	2010	$\beta^- ?; \beta^- n = 90\#; \beta^- 2n = 7\#$
<sup>87</sup> Ge	-44080# 300#		150# ms (>300 ns)	5/2 <sup>+</sup> #		15 97Be70 I	1997	$\beta^- ?; \beta^- n = 3\#; \beta^- 2n = 1\#$
<sup>87</sup> As	-55617.9 3.0		492 ms 25	(5/2 <sup>-</sup> , 3/2 <sup>-</sup> )		15 15Ko19 TJ	1970	$\beta^- = 100; \beta^- n = 15.4 22; \beta^- 2n = 0\#$ *
<sup>87</sup> Se	-66426.1 2.2		5.50 s 0.14	(3/2 <sup>+</sup> )		15 15Ko19 J	1968	$\beta^- = 100; \beta^- n = 0.36 8$
<sup>87</sup> Br	-73892 3		55.65 s 0.12	(5/2 <sup>-</sup> )		15	1943	$\beta^- = 100; \beta^- n = 2.60 4$
<sup>87</sup> Kr	-80709.52 0.25		76.3 m 0.5	5/2 <sup>+</sup>		15	1940	$\beta^- = 100$
<sup>87</sup> Rb	-84597.791 0.006		49.7 Gy 0.3	3/2 <sup>-</sup>		15	1921	IS=27.83 2; $\beta^- = 100$
<sup>87</sup> Sr	-84880.066 0.005		STABLE	9/2 <sup>+</sup>		15	1931	IS=7.00 1
<sup>87</sup> Sr <sup>m</sup>	-84491.537 0.006	388.5287 0.0023	2.815 h 0.012	1/2 <sup>-</sup>		15	1940	IT≈100; $\epsilon = 0.30 8$
<sup>87</sup> Y	-83018.4 1.1		79.8 h 0.3	1/2 <sup>-</sup>		15	1940	$\beta^+ = 100$
<sup>87</sup> Y <sup>m</sup>	-82637.6 1.1	380.82 0.07	13.37 h 0.03	9/2 <sup>+</sup>		15	1940	IT=98.43 11; $\beta^+ = 1.57 11$
<sup>87</sup> Zr	-79347 4		1.68 h 0.01	9/2 <sup>+</sup>		15	1948	$\beta^+ = 100$
<sup>87</sup> Zr <sup>m</sup>	-79011 4	335.84 0.19	14.0 s 0.2	1/2 <sup>-</sup>		15	1972	IT=100
<sup>87</sup> Nb	-73874 7		3.7 m 0.1	(1/2 <sup>-</sup> )		15	1971	$\beta^+ = 100$
<sup>87</sup> Nb <sup>m</sup>	-73870 7	3.9 0.1	2.6 m 0.1	(9/2 <sup>+</sup> )		15	1972	$\beta^+ = 100$
<sup>87</sup> Mo	-66884.8 2.9		14.1 s 0.3	7/2 <sup>+</sup> #		15	1977	$\beta^+ = 100; \beta^+ p = 15 5$
<sup>87</sup> Tc	-57690 4		2.2 s 0.2	9/2 <sup>+</sup> #		15	1991	$\beta^+ = 100; \beta^+ p ?$
<sup>87</sup> Tc <sup>m</sup>	-57683 4	7 1	2# s	1/2 <sup>-</sup> #		09Ga40 E		$\beta^+ ?; IT ?$ *
<sup>87</sup> Tc <sup>n</sup>	-57619 4	71 1	647 ns 24	7/2 <sup>+</sup> #		15	2009	IT=100
<sup>87</sup> Ru	-45520# 400#		50# ms (>1.5 $\mu$ s)	1/2 <sup>-</sup> #		15 95Ry03 I	1995	$\beta^+ ?; \beta^+ p ?$
* <sup>87</sup> As	T : average 15Ko19=560(80) 13Ma22=484(35) 93Ru01=485(40) **							
* <sup>87</sup> As	T : others 12Qu01=1450(550)(+3900-1100) 78Cr03=730(60) **							
* <sup>87</sup> Tc <sup>m</sup>	E : observed 64 keV $\gamma$ ray in parallel to 71 keV one depopulating <sup>87</sup> Tc <sup>m</sup> **							
<sup>88</sup> Ge	-40140# 400#		100# ms (>300 ns)	0 <sup>+</sup>		14	1997	$\beta^- ?; \beta^- n = 6\#; \beta^- 2n = 0.1\#$
<sup>88</sup> As	-50720# 200#		270 ms 150			14 12Qu01 T	1994	$\beta^- = 100; \beta^- n = 30\#$ *
<sup>88</sup> Se	-63884 3		1.53 s 0.06	0 <sup>+</sup>		14	1970	$\beta^- = 100; \beta^- n = 0.99 10$
<sup>88</sup> Br	-70716 3		16.34 s 0.08	(1 <sup>-</sup> )		14 15Cz01 J	1948	$\beta^- = 100; \beta^- n = 6.58 18$
<sup>88</sup> Br <sup>m</sup>	-70446 3	270.1 0.5	5.51 $\mu$ s 0.04	(4 <sup>-</sup> )		14 11Ru.A T	1970	IT=100 *
<sup>88</sup> Kr	-79691.3 2.6		2.825 h 0.019	0 <sup>+</sup>		14	1939	$\beta^- = 100$
<sup>88</sup> Rb	-82608.99 0.16		17.773 m 0.018	2 <sup>-</sup>		14	1939	$\beta^- = 100$
<sup>88</sup> Rb <sup>m</sup>	-81235.2 0.3	1373.8 0.3	123 ns 13	(7 <sup>+</sup> )		14	2000	IT=100
<sup>88</sup> Sr	-87921.618 0.006		STABLE	0 <sup>+</sup>		14	1923	IS=82.58 1
<sup>88</sup> Y	-84299.0 1.5		106.626 d 0.021	4 <sup>-</sup>		14	1948	$\beta^+ = 100$
<sup>88</sup> Y <sup>m</sup>	-83906.1 1.5	392.86 0.09	301 $\mu$ s 3	1 <sup>+</sup>		14	1955	IT=100
<sup>88</sup> Y <sup>n</sup>	-83624.5 1.5	674.55 0.04	13.98 ms 0.17	8 <sup>+</sup>		14	1962	IT=100
<sup>88</sup> Zr	-83629 5		83.4 d 0.3	0 <sup>+</sup>		14	1951	$\epsilon = 100$
<sup>88</sup> Zr <sup>m</sup>	-80741 5	2887.79 0.06	1.320 $\mu$ s 0.025	8 <sup>+</sup>		14	1978	IT=100
<sup>88</sup> Nb	-76170 60		14.50 m 0.11	(8 <sup>+</sup> )		14	1964	$\beta^+ = 100$
<sup>88</sup> Nb <sup>m</sup>	-76040 100	130 120	7.7 m 0.1	(4 <sup>-</sup> )		14	1971	$\beta^+ = 100$
<sup>88</sup> Mo	-72687 4		8.0 m 0.2	0 <sup>+</sup>		14	1971	$\beta^+ = 100$

... A-group is continued on next page ...



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>88</sup> Tc	-61680	150			*			$\beta^+=100; \beta^+p?$
<sup>88</sup> Tc <sup>m</sup>	-61680#	340#	0# 300#		*			$\beta^+=100; \beta^+p?$ *
<sup>88</sup> Tc <sup>n</sup>	-61580#	160#	100# 50#			14 09Ga40 TJ	2009	IT=100
<sup>88</sup> Ru	-54340#	300#				14	1994	$\beta^+=100; \beta^+p?$ *
<sup>88</sup> Rh	-36860#	400#			1# ms			$\beta^+?$
* <sup>88</sup> As	T: symmetrized from 12Qu01=200(5)(+200-90)							
* <sup>88</sup> Br <sup>m</sup>	J: 15Cz01=(4 <sup>-</sup> )							
* <sup>88</sup> Tc <sup>m</sup>	J: 09Ga40 suggest this state to be 2 <sup>+</sup> , plus existence of an isomer 95 keV							
* <sup>88</sup> Tc <sup>m</sup>	J: above this 2 <sup>+</sup> , that decays by E2, with half-life=146(12) ns							
* <sup>88</sup> Ru	T: symmetrized from 01Ki13=1.2(+0.3-0.2)							
<sup>89</sup> Ge	-33730#	400#						$\beta^-?; \beta^-n=20\#; \beta^-2n=2\#$
<sup>89</sup> As	-46800#	300#						$\beta^-?; \beta^-n=100\#; \beta^-2n=0.3\#$
<sup>89</sup> Se	-58992	4						$\beta^+=100; \beta^-n=7.8$ 25
<sup>89</sup> Br	-68274	3						$\beta^+=100; \beta^-n=13.8$ 4
<sup>89</sup> Kr	-76535.8	2.1						$\beta^+=100$ *
<sup>89</sup> Rb	-81712	5						$\beta^+=100$
<sup>89</sup> Sr	-86209.02	0.09						$\beta^+=100$
<sup>89</sup> Y	-87708.4	1.6						IS=100.
<sup>89</sup> Y <sup>m</sup>	-86799.4	1.6	908.97	0.03				IT=100
<sup>89</sup> Zr	-84876	3						$\beta^+=100$
<sup>89</sup> Zr <sup>m</sup>	-84288	3	587.82	0.10				IT=93.77 12; $\beta^+=6.23$ 12
<sup>89</sup> Nb	-80625	24			*			$\beta^+=100$
<sup>89</sup> Nb <sup>m</sup>	-80630#	40#	0# 30#		*			$\beta^+=100$
<sup>89</sup> Mo	-75015	4						$\beta^+=100$
<sup>89</sup> Mo <sup>m</sup>	-74628	4	387.5	0.2				IT=100
<sup>89</sup> Tc	-67395	4						$\beta^+=100$
<sup>89</sup> Tc <sup>m</sup>	-67332	4	62.6	0.5				$\beta^+\approx 100; IT<0.01$
<sup>89</sup> Ru	-58260#	300#						$\beta^+=100; \beta^+p=3.1$ 18 *
<sup>89</sup> Rh	-45860#	360#						$\beta^+?; \beta^+p?; p?$ *
* <sup>89</sup> Kr	J: positive parity, since no $\beta^-$ transition to <sup>89</sup> Rb ground-state							
* <sup>89</sup> Ru	D: $\beta^+p$ symmetrized from 3.0(+1.9-1.7)% T: other recent 12Lo08=2.2(1.2)							
* <sup>89</sup> Rh	T: post-deadline 16Ce02<120 ns							
<sup>90</sup> Ge	-29220#	500#						$\beta^-?; \beta^-n=50\#; \beta^-2n=2\#$
<sup>90</sup> As	-41330#	400#						$\beta^-?; \beta^-n=30\#; \beta^-2n=3\#$
<sup>90</sup> As <sup>m</sup>	-41210#	400#	124.5	0.5				IT=100 *
<sup>90</sup> Se	-55800	330						$\beta^+=100; \beta^-n=1\#$ *
<sup>90</sup> Br	-64000	3						$\beta^+=100; \beta^-n=25.2$ 9 *
<sup>90</sup> Kr	-74959.2	1.9						$\beta^+=100$
<sup>90</sup> Rb	-79364	6						$\beta^+=100$
<sup>90</sup> Rb <sup>m</sup>	-79257	6	106.90	0.03				$\beta^+=100$
<sup>90</sup> Rb <sup>x</sup>	-79293	14	71	12				$\beta^+=97.4$ 4; IT=2.6 4
<sup>90</sup> Sr	-85948.1	2.1						$\beta^+=100$
<sup>90</sup> Y	-86494.1	1.6						$\beta^+=100$
<sup>90</sup> Y <sup>m</sup>	-85812.4	1.6	681.67	0.10				IT $\approx$ 100; $\beta^-=0.0018$ 2
<sup>90</sup> Zr	-88772.54	0.12						IS=51.45 40
<sup>90</sup> Zr <sup>m</sup>	-86453.54	0.12	2319.000	0.010				IT=100
<sup>90</sup> Zr <sup>n</sup>	-85183.12	0.12	3589.419	0.016				IT=100
<sup>90</sup> Nb	-82662	3						$\beta^+=100$
<sup>90</sup> Nb <sup>m</sup>	-82540	3	122.370	0.022				IT=100
<sup>90</sup> Nb <sup>n</sup>	-82537	3	124.67	0.25				IT=100
<sup>90</sup> Nb <sup>p</sup>	-82491	3	171.10	0.10				IT=100
<sup>90</sup> Nb <sup>q</sup>	-82280	3	382.01	0.25				IT=100
<sup>90</sup> Nb <sup>r</sup>	-80782	3	1880.21	0.20				IT=100 *
<sup>90</sup> Mo	-80173	3						$\beta^+=100$
<sup>90</sup> Mo <sup>m</sup>	-77298	3	2874.73	0.15				IT=100
<sup>90</sup> Tc	-70724.7	1.0						$\beta^+=100$
<sup>90</sup> Tc <sup>m</sup>	-70580.7	1.3	144.0	1.7	MD			$\beta^+=100$
<sup>90</sup> Ru	-64884	4						$\beta^+=100$
<sup>90</sup> Rh	-51700#	300#			*			$\beta^+=100; \beta^+p?$ *
<sup>90</sup> Rh <sup>m</sup>	-51700#	580#	0# 500#		*			$\beta^+=100; \beta^+p?$ *
<sup>90</sup> Pd	-39710#	400#						$\beta^+?$
* <sup>90</sup> As <sup>m</sup>	T: symmetrized from 200(+120-90)							
* <sup>90</sup> Se	T: symmetrized from 12Qu01=195(7)(+95-65)							
* <sup>90</sup> Br	T: supersedes 80A115=1.92(0.02) same grp; other 12Qu01=1850(110)(+190-170)							
* <sup>90</sup> Nb <sup>r</sup>	T: average 05Ch65=470(10) 81Fi02=440(20) 78Ha52=477(8)							
* <sup>90</sup> Rh	T: symmetrized from 12(+9-4)							
* <sup>90</sup> Rh <sup>m</sup>	T: symmetrized from 1.0(+0.3-0.2)							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
<sup>91</sup> As	-36900#	400#	50# ms (>300 ns)	5/2 <sup>-</sup> #	13	97Be70 I	1997	$\beta^-$ ?; $\beta^-n=90\#$ ; $\beta^-2n=3\#$			
<sup>91</sup> Se	-50580	430	270 ms	50	1/2 <sup>+</sup> #	13	1975	$\beta^-n=100$ ; $\beta^-n=21$ 10; $\beta^-2n=0.03\#$			
<sup>91</sup> Br	-61107	4	543 ms	4	5/2 <sup>-</sup> #	13	1974	$\beta^-n=100$ ; $\beta^-n=19.5$ 26			
<sup>91</sup> Kr	-70974.0	2.2	8.57 s	0.04	5/2 <sup>(+)</sup>	13	1951	$\beta^-n=100$ ; $\beta^-n=0\#$			
<sup>91</sup> Rb	-77745	8	58.2 s	0.3	3/2 <sup>(-)</sup>	13	1951	$\beta^-n=100$ ; $\beta^-n=0\#$			
<sup>91</sup> Sr	-83652	5	9.65 h	0.06	5/2 <sup>+</sup>	13	1943	$\beta^-n=100$			
<sup>91</sup> Y	-86351.3	1.8	58.51 d	0.06	1/2 <sup>-</sup>	13	1943	$\beta^-n=100$			
<sup>91</sup> Y <sup>m</sup>	-85795.7	1.8	555.58	0.05	49.71 m	0.04	1953	IT>98.5; $\beta^-n<1.5$			
<sup>91</sup> Zr	-87895.57	0.10	STABLE		5/2 <sup>+</sup>	13	1934	IS=11.22 5			
<sup>91</sup> Zr <sup>m</sup>	-84728.3	0.4	3167.3	0.4	4.35 $\mu$ s	0.14	1985	IT=100			
<sup>91</sup> Nb	-86638.0	2.9	680 y	130	9/2 <sup>+</sup>	13	1951	$\epsilon\approx 100$ ; $e^+=0.0138$ 25			
<sup>91</sup> Nb <sup>m</sup>	-86533.4	2.9	104.60	0.05	60.86 d	0.22	1950	IT=96.6 5; $\epsilon=3.4$ 5; ; $e^+=0.0028$ 2			
<sup>91</sup> Nb <sup>n</sup>	-84603.6	2.9	2034.42	0.20	3.76 $\mu$ s	0.12	1974	IT=100			
<sup>91</sup> Mo	-82209	6	15.49 m	0.01	9/2 <sup>+</sup>	13	1948	$\beta^+n=100$			
<sup>91</sup> Mo <sup>m</sup>	-81556	6	653.01	0.09	64.6 s	0.6	1953	IT=50.0 16; $\beta^+n=50.0$ 16			
<sup>91</sup> Tc	-75986.6	2.4	3.14 m	0.02	(9/2 <sup>+</sup> )	13	1974	$\beta^+n=100$			
<sup>91</sup> Tc <sup>m</sup>	-75847.3	2.4	139.3	0.3	3.3 m	0.1	1975	$\beta^+n>99$ ; IT<1			
<sup>91</sup> Ru	-68239.8	2.2	8.0 s	0.4	(9/2 <sup>+</sup> )	13	1983	$\beta^+n=100$ ; $\beta^+p$ ?			
<sup>91</sup> Ru <sup>m</sup>	-68580	500	-340	500	BD *	7.6 s	0.8	1983	$\beta^+n\approx 100$ ; $\beta^+p=?$ ; IT?		
<sup>91</sup> Rh	-58570#	300#	1.60 s	0.15	7/2 <sup>+</sup> #	13	1994	$\beta^+n=100$ ; $\beta^+p=1.3$ 5			
<sup>91</sup> Rh <sup>m</sup>	-58400#	300#	172.9	0.4	1.46 s	0.11	2004	$\beta^+n=100$ ; IT?			
<sup>91</sup> Pd	-45930#	400#	10# ms (>1.5 $\mu$ s)	7/2 <sup>+</sup> #	13	95Ry03 I	1995	$\beta^+n$ ?; $\beta^+p$ ?			
* <sup>91</sup> Rh	T : average 04De40=1.7(0.2) 01Ki13=1.47(0.22); 00We.A(same group)=1.74(0.14)										
<sup>92</sup> As	-30980#	500#	30# ms (>300 ns)			12	97Be70 I	1997	$\beta^-$ ?; $\beta^-n=60\#$ ; $\beta^-2n=40\#$		
<sup>92</sup> Se	-46720#	400#	100# ms (>300 ns)	0 <sup>+</sup>		12	97Be70 I	1997	$\beta^-$ ?; $\beta^-n=2\#$ ; $\beta^-2n=0\#$		
<sup>92</sup> Se <sup>m</sup>	-44780#	400#	1940	50	12 $\mu$ s	4	12Ka36 ET	2012	IT=100		
<sup>92</sup> Br	-56233	7	0.314s	0.016	(2 <sup>-</sup> )	12	1974	$\beta^-n=100$ ; $\beta^-n=33.1$ 25; $\beta^-2n=0.01\#$			
<sup>92</sup> Br <sup>m</sup>	-55571	7	88 ns	8		12Ka36 ET	2012	IT=100			
<sup>92</sup> Br <sup>n</sup>	-55095	7	1138	1	85 ns	10	12Ka36 ET	2012	IT=100		
<sup>92</sup> Kr	-68769.3	2.7	1.840 s	0.008	0 <sup>+</sup>	12	1951	$\beta^-n=100$ ; $\beta^-n=0.0332$ 25			
<sup>92</sup> Rb	-74772	6	4.48 s	0.03	0 <sup>-</sup>	12	1960	$\beta^-n=100$ ; $\beta^-n=0.0107$ 5			
<sup>92</sup> Sr	-82867	3	2.611 h	0.017	0 <sup>+</sup>	12	1956	$\beta^-n=100$			
<sup>92</sup> Y	-84816	9	3.54 h	0.01	2 <sup>-</sup>	12	1940	$\beta^-n=100$			
<sup>92</sup> Y <sup>m</sup>	-84010	50	807	50	3.7 $\mu$ s	0.5	7 <sup>+</sup> #	12	11Ru.A ET	2009	IT=100
<sup>92</sup> Zr	-88459.03	0.10	STABLE		0 <sup>+</sup>	12	1924	IS=17.15 8			
<sup>92</sup> Nb	-86453.3	1.8	34.7 My	2.4	7 <sup>+</sup>	12	1938	$\beta^+n=100$			
<sup>92</sup> Nb <sup>m</sup>	-86317.8	1.8	135.5	0.4	10.15 d	0.02	(2 <sup>+</sup> )	12	1959	$\beta^+n=100$	
<sup>92</sup> Nb <sup>n</sup>	-86227.5	1.8	225.8	0.4	5.9 $\mu$ s	0.2	(2 <sup>-</sup> )	12	1958	IT=100	
<sup>92</sup> Nb <sup>p</sup>	-84250.0	1.8	2203.3	0.4	167 ns	4	(11 <sup>-</sup> )	12	1989	IT=100	
<sup>92</sup> Mo	-86808.58	0.16	STABLE	(>190 Ey)	0 <sup>+</sup>	12	97Ba35 T	1930	IS=14.53 30; $2\beta^+$ ?		
<sup>92</sup> Mo <sup>m</sup>	-84048.06	0.21	2760.52	0.14	190 ns	3	8 <sup>+</sup>	12	1964	IT=100	
<sup>92</sup> Tc	-78926	3	4.25 m	0.15	(8 <sup>+</sup> )	12	1964	$\beta^+n=100$			
<sup>92</sup> Tc <sup>m</sup>	-78656	3	270.09	0.08	1.03 $\mu$ s	0.07	(4 <sup>+</sup> )	12	1976	IT=100	
<sup>92</sup> Tc <sup>n</sup>	-78397	3	529.42	0.13	< 0.1 $\mu$ s		(3 <sup>+</sup> )	12	1976	IT=100	
<sup>92</sup> Tc <sup>p</sup>	-78215	3	711.33	0.15	< 0.1 $\mu$ s		1 <sup>+</sup>	12	1976	IT=100	
<sup>92</sup> Ru	-74301.2	2.7	3.65 m	0.05	0 <sup>+</sup>	12	1971	$\beta^+n=100$			
<sup>92</sup> Rh	-62999	4	4.66 s	0.25	(6 <sup>+</sup> )	12	04De40 J	1994	$\beta^+n=100$ ; $\beta^+p=1.9$ 1		
<sup>92</sup> Rh <sup>m</sup>	-62950#	100#	50#	100#	0.53 s	0.37	(2 <sup>+</sup> )	12	04De40 TJD	2004	$\beta^+n=100$ ; $\beta^+p=?$
<sup>92</sup> Pd	-54580#	300#	1.1 s	0.3	0 <sup>+</sup>	12	01Ki13 TD	1994	$\beta^+n=100$ ; $\beta^+p$ ?		
<sup>92</sup> Ag	-37130#	500#	1# ms (>400 ns)			16Ce02 I	2016	$\beta^+n$ ?			
* <sup>92</sup> Se <sup>m</sup>	E : 12Ka36=503.4(0.5), 538.8(0.5) and 897.8(0.5) $\gamma$ rays in cascade to										
* <sup>92</sup> Se <sup>m</sup>	E : ground-state =1940(1); error increased for possible missing $\gamma$										
* <sup>92</sup> Se <sup>m</sup>	T : symmetrized from 10.3(+5.5-2.8)										
* <sup>92</sup> Br	I : also an isomer with T<500 ns decaying by $\gamma$ -rays 1039, 780, 301... keV										
* <sup>92</sup> Br <sup>m</sup>	T : symmetrized from 89(+7-8)										
* <sup>92</sup> Br <sup>n</sup>	T : symmetrized from 84(+10-9); other 09Fo05<500 ns assuming single isomer										
* <sup>92</sup> Y <sup>m</sup>	T : average 11Ru.A=3.3(0.6) 09Fo05=4.2(+0.8-0.6)										
* <sup>92</sup> Y <sup>m</sup>	E : observed 315 and 419 $\gamma$ rays; low energy transition may directly										
* <sup>92</sup> Y <sup>m</sup>	E : depopulate the isomer										
* <sup>92</sup> Mo	T : T>190 Ey (2 $\sigma$ )										
* <sup>92</sup> Rh	D : from 12Lo08										
* <sup>92</sup> Rh <sup>m</sup>	I : this state is not observed in 12Lo08										
* <sup>92</sup> Pd	T : symmetrized from 1.0(+0.3-0.2)										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>93</sup> Se	-40720# 400#		50# ms (>300 ns)	1/2 <sup>+</sup> #	11	97Be70 I	1997	$\beta^-$ ?; $\beta^-n=30\#$ ; $\beta^-2n=2\#$
<sup>93</sup> Se <sup>m</sup>	-40040# 400#	678.2 0.7	420 ns 100			12Ka36 ET	2012	IT=100
<sup>93</sup> Br	-52890 430		152 ms 8	5/2 <sup>-</sup> #	11	13Mi13 TD	1981	$\beta^-n=100$ ; $\beta^-n=55$ 10; $\beta^-2n=0.01\#$
<sup>93</sup> Kr	-64136.0 2.5		1.286 s 0.010	1/2 <sup>+</sup>	11		1951	$\beta^-n=100$ ; $\beta^-n=1.95$ 11
<sup>93</sup> Rb	-72620 8		5.84 s 0.02	5/2 <sup>-</sup>	11		1960	$\beta^-n=100$ ; $\beta^-n=1.39$ 7
<sup>93</sup> Rb <sup>m</sup>	-68197 8	4423.1 1.5	111 ns 11	(27/2 <sup>-</sup> )	11		2010	IT=100
<sup>93</sup> Rb <sup>t</sup>	-72367 8	253.39 0.03	< 0.5 ns	3/2 <sup>-</sup>	11	86Si20 T	1970	IT=100
<sup>93</sup> Sr	-80086 8		7.43 m 0.03	5/2 <sup>+</sup>	11		1959	$\beta^-n=100$
<sup>93</sup> Y	-84227 10		10.18 h 0.08	1/2 <sup>-</sup>	11		1948	$\beta^-n=100$
<sup>93</sup> Y <sup>m</sup>	-83468 10	758.719 0.021	820 ms 40	9/2 <sup>+</sup>	11	07Ch07 J	1974	IT=100
<sup>93</sup> Zr	-87122.0 0.5		1.61 My 0.05	5/2 <sup>+</sup>	11		1950	$\beta^-n=100$
<sup>93</sup> Nb	-87212.8 1.5		STABLE	9/2 <sup>+</sup>	11		1932	IS=100.
<sup>93</sup> Nb <sup>m</sup>	-87182.0 1.5	30.77 0.02	16.12 y 0.12	1/2 <sup>-</sup>	11		1965	IT=100
<sup>93</sup> Nb <sup>n</sup>	-79753 17	7460 17	1.5 $\mu$ s 0.5		11		2007	IT ?
<sup>93</sup> Mo	-86807.07 0.18		4.0 ky 0.8	5/2 <sup>+</sup>	11		1946	$\epsilon=100$
<sup>93</sup> Mo <sup>m</sup>	-84382.12 0.18	2424.95 0.04	6.85 h 0.07	21/2 <sup>+</sup>	11		1950	IT $\approx$ 100; $\beta^+=0.12$ 1
<sup>93</sup> Mo <sup>n</sup>	-77112 17	9695 17	1.8 $\mu$ s 1.0	(39/2 <sup>-</sup> )	11		2005	IT ?
<sup>93</sup> Tc	-83606.1 1.0		2.75 h 0.05	9/2 <sup>+</sup>	11		1948	$\beta^+=100$
<sup>93</sup> Tc <sup>m</sup>	-83214.3 1.0	391.84 0.08	43.5 m 1.0	1/2 <sup>-</sup>	11		1939	IT=77.4 6; $\beta^+=22.6$ 6
<sup>93</sup> Tc <sup>n</sup>	-81420.9 1.0	2185.16 0.15	10.2 $\mu$ s 0.3	(17/2 <sup>-</sup> )	11		1973	IT=100
<sup>93</sup> Ru	-77216.7 2.1		59.7 s 0.6	(9/2 <sup>+</sup> )	11		1972	$\beta^+=100$
<sup>93</sup> Ru <sup>m</sup>	-76482.3 2.1	734.40 0.10	10.8 s 0.3	(1/2 <sup>-</sup> )	11		1983	$\beta^+=78.0$ 23; IT=22.0 23; $\beta^+p=0.027$ 5
<sup>93</sup> Ru <sup>n</sup>	-75134.2 2.3	2082.5 0.9	2.49 $\mu$ s 0.15	(21/2 <sup>+</sup> )	11		1983	IT=100
<sup>93</sup> Rh	-69011.8 2.6		13.9 s 1.6	9/2 <sup>+</sup> #	11		1994	$\beta^+=100$
<sup>93</sup> Pd	-59000# 300#		1.15 s 0.05	(9/2 <sup>+</sup> )	11	12Lo08 TD	1994	$\beta^+=100$ ; $\beta^+p=7.5$ 5
<sup>93</sup> Ag	-46270# 400#		20# ms (>1.5 $\mu$ s)	9/2 <sup>+</sup> #	11	95Ry03 I	1994	$\beta^+ ?$ ; $p ?$ ; $\beta^+p ?$
<sup>93</sup> Se <sup>m</sup>	E : 12Ka36=208.3(0.5) and 469.9(0.5) $\gamma$ rays in cascade to ground-state							
<sup>93</sup> Se <sup>m</sup>	T : symmetrized from 390(+120-80)							
<sup>93</sup> Br	D : symmetrized from 13Mi13 $\beta^-n=53(+11-8)\%$							
<sup>93</sup> Kr	T : also 13Mi13=1.298(0.054) outweighed D : also 13Mi13=1.9(+0.6-0.2)							
<sup>93</sup> Rb <sup>t</sup>	T : 70Gr38=57(15) $\mu$ s not confirmed in 14Mi12; most likely <sup>95</sup> Y <sup>m</sup>							
<sup>93</sup> Rb <sup>t</sup>	J : 253.4 keV M1 (and E2) $\gamma$ ray to 5/2 <sup>-</sup> ; $\beta^-$ feeding from 1/2 <sup>+</sup> <sup>93</sup> Kr							
<sup>93</sup> Nb <sup>n</sup>	E : ENSDF2011 : x keV above 7435.3(2.1) 37/2 <sup>-</sup> level; NUBASE assumes x<50							
<sup>93</sup> Mo <sup>n</sup>	E : ENSDF2011 : x keV above 9670.0(2.3) (35/2,37/2) level; NUBASE assumes x<50							
<sup>93</sup> Mo <sup>n</sup>	T : symmetrized from 1.1(+1.5-0.4)							
<sup>93</sup> Ag	I : the few events reported in <sup>94</sup> He28 are not trusted by NUBASE							
<sup>93</sup> Ag	I : 10St.A>0.2 $\mu$ s							
<sup>93</sup> Ag	T : estimated half-life is for $\beta^+$ decay; p-decay would be much shorter							
<sup>93</sup> Ag	T : post-deadline 16Ce02=228#(16#) ns							
<sup>94</sup> Se	-36800# 500#		20# ms (>300 ns)	0 <sup>+</sup>	06	97Be70 I	1997	$\beta^-$ ?; $\beta^-n=20\#$ ; $\beta^-2n=0.2\#$
<sup>94</sup> Br	-47400# 300#		70 ms 20	2 <sup>-</sup> #	06		1981	$\beta^-n=100$ ; $\beta^-n=68$ 16; $\beta^-2n=3\#$
<sup>94</sup> Br <sup>m</sup>	-47110# 300#	294.6 0.5	530 ns 15			12Ka36 ET	2012	IT=100
<sup>94</sup> Kr	-61348 12		212 ms 5	0 <sup>+</sup>	11		1972	$\beta^-n=100$ ; $\beta^-n=1.11$ 7
<sup>94</sup> Rb	-68562.8 2.0		2.702 s 0.005	3 <sup>(-)</sup>	11		1961	$\beta^-n=100$ ; $\beta^-n=10.5$ 4
<sup>94</sup> Rb <sup>m</sup>	-66487.9 2.4	2074.9 1.4	107 ns 16	(10 <sup>-</sup> )	11		2008	IT=100
<sup>94</sup> Sr	-78845.7 1.7		75.3 s 0.2	0 <sup>+</sup>	11		1959	$\beta^-n=100$
<sup>94</sup> Y	-82351 6		18.7 m 0.1	2 <sup>-</sup>	06		1948	$\beta^-n=100$
<sup>94</sup> Y <sup>m</sup>	-81149 6	1202.3 1.0	1.295 $\mu$ s 0.005	(5 <sup>+</sup> )	06	11Ru.A T	1999	IT=100
<sup>94</sup> Zr	-87269.32 0.16		STABLE	(>110 Py)	0 <sup>+</sup>	06 99Ar25 T	1924	IS=17.38 28; 2 $\beta^-$ ?
<sup>94</sup> Nb	-86369.1 1.5		20.4 ky 0.4	6 <sup>+</sup>	06	12He11 T	1938	$\beta^-n=100$
<sup>94</sup> Nb <sup>m</sup>	-86328.2 1.5	40.892 0.012	6.263 m 0.004	3 <sup>+</sup>	06		1962	IT=99.50 6; $\beta^-n=0.50$ 6
<sup>94</sup> Mo	-88414.06 0.14		STABLE	0 <sup>+</sup>	06		1930	IS=9.15 9
<sup>94</sup> Tc	-84158 4		293 m 1	7 <sup>+</sup>	06		1948	$\beta^+=100$
<sup>94</sup> Tc <sup>m</sup>	-84082 5	76 3	52.0 m 1.0	(2) <sup>+</sup>	06		1948	$\beta^+\approx 100$ ; IT<0.1
<sup>94</sup> Ru	-82584 3		51.8 m 0.6	0 <sup>+</sup>	06		1952	$\beta^+=100$
<sup>94</sup> Ru <sup>m</sup>	-79940 3	2644.1 0.4	71 $\mu$ s 4	8 <sup>+</sup>	06		1971	IT=100
<sup>94</sup> Rh	-72908 3		70.6 s 0.6	(4 <sup>+</sup> )	06	06Ba55 J	1979	$\beta^+=100$ ; $\beta^+p=1.8$ 5
<sup>94</sup> Rh <sup>m</sup>	-72853 3	54.60 0.20	480 ns 30	(2 <sup>+</sup> )	06		2004	IT=100
<sup>94</sup> Rh <sup>n</sup>	-72610# 200#	300# 200#	25.8 s 0.2	(8 <sup>+</sup> )	06		1973	$\beta^+=100$
<sup>94</sup> Pd	-66102 4		9.0 s 0.5	0 <sup>+</sup>	06		1982	$\beta^+=100$
<sup>94</sup> Pd <sup>m</sup>	-61219 4	4883.1 0.4	511.0 ns 7.3	(14 <sup>+</sup> )	06	11Br01 T	1995	IT=100
<sup>94</sup> Pd <sup>n</sup>	-58893 4	7209.1 1.8	197 ns 22	(19 <sup>-</sup> )	06	11Br01 TJ	2011	IT=100

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)				
... A-group continued ...													
<sup>94</sup> Ag	-52410#	400#		37 ms	18	0 <sup>+</sup> #	06	1994	$\beta^+=100; \beta^+p?$	*			
<sup>94</sup> Ag <sup>m</sup>	-51060#	570#	1350#	400#	550 ms	60	(7 <sup>+</sup> )	06	$\beta^+=100; \beta^+p=20$	**			
<sup>94</sup> Ag <sup>n</sup>	-45920#	300#	6490#	500#	400 ms	40	(21 <sup>+</sup> )	06	$\beta^+=95.4\ 7; \beta^+p=27; p=4.1\ 6; 2p=0.5\ 3$	*			
<sup>94</sup> Cd	-40140#	500#		80# ms	(>400 ns)	0 <sup>+</sup>	16Ce02	I	2016	$\beta^+?$	**		
<sup>94</sup> Pd <sup>m</sup>	T: average 11Br01=499(13) 09Ga40=468(19) 02La18=530(10)												
<sup>94</sup> Pd <sup>n</sup>	E: from 4883.1(0.4) for the 14 <sup>+</sup> state and 1651(1), 267(1) and 408(1) keV												
<sup>94</sup> Pd <sup>n</sup>	E: $\gamma$ rays in a cascade from (19 <sup>-</sup> ); uncertainties added in quadrature												
<sup>94</sup> Ag	T: symmetrized from 26(+26-9)												
<sup>94</sup> Ag <sup>n</sup>	D: p=1.9(5) + 2.2(4) from 05Mu15, 2p from 06Mu03												
<sup>95</sup> Se	-30460#	500#		10# ms	(>400 ns)	3/2 <sup>+</sup> #	12	100h02	I	2010	$\beta^-?; \beta^-n=10\#; \beta^-2n=3\#$	**	
<sup>95</sup> Br	-43770#	300#		50# ms	(>300 ns)	5/2 <sup>-</sup> #	10	97Be70	I	1997	$\beta^-?; \beta^-n=70\#; \beta^-2n=0.2\#$	**	
<sup>95</sup> Br <sup>m</sup>	-43230#	300#	537.9	0.5	6.8 $\mu$ s	1.0		12Ka36	ET	2012	IT=100	*	
<sup>95</sup> Kr	-56159	19			114 ms	3	1/2 <sup>(+)</sup>	10		1994	$\beta^-n=100; \beta^-n=2.87\ 18; \beta^-2n=0\#$	**	
<sup>95</sup> Kr <sup>m</sup>	-55964	19	195.5	0.3	1.582 $\mu$ s	0.022	(7/2 <sup>+</sup> )	10	12Ka36	T	2006	IT=100	*
<sup>95</sup> Rb	-65891	20			377.7 ms	0.8	5/2 <sup>-</sup>	10		1967	$\beta^-n=100; \beta^-n=8.7\ 3$	**	
<sup>95</sup> Rb <sup>m</sup>	-65056	20	835.0	0.6	< 500 ns		9/2 <sup>+</sup> #	10		2009	IT=100	**	
<sup>95</sup> Sr	-75120	6			23.90 s	0.14	1/2 <sup>+</sup>	10		1961	$\beta^-n=100$	**	
<sup>95</sup> Y	-81209	7			10.3 m	0.1	1/2 <sup>-</sup>	10		1959	$\beta^-n=100$	**	
<sup>95</sup> Y <sup>m</sup>	-80121	7	1087.6	0.6	48.6 $\mu$ s	0.5	9/2 <sup>+</sup>	10	11Ru.A	T	1981	IT=100	**
<sup>95</sup> Zr	-85659.9	0.9			64.032 d	0.006	5/2 <sup>+</sup>	10		1946	$\beta^-n=100$	**	
<sup>95</sup> Nb	-86786.3	0.5			34.991 d	0.006	9/2 <sup>+</sup>	10		1951	$\beta^-n=100$	**	
<sup>95</sup> Nb <sup>m</sup>	-86550.6	0.5	235.69	0.02	3.61 d	0.03	1/2 <sup>-</sup>	10		1969	IT=94.4 6; $\beta^-n=5.6\ 6$	**	
<sup>95</sup> Mo	-87711.86	0.12			STABLE		5/2 <sup>+</sup>	10		1930	IS=15.84 11	**	
<sup>95</sup> Tc	-86021	5			20.0 h	0.1	9/2 <sup>+</sup>	10		1947	$\beta^+n=100$	**	
<sup>95</sup> Tc <sup>m</sup>	-85982	5	38.91	0.04	61 d	2	1/2 <sup>-</sup>	10		1959	$\beta^+n=96.12\ 32; IT=3.88\ 32$	**	
<sup>95</sup> Ru	-83458	10			1.643 h	0.013	5/2 <sup>+</sup>	10		1948	$\beta^+n=100$	**	
<sup>95</sup> Rh	-78341	4			5.02 m	0.10	(9/2 <sup>+</sup> )	10		1967	$\beta^+n=100$	**	
<sup>95</sup> Rh <sup>m</sup>	-77798	4	543.3	0.3	1.96 m	0.04	(1/2 <sup>-</sup> )	10		1974	IT=88 5; $\beta^+n=12\ 5$	**	
<sup>95</sup> Pd	-69966	3			7.5 s	0.5	9/2 <sup>+</sup> #	10	12Lo08	T	1980	$\beta^+n=100; \beta^+p?$	**
<sup>95</sup> Pd <sup>m</sup>	-68091	3	1875.13	0.14	13.3 s	0.3	(21/2 <sup>+</sup> )	10		1982	$\beta^+n=?; IT=11\ 3; \beta^+p=0.93\ 15$	**	
<sup>95</sup> Ag	-59600#	300#			1.76 s	0.09	(9/2 <sup>+</sup> )	10	12Lo08	TD	1994	$\beta^+n=100; \beta^+p=2.5\ 3$	*
<sup>95</sup> Ag <sup>m</sup>	-59260#	300#	344.2	0.3	< 500 ms		(1/2 <sup>-</sup> )	10		2003	IT=100	**	
<sup>95</sup> Ag <sup>n</sup>	-57070#	300#	2531.3	1.5	< 16 ms		(23/2 <sup>+</sup> )	10		2003	IT=100	**	
<sup>95</sup> Ag <sup>p</sup>	-54740#	300#	4860.0	1.5	< 40 ms		(37/2 <sup>+</sup> )	10		2003	IT=100	**	
<sup>95</sup> Cd	-46630#	400#			90 ms	40	9/2 <sup>+</sup> #	10	10St.A	T	2011	$\beta^+?; \beta^+p?$	*
<sup>95</sup> Br <sup>m</sup>	T: symmetrized from 6.7(+1.1-0.9)												
<sup>95</sup> Kr <sup>m</sup>	E: also 12Ka36=82.6(0.5) and 114.3(0.5) $\gamma$ rays in cascade to ground-state												
<sup>95</sup> Kr <sup>m</sup>	T: other 11Ru.A=1.28(0.05) 06Ge05=1.4(0.2)												
<sup>95</sup> Ag	T: average 12Lo08=1.85(0.08) 05Ha45=1.76(0.13) 03Do09=1.85(0.34) and												
<sup>95</sup> Cd	T: symmetrized from 73(+53-28)												
<sup>96</sup> Br	-38160#	300#			20# ms	(>300 ns)		08	97Be70	I	1997	$\beta^-?; \beta^-n=50\#; \beta^-2n=6\#$	**
<sup>96</sup> Br <sup>m</sup>	-37850#	300#	311.5	0.5	3.0 $\mu$ s	0.9			12Ka36	ET	2012	IT=100	*
<sup>96</sup> Kr	-53080	20			80 ms	8	0 <sup>+</sup>	12		1994	$\beta^-n=100; \beta^-n=3.7\ 4$	**	
<sup>96</sup> Rb	-61354	3			201 ms	1	2 <sup>-</sup>	08	93Ru01	T	1967	$\beta^-n=100; \beta^-n=13.3\ 7; \beta^-2n=0.02\#$	*
<sup>96</sup> Rb <sup>m</sup>	-61350#	200#	0#	200#	200# ms	(>1 ms)	1 <sup>(+)</sup> #	08	81Bo30	JI	1981	$\beta^-?; IT?; \beta^-n=10\#; \beta^-2n=0.02\#$	*
<sup>96</sup> Rb <sup>n</sup>	-60219	3	1134.6	1.1	1.80 $\mu$ s	0.04	(10 <sup>-</sup> )	08		1999	IT=100	*	
<sup>96</sup> Sr	-72924	8			1.07 s	0.01	0 <sup>+</sup>	08		1971	$\beta^-n=100; \beta^-n=0\#$	**	
<sup>96</sup> Y	-78336	6			5.34 s	0.05	0 <sup>-</sup>	08		1975	$\beta^-n=100$	**	
<sup>96</sup> Y <sup>m</sup>	-76796	6	1540	9	9.6 s	0.2	8 <sup>+</sup>	08	07Ch07	J	1974	$\beta^-n=100$	**
<sup>96</sup> Zr	-85438.85	0.11			23 Ey	2	0 <sup>+</sup>	08	15Ba11	T	1934	IS=2.80 9; $2\beta^-n=100$	**
<sup>96</sup> Nb	-85602.82	0.15			23.35 h	0.05	6 <sup>+</sup>	08		1949	$\beta^-n=100$	**	
<sup>96</sup> Mo	-88794.88	0.12			STABLE		0 <sup>+</sup>	08		1930	IS=16.67 15	**	
<sup>96</sup> Tc	-85822	5			4.28 d	0.07	7 <sup>+</sup>	08		1947	$\beta^+n=100$	**	
<sup>96</sup> Tc <sup>m</sup>	-85788	5	34.23	0.04	51.5 m	1.0	4 <sup>+</sup>	08		1950	IT=98.0 5; $\beta^+n=2.0\ 5$	**	
<sup>96</sup> Ru	-86080.37	0.17			STABLE	(>80 Ey)	0 <sup>+</sup>	08	13Be09	T	1931	IS=5.54 14; $2\beta^+?$	*
<sup>96</sup> Rh	-79688	10			9.90 m	0.10	6 <sup>+</sup>	08		1967	$\beta^+n=100$	**	
<sup>96</sup> Rh <sup>m</sup>	-79636	10	51.98	0.09	1.51 m	0.02	3 <sup>+</sup>	08		1966	IT=60 5; $\beta^+n=40\ 5$	**	
<sup>96</sup> Pd	-76183	4			122 s	2	0 <sup>+</sup>	08		1980	$\beta^+n=100$	**	
<sup>96</sup> Pd <sup>m</sup>	-73652	4	2530.57	0.23	1.81 $\mu$ s	0.01	8 <sup>+</sup> #	08	98Gr.B	TD	1983	IT=100	*
<sup>96</sup> Ag	-64510	90			4.44 s	0.04	(8 <sup>+</sup> )	08	12Lo08	TD	1982	$\beta^+n=100; \beta^+p=6.9\ 7$	*
<sup>96</sup> Ag <sup>m</sup>	-64510#	100#	0#	50#	6.9 s	0.5	(2 <sup>-</sup> )	08	12Lo08	TD	2003	$\beta^+n=100; \beta^+p=15.1\ 26$	*
<sup>96</sup> Ag <sup>n</sup>	-62050	90	2461.4	0.3	100 $\mu$ s	10	(13 <sup>-</sup> )	08	11Bo23	TJD	2011	IT=100	*
<sup>96</sup> Ag <sup>p</sup>	-61830	90	2680	7	1.543 $\mu$ s	0.028	(15 <sup>+</sup> )	08	11Bo23	ETJ	2011	IT=100	*
<sup>96</sup> Ag <sup>d</sup>	-57570	90	6945	7	160 ns	30	(19 <sup>+</sup> )	08	11Bo23	ETJ	2011	IT=100	*
... A-group is continued on next page ...													

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...										
<sup>96</sup> Cd	-55570#	400#	880 ms	90	0 <sup>+</sup>	10 12Lo08 D	2008	$\beta^+=100; \beta^+p=5.5$ 40	*	
<sup>96</sup> Cd <sup>m</sup>	-50270#	400# 5300# 2000#	300 ms	110	16 <sup>+</sup>	10 11Na34 TJD	2011	$\beta^+=100; \beta^+p?$	*	
<sup>96</sup> In	-37890#	500#	1# ms	(>400 ns)		16Ce02 I	2016	$\beta^+?$	**	
<sup>96</sup> Br <sup>m</sup>	T: symmetrized from 2.7(+1.1-0.7)									
<sup>96</sup> Rb	J: measured magnetic moment consistent with 2 <sup>-</sup>									
<sup>96</sup> Rb <sup>m</sup>	I: non-observation in 81Th04 is not in contradiction with 81Bo30 experiment									
<sup>96</sup> Rb <sup>n</sup>	T: average 12Ka36=1.72(+0.16-0.14) 11Ru.A=1.77(0.05) 05Pi13=2.0(0.1)									
<sup>96</sup> Rb <sup>n</sup>	T: <sup>99</sup> Ge01=1.65(0.15)									
<sup>96</sup> Ru	T: 13Be09: 2v- $\beta^+e>80$ Ey (theor. most probable); 2nu $\beta^+>140$ Ey 0nu2K>1Zy									
<sup>96</sup> Pd <sup>m</sup>	T: supersedes 97Gr02=1.7(0.1); others 09Ga40=1.76(0.05) 07My02=2.10(0.21)									
<sup>96</sup> Pd <sup>m</sup>	T: <sup>83</sup> Gr01=2.2(0.3) J: from 03Ba39									
<sup>96</sup> Ag	T: average 12Lo08=4.40(0.09) 03Ba39=4.40(0.06) 97Sc30=4.50(0.06)									
<sup>96</sup> Ag	D: $\beta^+p$ average 12Lo08=6.5(0.8) 03Ba39=8.5(1.5)									
<sup>96</sup> Ag <sup>m</sup>	T: average 12Lo08=6.8(1.0) 03Ba39=6.9(0.6) D: average 12Lo08=14(3) 03Ba39=18(5)									
<sup>96</sup> Ag <sup>n</sup>	E: from least-squares fit to $\gamma$ -ray energies using 11Bo23 level scheme									
<sup>96</sup> Ag <sup>n</sup>	T: other 11Be34=8.6(6.3) $\mu$ s using a collection time of 12 $\mu$ s									
<sup>96</sup> Ag <sup>p</sup>	E: 25-50 keV above the 2643 13 <sup>+</sup> level									
<sup>96</sup> Ag <sup>p</sup>	T: average 11Bo23=1.56(0.03) 11Be34=1.45(0.07)									
<sup>96</sup> Ag <sup>q</sup>	E: 4265 above the <sup>96</sup> Ag <sup>p</sup>									
<sup>96</sup> Cd	T: average 11Na34=670(150) 10St.A=990(130) 08Ba53=1030(+240-210)									
<sup>96</sup> Cd <sup>m</sup>	T: symmetrized from 11Na34=290(+110-100)									
<sup>97</sup> Br	-34060#	400#	10# ms	(>300 ns)	5/2 <sup>-</sup> #	10	1997	$\beta^-?$ ; $\beta^-n=90\#; \beta^-2n=5\#$	*	
<sup>97</sup> Kr	-47420	130	62.2 ms	3.2	3/2 <sup>+</sup> #	10 11Ni01 T	1997	$\beta^-n=100; \beta^-n=6.7$ 6; $\beta^-2n=0.1\#$	*	
<sup>97</sup> Rb	-58519.1	1.9	169.1 ms	0.6	3/2 <sup>+</sup>	15	1969	$\beta^-n=100; \beta^-n=25.5$ 9; $\beta^-2n=0\#$	*	
<sup>97</sup> Rb <sup>m</sup>	-58442.5	1.9	76.6	0.2	5.7 $\mu$ s	0.6 (1/2,3/2) <sup>-</sup>	15	2012	IT=100	
<sup>97</sup> Sr	-68581	3	429 ms	5	1/2 <sup>+</sup>	10	1978	$\beta^-n=100; \beta^-n<0.05$	*	
<sup>97</sup> Sr <sup>m</sup>	-68273	3	308.13	0.11	165 ns	4 7/2 <sup>+</sup>	10 15Cz01 T	1990	IT=100	
<sup>97</sup> Sr <sup>n</sup>	-67750	3	830.83	0.23	515 ns	10 (9/2 <sup>+</sup> )	10 13Ru07 TJ	1974	IT=100	
<sup>97</sup> Y	-76121	7	3.75 s	0.03	1/2 <sup>-</sup>	10 07Ch07 J	1970	$\beta^-n=100; \beta^-n=0.055$ 4	*	
<sup>97</sup> Y <sup>m</sup>	-75453	7	667.52	0.23	1.17 s	0.03 9/2 <sup>+</sup>	10 07Ch07 J	1970	$\beta^->99.3; IT<0.7; \beta^-n<0.08$	
<sup>97</sup> Y <sup>n</sup>	-72598	7	3522.6	0.4	142 ms	8 (27/2 <sup>-</sup> )	10	1986	IT=94.8 9; $\beta^-n=5.2$ 9	
<sup>97</sup> Zr	-82942.7	0.4	16.749 h	0.008	1/2 <sup>+</sup>	10	1951	$\beta^-n=100$	*	
<sup>97</sup> Zr <sup>m</sup>	-81678.3	0.4	1264.35	0.16	104.8 ns	1.7 7/2 <sup>+</sup>	10 11Ru.A T	1976	IT=100	
<sup>97</sup> Nb	-85606	4	72.1 m	0.7	9/2 <sup>+</sup>	10	1951	$\beta^-n=100$	*	
<sup>97</sup> Nb <sup>m</sup>	-84863	4	743.35	0.03	58.7 s	1.8 1/2 <sup>-</sup>	10	1950	IT=100	
<sup>97</sup> Mo	-87544.69	0.16	STABLE		5/2 <sup>+</sup>	10	1930	IS=9.60 14	*	
<sup>97</sup> Tc	-87224	4	4.21 My	0.16	9/2 <sup>+</sup>	10	1946	$\epsilon=100$	*	
<sup>97</sup> Tc <sup>m</sup>	-87127	4	96.57	0.06	91.0 d	0.6 1/2 <sup>-</sup>	10	1954	IT=96.06 18; $\epsilon=3.94$ 18	
<sup>97</sup> Ru	-86120.6	2.8	2.8370 d	0.0014	5/2 <sup>+</sup>	10 09Go29 T	1946	$\beta^+=100$	*	
<sup>97</sup> Rh	-82600	40	30.7 m	0.6	9/2 <sup>+</sup>	10	1955	$\beta^+=100$	*	
<sup>97</sup> Rh <sup>m</sup>	-82340	40	258.76	0.18	46.2 m	1.6 1/2 <sup>-</sup>	10	1971	$\beta^+=94.4$ 6; IT=5.6 6	
<sup>97</sup> Pd	-77806	5	3.10 m	0.09	5/2 <sup>+</sup> #	10	1969	$\beta^+=100$	*	
<sup>97</sup> Ag	-70830	110	25.5 s	0.3	(9/2 <sup>+</sup> ) <sup>+</sup>	10 14Fe01 J	1978	$\beta^+=100$	*	
<sup>97</sup> Ag <sup>m</sup>	-70430#	230#	400#	200#	100# ms	1/2 <sup>-</sup> #		IT?	*	
<sup>97</sup> Cd	-60450#	300#	1.10 s	0.08	(9/2 <sup>+</sup> )	10 11Lo09 TJD	1978	$\beta^+=100; \beta^+p=11.8$ 20	*	
<sup>97</sup> Cd <sup>m</sup>	-58950#	580#	1500#	500#	3.8 s	0.2 (25/2 <sup>+</sup> )	10 11Lo09 TJD	1982	$\beta^+=100; \beta^+p=25$ 4	
<sup>97</sup> In	-47190#	400#	50 ms	30	9/2 <sup>+</sup> #	10St.A TD	2011	$\beta^+=100; p?; \beta^+p?$	*	
<sup>97</sup> Kr	T: average 11Ni01=60(+6-5) 03Be05=63(4)									
<sup>97</sup> Sr <sup>m</sup>	E: also 12Ka36=141.3(0.5) and 167.6(0.5) $\gamma$ rays in cascade to ground-state									
<sup>97</sup> Sr <sup>m</sup>	T: others 11Ru.A=180.9(2.8) 06Hw01=165(25) 83Kr11=170(10)									
<sup>97</sup> Zr <sup>m</sup>	T: average 11Ru.A=106.1(2.1) 85Be20=102(3); others outweighed 06Hw01=97(16)									
<sup>97</sup> Zr <sup>m</sup>	T: <sup>96</sup> Lh03=106(7)									
<sup>97</sup> In	T: symmetrized from 26(+47-10)									
<sup>98</sup> Br	-28250#	400#	5# ms	(>400 ns)		10 10Oh02 I	2010	$\beta^-?$ ; $\beta^-n=70\#; \beta^-2n=20\#$	*	
<sup>98</sup> Kr	-44310#	300#	42.8 ms	3.6	0 <sup>+</sup>	03 11Ni01 T	1997	$\beta^-n=100; \beta^-n=7.0$ 10; $\beta^-2n=0\#$	*	
<sup>98</sup> Rb	-54369	16	114 ms	5	0 <sup>(-#)</sup>	03 81Th04 J	1971	$\beta^-n=100; \beta^-n=13.8$ 6; $\beta^-2n=0.051$ 7	*	
<sup>98</sup> Rb <sup>m</sup>	-54296	20	73	26	BD	96 ms	3 (3,4) <sup>(+#)</sup>	03	1980	$\beta^-n=100; \beta^-n=10\#; \beta^-2n=0.05\#$
<sup>98</sup> Rb <sup>n</sup>	-54191	16	178.3	0.4	358 ns	7	09 12Ka36 ET	2009	IT=100	
<sup>98</sup> Sr	-66423	3	653 ms	2	0 <sup>+</sup>	03	1971	$\beta^-n=100; \beta^-n=0.25$ 5	*	
<sup>98</sup> Y	-72295	8	548 ms	2	(0) <sup>-</sup>	03	1970	$\beta^-n=100; \beta^-n=0.331$ 24	*	
<sup>98</sup> Y <sup>m</sup>	-72054	28	241	29	BD	2.0 s	0.2 (5 <sup>+</sup> ,4 <sup>-</sup> )	03	1977	$\beta^-n=100; IT=10\#; \beta^-n=3.4$ 10
<sup>98</sup> Y <sup>n</sup>	-72124	8	170.74	0.06	610 ns	9 (2) <sup>-</sup>	03 11Ru.A T	1972	IT=100	
<sup>98</sup> Y <sup>p</sup>	-71799	8	496.19	0.15	6.87 $\mu$ s	0.05 (4) <sup>-</sup>	03 11Ru.A T	1970	IT=100	
<sup>98</sup> Y <sup>q</sup>	-71114	8	1181.1	0.4	806 ns	21 (10) <sup>-</sup>	03 11Ru.A T	1972	IT=100	
... A-group is continued on next page ...										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...										
<sup>98</sup> Zr	-81287	8	30.7 s	0.4	0 <sup>+</sup>	03	1967	$\beta^-$ =100		
<sup>98</sup> Zr <sup>m</sup>	-74683	8	6603.7	0.3	1.9 $\mu$ s	0.2 (17 <sup>-</sup> )	06Si36 EJT	2005 IT=100		
<sup>98</sup> Nb	-83525	5	2.86 s	0.06	1 <sup>+</sup>	03	1960	$\beta^-$ =100		
<sup>98</sup> Nb <sup>m</sup>	-83441	6	84	4	51.3 m	0.4 (5 <sup>+</sup> )	03	1948 $\beta^- \approx 100$ ; IT=0.1#		
<sup>98</sup> Mo	-88115.97	0.17	STABLE	(>100 Ty)	0 <sup>+</sup>	03	52Fr23 T	1930 IS=24.39 37; 2 $\beta^-$ ?		
<sup>98</sup> Tc	-86432	3	4.2 My	0.3	(6 <sup>+</sup> )	03	1955	$\beta^-$ =100; $\beta^+$ =0		
<sup>98</sup> Tc <sup>m</sup>	-86341	3	90.76	0.16	14.7 $\mu$ s	0.3 (2 <sup>-</sup> )	03	1976 IT=100		
<sup>98</sup> Ru	-88225	6	STABLE		0 <sup>+</sup>	03	1944	IS=1.87 3		
<sup>98</sup> Rh	-83175	12	* 8.72 m	0.12	(2 <sup>+</sup> )	03	1955	$\beta^+$ =100		
<sup>98</sup> Rh <sup>m</sup>	-83120#	50#	* 60#	50#	* 3.6 m	0.2 (5 <sup>+</sup> )	03	1966 IT=89 5; $\beta^+$ =11 5		
<sup>98</sup> Pd	-81321	5	17.7 m	0.3	0 <sup>+</sup>	03	1955	$\beta^+$ =100		
<sup>98</sup> Ag	-73070	30	47.5 s	0.3	(5,6) <sup>+</sup>	03	14Fe01 J	1978 $\beta^+$ =100; $\beta^+$ p=0.0012 5		
<sup>98</sup> Ag <sup>m</sup>	-72900	30	167.83	0.15	220 ns	20 (3 <sup>+</sup> )	03	98Gr.B ETD	1998 IT=100	
<sup>98</sup> Cd	-67640	50	9.2 s	0.3	0 <sup>+</sup>	03	1978	$\beta^+$ =100; $\beta^+$ p<0.025		
<sup>98</sup> Cd <sup>m</sup>	-65210	50	2427.5	0.6	189 ns	19 (8 <sup>+</sup> )	03	04B110 TJ	1996 IT=100	
<sup>98</sup> Cd <sup>n</sup>	-61010	50	6635	2	240 ns	40 (12 <sup>+</sup> )	03	04B110 ETJ	2004 IT=100	
<sup>98</sup> In	-53900#	300#	* 37 ms	5	0 <sup>+</sup> #	03	12Lo08 TD	1994 $\beta^+$ =100; $\beta^+$ p=5.6 3		
<sup>98</sup> In <sup>m</sup>	-53900#	580#	* 0#	500#	* 1.01 s	0.13	03	12Lo08 TD	2001 $\beta^+$ =100; $\beta^+$ p=19 2	
<sup>98</sup> Kr	T : average 11Ni01=42(4) 03Be05=46(8)									
<sup>98</sup> Rb	T : also 11Ni01=102(4), maybe mixture									
<sup>98</sup> Rb <sup>m</sup>	I : also an isomer with T=700(+60-50) ns decaying by $\gamma$ -rays of 178, 124 keV									
<sup>98</sup> Rb <sup>n</sup>	E : average 12Ka36=178.4(0.5) 09Fo05=178.0(0.7) T : other 09Fo05=700(+60-50)									
<sup>98</sup> Y <sup>m</sup>	J : 04Br14=(5 <sup>+</sup> ) 95Ha.B=(4 <sup>-</sup> ) 94St31=(5 <sup>+</sup> )									
<sup>98</sup> Y <sup>p</sup>	J : from 04Br14; ENSDF=(2 <sup>-</sup> ) and (p1/2[303]+n9/2[404]) config (in error)									
<sup>98</sup> Y <sup>q</sup>	J : from 04Br14; ENSDF=(8 <sup>-</sup> ) from (2 <sup>-</sup> ) for 496 keV isomer									
<sup>98</sup> Mo	T : 52Fr23 : 0v- $\beta\beta$ >100 Ty (theoretically faster, see text)									
<sup>98</sup> Ag	D : symmetrized from $\beta^+$ p=0.0011(+5-4)%									
<sup>98</sup> Cd <sup>m</sup>	T : average 04B110=170(+60-40) 98Gr.B=190(20), the latter supersedes									
<sup>98</sup> Cd <sup>m</sup>	T : 97Gr02=200(+300-170); other 97Go18=480(160) outweighed									
<sup>98</sup> Cd <sup>n</sup>	T : symmetrized from 230(+40-30) E : unc. estimated by evaluator									
<sup>98</sup> In	T : average 12Lo08=47(13) 10St.A=32(6) 08Ba53=44(+13-12) 01Ki13=32(+32-11)									
<sup>98</sup> In	D : $\beta^+$ p symmetrized from 12Lo08=5.5(+0.3-0.2)									
<sup>98</sup> In <sup>m</sup>	T : average 12Lo08=1.27(0.30) 10St.A=0.86(0.21) 08Ba53=0.92(+0.27-0.17) and									
<sup>98</sup> In <sup>m</sup>	T : 01Ki13=1.2(+1.2-0.4)									
<sup>99</sup> Kr	-38760#	400#	40 ms	11	5/2 <sup>-</sup> #	11	03Be05 TD	1997 $\beta^-$ =100; $\beta^-$ n=11 7; $\beta^-$ 2n=2#		
<sup>99</sup> Rb	-51121	4	56.4 ms	1.2	(3/2 <sup>+</sup> )	15	1971	$\beta^-$ =100; $\beta^-$ n=15.8 24; $\beta^-$ 2n=0.01#		
<sup>99</sup> Sr	-62521	5	269 ms	1	3/2 <sup>+</sup>	11	1975	$\beta^-$ =100; $\beta^-$ n=0.100 19		
<sup>99</sup> Y	-70650	7	1.484 s	0.007	5/2 <sup>+</sup>	11	07Ch07 J	1975 $\beta^-$ =100; $\beta^-$ n=1.7 4		
<sup>99</sup> Y <sup>m</sup>	-68508	7	2141.65	0.19	8.6 $\mu$ s	0.8 (17/2 <sup>+</sup> )	11	1985 IT=100		
<sup>99</sup> Zr	-77621	11	2.1 s	0.1	1/2 <sup>+</sup>	11	02Ca37 J	1970 $\beta^-$ =100		
<sup>99</sup> Zr <sup>m</sup>	-77369	11	251.96	0.09	293 ns	10	7/2 <sup>+</sup>	11	FGK126 J	1970 IT=100
<sup>99</sup> Nb	-82335	12	15.0 s	0.2	9/2 <sup>+</sup>	11	1950	$\beta^-$ =100		
<sup>99</sup> Nb <sup>m</sup>	-81970	12	365.27	0.08	2.5 m	0.2	1/2 <sup>-</sup>	11	1960 $\beta^-$ =?; IT<3.8	
<sup>99</sup> Mo	-85970.10	0.23	65.976 h	0.024	1/2 <sup>+</sup>	11	1948	$\beta^-$ =100		
<sup>99</sup> Mo <sup>m</sup>	-85872.32	0.23	97.785	0.003	15.5 $\mu$ s	0.2	5/2 <sup>+</sup>	11	1958 IT=100	
<sup>99</sup> Mo <sup>n</sup>	-85286.00	0.30	684.10	0.19	760 ns	60	11/2 <sup>-</sup>	11	1975 IT=100	
<sup>99</sup> Tc	-87327.9	0.9	211.1 ky	1.2	9/2 <sup>+</sup>	11	1938	$\beta^-$ =100		
<sup>99</sup> Tc <sup>m</sup>	-87185.2	0.9	142.6832	0.0011	6.0067 h	0.0005	1/2 <sup>-</sup>	11	1958 IT $\approx$ 100; $\beta^-$ =0.0037 6	
<sup>99</sup> Ru	-87625.4	0.3	STABLE		5/2 <sup>+</sup>	11	1931	IS=12.76 14		
<sup>99</sup> Rh	-85581	7	16.1 d	0.2	(1/2 <sup>-</sup> )	11	1952	$\beta^+$ =100		
<sup>99</sup> Rh <sup>m</sup>	-85516	7	64.6	0.5	4.7 h	0.1	9/2 <sup>+</sup>	11	1952 $\beta^+ \approx 100$ ; IT<0.16	
<sup>99</sup> Pd	-82183	5	21.4 m	0.2	(5/2 <sup>+</sup> )	11	1955	$\beta^+$ =100		
<sup>99</sup> Ag	-76712	6	2.07 m	0.05	(9/2 <sup>+</sup> )	11	14Fe01 J	1967 $\beta^+$ =100		
<sup>99</sup> Ag <sup>m</sup>	-76206	6	506.1	0.4	10.5 s	0.5	(1/2 <sup>-</sup> )	11	14Fe01 J	1978 IT=100
<sup>99</sup> Cd	-69931.1	1.6	16 s	3	5/2 <sup>+</sup> #	11	1978	$\beta^+$ =100; $\beta^+$ p=0.21 8; $\beta^+ \alpha < 1e-4$		
<sup>99</sup> In	-61380#	300#	3.1 s	0.2	9/2 <sup>+</sup> #	11	12Lo08 TD	1994 $\beta^+$ =100; $\beta^+$ p=0.9 4		
<sup>99</sup> In <sup>m</sup>	-60980#	340#	400#	150#	1# s		1/2 <sup>-</sup> #			
<sup>99</sup> Sn	-47940#	500#	5# ms	(>0.2 $\mu$ s)	9/2 <sup>+</sup> #		10St.A I	2011 $\beta^+$ ?; $\beta^+$ p ?		
<sup>99</sup> Sn <sup>m</sup>	-47540#	510#	400#	100#	1/2 <sup>-</sup> #		Mirror I			
<sup>99</sup> Kr	T : also 11Ni01=13(+34-6)									
<sup>99</sup> Zr <sup>m</sup>	J : 130.2 $\gamma$ ray, E2 to 3/2 <sup>+</sup> and 121.7 keV, $\gamma$ ray, M1 to 1/2 <sup>+</sup>									
<sup>99</sup> Cd	D : symmetrized from $\beta^+$ p=0.17(+11-5)%									
<sup>99</sup> In	T : recent not used 01Ki13=3.0(+0.8-0.7)									
<sup>99</sup> Sn	I : the 3 events reported in 95Ry03 are not trusted by NUBASE									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>100</sup> Kr	-35050#	400#	12 ms	8	0 <sup>+</sup>	11 11Ni01 TD	1997	$\beta^-$ =100; $\beta^-$ -n=10#; $\beta^-$ -2n=0.4# *		
<sup>100</sup> Rb	-46247	20	48 ms	3	(3 <sup>+</sup> ), 4 <sup>-</sup> #	08 11Ni01 T	1978	$\beta^-$ =100; $\beta^-$ -n=6 3; $\beta^-$ -2n=0.16 8		
<sup>100</sup> Sr	-59821	7	202 ms	3	0 <sup>+</sup>	08	1978	$\beta^-$ =100; $\beta^-$ -n=0.78 13		
<sup>100</sup> Sr <sup>m</sup>	-58202	7	1618.72	0.20	122 ns	9 (4 <sup>-</sup> )	12Ka36 T	1995	IT=100 *	
<sup>100</sup> Y	-67327	11	735 ms	7	(1 <sup>-</sup> )	08 83Wo10 J	1977	$\beta^-$ =100; $\beta^-$ -n=0.92 8 *		
<sup>100</sup> Y <sup>m</sup>	-67183	11	144	16	MD	940 ms	30 4(+)	08 13Ma15 J	1977	$\beta^-$ =100; $\beta^-$ -n=1#
<sup>100</sup> Zr	-76377	8	7.1 s	0.4	0 <sup>+</sup>	08	1970	$\beta^-$ =100		
<sup>100</sup> Nb	-79797	8	1.5 s	0.2	1 <sup>+</sup>	08	1967	$\beta^-$ =100		
<sup>100</sup> Nb <sup>m</sup>	-79484.7	2.0	313	8	MD	2.99 s	0.11 (5 <sup>+</sup> )	08	1967	$\beta^-$ =100
<sup>100</sup> Nb <sup>n</sup>	-79450	11	347	8	460 ns	60 (4 <sup>-</sup> , 5 <sup>-</sup> , 6 <sup>-</sup> )	08	1986	IT=100 *	
<sup>100</sup> Nb <sup>p</sup>	-79063	11	734	8	12.43 $\mu$ s	0.26 (8 <sup>-</sup> )	08 11Ru.A T	1980	IT=100 *	
<sup>100</sup> Mo	-86193.0	0.3	7.1 Ey	0.4	0 <sup>+</sup>	08 15Ba11 T	1930	IS=9.82 31; 2 $\beta^-$ =100 *		
<sup>100</sup> Tc	-86020.9	1.4	15.46 s	0.19	1 <sup>+</sup>	08	1952	$\beta^-$ $\approx$ 100; $\epsilon$ =0.0018 9		
<sup>100</sup> Tc <sup>m</sup>	-85820.2	1.4	200.67	0.04	8.32 $\mu$ s	0.14 (4 <sup>+</sup> )	08	1958	IT=100	
<sup>100</sup> Tc <sup>n</sup>	-85777.0	1.4	243.95	0.04	3.2 $\mu$ s	0.2 (6 <sup>+</sup> )	08	1967	IT=100	
<sup>100</sup> Ru	-89227.4	0.3	STABLE		0 <sup>+</sup>	08	1931	IS=12.60 7		
<sup>100</sup> Rh	-85591	18	20.8 h	0.1	1 <sup>-</sup>	08	1948	$\epsilon$ =95.1 5; $\epsilon^+$ =4.9 5		
<sup>100</sup> Rh <sup>m</sup>	-85516	18	74.782	0.014	214.0 ns	2.0 (2 <sup>+</sup> )	08	1965	IT=100	
<sup>100</sup> Rh <sup>n</sup>	-85483	18	107.6	0.2	4.6 m	0.2 (5 <sup>+</sup> )	08	1973	IT $\approx$ 98.3; $\beta^+$ $\approx$ 1.7	
<sup>100</sup> Rh <sup>p</sup>	-85371	18	219.61	0.22	130 ns	10 (7 <sup>+</sup> )	08	1984	IT=100	
<sup>100</sup> Pd	-85213	18	3.63 d	0.09	0 <sup>+</sup>	08	1948	$\epsilon$ =100		
<sup>100</sup> Ag	-78138	5	2.01 m	0.09	(5 <sup>+</sup> )	08 14Fe01 J	1970	$\beta^+$ =100		
<sup>100</sup> Ag <sup>m</sup>	-78122	5	15.52	0.16	2.24 m	0.13 (2 <sup>+</sup> )	08	1980	$\beta^+$ =?; IT ?	
<sup>100</sup> Cd	-74194.6	1.7	49.1 s	0.5	0 <sup>+</sup>	10	1970	$\beta^+$ =100		
<sup>100</sup> In	-64310	180	5.83 s	0.17	6 <sup>+</sup> #	14 12Lo08 TD	1982	$\beta^+$ =100; $\beta^+$ p=1.64 24 *		
<sup>100</sup> Sn	-57280	300	1.16 s	0.16	0 <sup>+</sup>	14 12Hi07 T	1994	$\beta^+$ =100; $\beta^+$ p<17 *		
<sup>100</sup> Sr <sup>m</sup>	-52780#	360#	4500#	200#	100# ns	6 <sup>+</sup> #		p ?		
* <sup>100</sup> Kr	T: symmetrized from 11Ni01=7(+11-3)								**	
* <sup>100</sup> Sr <sup>m</sup>	E: also 12Ka36=129.6(0.5), 288.1(0.5) and 1201.8(0.5) $\gamma$ rays in cascade								**	
* <sup>100</sup> Sr <sup>m</sup>	E: to ground-state =1619.5(0.9) keV								**	
* <sup>100</sup> Sr <sup>m</sup>	T: other 95Pf04=85(7)								**	
* <sup>100</sup> Y	J: ENSDF=1 <sup>-</sup> , 2 <sup>-</sup> ; but 1 <sup>-</sup> is favored from (p5/2[303]+n3/2[411]), see 83Wo10								**	
* <sup>100</sup> Nb <sup>n</sup>	E: 34.3 keV above 5 <sup>+</sup> isomer								**	
* <sup>100</sup> Nb <sup>p</sup>	E: 420.7 keV above 5 <sup>+</sup> isomer								**	
* <sup>100</sup> Nb <sup>p</sup>	J: 28 keV, (E2) $\gamma$ to (6 <sup>-</sup> ). Mult. from intensity balances								**	
* <sup>100</sup> Mo	T: also 14Ca46=7.15(0.37stat)(0.66syst)								**	
* <sup>100</sup> Mo	T: and 15Ba11=670(+50-40) 14Ar08=750(60stat)(60syst) to first exc. 0 <sup>+</sup> state								**	
* <sup>100</sup> In	T: average 12Lo08=5.7(0.3) 02Pi03=5.9(0.2)								**	
* <sup>100</sup> In	D: $\beta^+$ p average 12Lo08=1.7(0.4) 02Pi03=1.6(0.3)								**	
* <sup>100</sup> Sn	T: average 12Hi07=1.16(0.20) 08Ba53=0.55(+0.70-0.31) 96Ki23=0.94(+0.54-0.26)								**	
<sup>101</sup> Kr	-29130#	500#	5# ms	(>400 ns)	5/2 <sup>+</sup> #	10 10Oh02 I	2010	$\beta^-$ ?; $\beta^-$ -n=20#; $\beta^-$ -2n=2#		
<sup>101</sup> Rb	-42850#	200#	31.8 ms	3.3	3/2 <sup>+</sup> #	06 11Ni01 T	1992	$\beta^-$ =100; $\beta^-$ -n=28 4; $\beta^-$ -2n=0.3# ? *		
<sup>101</sup> Sr	-55325	8	113.8 ms	1.7	(5/2 <sup>-</sup> )	06 11Ni01 T	1983	$\beta^-$ =100; $\beta^-$ -n=2.37 14 *		
<sup>101</sup> Y	-65061	7	426 ms	20	5/2 <sup>+</sup>	06 07Ch07 J	1983	$\beta^-$ =100; $\beta^-$ -n=1.94 18 *		
<sup>101</sup> Y <sup>m</sup>	-64730	7	331.5	0.7	190 ns	40 12Ka36 ETD	2012	IT=100		
<sup>101</sup> Y <sup>n</sup>	-63854	7	1207.0	1.6	870 ns	90 09Fo05 ETD	2009	IT=100 *		
<sup>101</sup> Zr	-73166	8	2.3 s	0.1	3/2 <sup>+</sup>	06 02Ca37 J	1972	$\beta^-$ =100		
<sup>101</sup> Nb	-78891	4	7.1 s	0.3	(5/2#) <sup>+</sup>	06	1970	$\beta^-$ =100		
<sup>101</sup> Mo	-83519.9	0.3	14.61 m	0.03	1/2 <sup>+</sup>	06	1941	$\beta^-$ =100		
<sup>101</sup> Mo <sup>m</sup>	-83506.4	0.3	13.497	0.009	226 ns	7 3/2 <sup>+</sup>	06	1977	IT=100	
<sup>101</sup> Mo <sup>n</sup>	-83462.9	0.3	57.015	0.011	133 ns	70 5/2 <sup>+</sup>	06	1977	IT=100	
<sup>101</sup> Tc	-86345	24	14.22 m	0.01	9/2 <sup>+</sup>	06	1941	$\beta^-$ =100		
<sup>101</sup> Tc <sup>m</sup>	-86137	24	207.526	0.020	636 $\mu$ s	8 1/2 <sup>-</sup>	06	1964	IT=100	
<sup>101</sup> Ru	-87958.1	0.4	STABLE		5/2 <sup>+</sup>	06	1931	IS=17.06 2		
<sup>101</sup> Ru <sup>m</sup>	-87430.5	0.4	527.56	0.10	17.5 $\mu$ s	0.4 11/2 <sup>-</sup>	06	1974	IT=100	
<sup>101</sup> Rh	-87412	6	3.3 y	0.3	1/2 <sup>-</sup>	06	1948	$\epsilon$ =100		
<sup>101</sup> Rh <sup>m</sup>	-87255	6	157.32	0.03	4.34 d	0.01 9/2 <sup>+</sup>	06	1944	$\epsilon$ =92.80 25; IT=7.20 25	
<sup>101</sup> Pd	-85432	5	8.47 h	0.06	5/2 <sup>+</sup>	06	1948	$\beta^+$ =100		
<sup>101</sup> Ag	-81334	5	11.1 m	0.3	9/2 <sup>+</sup>	06 14Fe01 J	1966	$\beta^+$ =100		
<sup>101</sup> Ag <sup>m</sup>	-81060	5	274.1	0.3	3.10 s	0.10 (1/2) <sup>-</sup>	06 14Fe01 J	1975	IT=100 *	
<sup>101</sup> Cd	-75836.5	1.5	1.36 m	0.05	5/2 <sup>+</sup> #	06	1969	$\beta^+$ =100		

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>101</sup> In	-68610# 200#		15.1 s 1.1	9/2 <sup>+</sup> #	06	97Sz04 T	1988	$\beta^+$ =100; $\beta^+$ p=? *
<sup>101</sup> In <sup>m</sup>	-68060# 220# 550# 100#		10# s	1/2 <sup>-</sup> #				$\beta^+$ =95#; IT=5#
<sup>101</sup> Sn	-60310 300		1.97 s 0.16	(7/2 <sup>+</sup> )	07	12Lo08 TD	1994	$\beta^+$ =100; $\beta^+$ p=21.0 7 *
* <sup>101</sup> Rb	T : average 11Ni01=31(+5-4) 95Lh04=32(5) **							
* <sup>101</sup> Sr	T : average 11Ni01=113(2) 86Wa17=114(4) 83Wo10=121(6) **							
* <sup>101</sup> Y	T : average 96Me09=400(20) 86Wa17=440(20) 83Wo10=500(50) **							
* <sup>101</sup> Y	T : <sup>93</sup> Ru01=279(9) conflicting, not used **							
* <sup>101</sup> Y <sup>m</sup>	E : 12Ka36=128.0(0.5) and 203.5(0.5) $\gamma$ rays in cascade to ground-state **							
* <sup>101</sup> Y <sup>m</sup>	T : symmetrized from 187(+49-38) **							
* <sup>101</sup> Y <sup>n</sup>	T : symmetrized from 860(+90-80) **							
* <sup>101</sup> Y <sup>n</sup>	E : from a least-squares fit to Eg using 09Fo05 level scheme **							
* <sup>101</sup> Nb	J : positive parity due to M1 + E2 $\gamma$ from a + exc. level **							
* <sup>101</sup> Ag <sup>m</sup>	J : from ENSDF : E3 $\gamma$ to (7/2) <sup>+</sup> level **							
* <sup>101</sup> In	T : average 97Sz04=14.9(1.2) 88Hu07=16(3) **							
* <sup>101</sup> Sn	T : average 12Lo08=2.1(0.2) 07Se04=1.3(0.5) 07Ka15=1.9(0.3) **							
* <sup>101</sup> Sn	D : $\beta^+$ p average 12Lo08=22(1) 10St.A=20(1) J : from 10Da17 **							
<sup>102</sup> Rb	-37710# 300#		37 ms 3	(4 <sup>+</sup> )	09	16Wa16 JD	1995	$\beta^-$ =100; $\beta^-$ n=65 22; $\beta^-$ 2n=2# *
<sup>102</sup> Sr	-52160 70		69 ms 6	0 <sup>+</sup>	09		1986	$\beta^-$ =100; $\beta^-$ n=5.5 15 *
<sup>102</sup> Y	-61173 4		298 ms 9	(2 <sup>-</sup> )	09	11Ha48 J	1983	$\beta^-$ =100; $\beta^-$ n=4.9 12 *
<sup>102</sup> Y <sup>m</sup>	-60970# 200# 200#		360 ms 40	(> 5)	09	11Ha48 J	1980	$\beta^-$ =100; $\beta^-$ n=4.9 12 *
<sup>102</sup> Zr	-71588 9		2.9 s 0.2	0 <sup>+</sup>	09		1970	$\beta^-$ =100
<sup>102</sup> Nb	-76304.5 2.5		4.3 s 0.4	(4 <sup>+</sup> )	09		1972	$\beta^-$ =100
<sup>102</sup> Nb <sup>m</sup>	-76210 8 94 7 MD		1.3 s 0.2	1 <sup>+</sup>	09		1976	$\beta^-$ =100
<sup>102</sup> Mo	-83566 8		11.3 m 0.2	0 <sup>+</sup>	09		1954	$\beta^-$ =100
<sup>102</sup> Tc	-84573 9		5.28 s 0.15	1 <sup>+</sup>	09		1954	$\beta^-$ =100
<sup>102</sup> Tc <sup>m</sup>	-84553 13 20 10		4.35 m 0.07	(4, 5)	09		1954	$\beta^-$ =98 2; IT=2 2
<sup>102</sup> Ru	-89106.4 0.4		STABLE	0 <sup>+</sup>	09		1931	IS=31.55 14
<sup>102</sup> Rh	-86783 6		207.0 d 1.5	(1 <sup>-</sup> , 2 <sup>-</sup> )	09	98Sh21 T	1941	$\beta^+$ =78 5; $\beta^-$ =22 5 *
<sup>102</sup> Rh <sup>m</sup>	-86642 6 140.73 0.09		3.742 y 0.010	6 <sup>+</sup>	09	99Gi14 J	1962	$\beta^+$ ≈100; IT=0.233 24
<sup>102</sup> Pd	-87903.2 0.6		STABLE	0 <sup>+</sup>	09		1935	IS=1.02 1; 2 $\beta^+$ ?
<sup>102</sup> Ag	-82247 8		12.9 m 0.3	5 <sup>(+)</sup>	09		1960	$\beta^+$ =100
<sup>102</sup> Ag <sup>m</sup>	-82238 8 9.40 0.07		7.7 m 0.5	2 <sup>+</sup>	09		1967	$\beta^+$ =51 5; IT=49 5
<sup>102</sup> Cd	-79659.7 1.7		5.5 m 0.5	0 <sup>+</sup>	09		1969	$\beta^+$ =100
<sup>102</sup> In	-70695 5		23.3 s 0.1	(6 <sup>+</sup> )	09	95Sz01 J	1981	$\beta^+$ =100; $\beta^+$ p=0.0093 13
<sup>102</sup> Sn	-64930 100		3.8 s 0.2	0 <sup>+</sup>	09		1994	$\beta^+$ =100
<sup>102</sup> Sn <sup>m</sup>	-62910 100 2017 2		367 ns 8	(6 <sup>+</sup> )	09	98Li50 E	1996	IT=100 *
* <sup>102</sup> Rb	T : also 15Lo04=37(10) 11Ni01=35(+15-8) D : $\beta^-$ n=18(8)% in 85Pf.A **							
* <sup>102</sup> Sr	T : also 11Ni01=85(15) **							
* <sup>102</sup> Y	J : in 11Ha48, combining 07Ch07=(2,3) with spectroscopy data from 91Hill **							
* <sup>102</sup> Rh	T : average 98Sh21=207.3(1.7) 61Hi06=206(3) **							
* <sup>102</sup> Sn	T : 95Fa.A=4.6(1.4) supersedes 95Sc28=4.5(0.7), preliminary from same group **							
* <sup>102</sup> Sn <sup>m</sup>	T : from 11Hi.A **							
<sup>103</sup> Rb	-33610# 400#		26 ms 11	3/2 <sup>+</sup> #	15	15Lo04 TD	2010	$\beta^-$ =100; $\beta^-$ n=50#; $\beta^-$ 2n=2# *
<sup>103</sup> Sr	-47420# 200#		53 ms 10	5/2 <sup>+</sup> #	15		1997	$\beta^-$ =100; $\beta^-$ n=2#; $\beta^-$ 2n=0.01#
<sup>103</sup> Y	-58458 11		239 ms 12	5/2 <sup>+</sup> #	09	11Ni01 T	1994	$\beta^-$ =100; $\beta^-$ n=8.0 17 *
<sup>103</sup> Zr	-67815 9		1.38 s 0.07	5/2 <sup>-</sup> #	09	09Pe06 TD	1987	$\beta^-$ =100; $\beta^-$ n<1
<sup>103</sup> Nb	-75029 4		1.5 s 0.2	5/2 <sup>+</sup> #	09		1971	$\beta^-$ =100; $\beta^-$ n=0#
<sup>103</sup> Mo	-80961 9		67.5 s 1.5	3/2 <sup>+</sup>	09	09Ch09 J	1963	$\beta^-$ =100
<sup>103</sup> Tc	-84604 10		54.2 s 0.8	5/2 <sup>+</sup>	09		1957	$\beta^-$ =100
<sup>103</sup> Ru	-87267.2 0.4		39.247 d 0.013	3/2 <sup>+</sup>	09		1945	$\beta^-$ =100 *
<sup>103</sup> Ru <sup>m</sup>	-87029.0 0.8 238.2 0.7		1.69 ms 0.07	11/2 <sup>-</sup>	09		1964	IT=100
<sup>103</sup> Rh	-88031.7 2.3		STABLE	1/2 <sup>-</sup>	09		1934	IS=100
<sup>103</sup> Rh <sup>m</sup>	-87991.9 2.3 39.753 0.006		56.114 m 0.009	7/2 <sup>+</sup>	09		1943	IT=100
<sup>103</sup> Pd	-87457.2 0.9		16.991 d 0.019	5/2 <sup>+</sup>	09		1950	$\epsilon$ =100
<sup>103</sup> Ag	-84803 4		65.7 m 0.7	7/2 <sup>+</sup>	09		1954	$\beta^+$ =100
<sup>103</sup> Ag <sup>m</sup>	-84669 4 134.45 0.04		5.7 s 0.3	1/2 <sup>-</sup>	09		1962	IT=100
<sup>103</sup> Cd	-80651.6 1.8		7.3 m 0.1	5/2 <sup>+</sup> #	09		1960	$\beta^+$ =100
<sup>103</sup> In	-74633 10		60 s 1	9/2 <sup>+</sup> #	09	97Sz04 T	1978	$\beta^+$ =100
<sup>103</sup> In <sup>m</sup>	-74001 10 631.7 0.1		34 s 2	1/2 <sup>-</sup> #	09		1988	$\beta^+$ =67; IT=33
<sup>103</sup> Sn	-66970 70		7.0 s 0.2	5/2 <sup>+</sup> #	09		1981	$\beta^+$ =100; $\beta^+$ p=1.2 1
<sup>103</sup> Sb	-56180# 300#		<49 ns	5/2 <sup>+</sup> #	15			p ?
* <sup>103</sup> Rb	T : symmetrized from 15Lo04=23(+13-9) **							
* <sup>103</sup> Y	T : average 11Ni01=234(+18-15) 09Pe06=260(+40-20) 96Me09=230(20) and **							
* <sup>103</sup> Y	T : <sup>96</sup> Lh04=190(50) D : average 09Pe06=8(2)% 96Me09=8(3)% **							
* <sup>103</sup> Ru	T : other recent 09Go29=39.210(0.038) **							



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{104}\text{Sr}$	-44110# 300#			50.6 ms 4.2	$0^+$	15	15Lo04 T	1997	$\beta^- = 100; \beta^- n = 9\#; \beta^- 2n = 0\#$ *
$^{104}\text{Y}$	-54060# 400#			197 ms 4		15		1994	$\beta^- = 100; \beta^- n = 34\ 10; \beta^- 2n = 0\#$ *
$^{104}\text{Zr}$	-65724 9			920 ms 28		07	09Pe06 TD	1990	$\beta^- = 100; \beta^- n < 1$ *
$^{104}\text{Nb}$	-71819.0 2.7			4.9 s 0.3	$(1^+)$	07		1971	$\beta^- = 100; \beta^- n = 0.06\ 3$ *
$^{104}\text{Nb}^m$	-71600 120 210 120		BD *	940 ms 40	high	07		1976	$\beta^- = 100; \beta^- n = 0.05\ 3$ *
$^{104}\text{Mo}$	-80350 9			60 s 2	$0^+$	07		1962	$\beta^- = 100$ *
$^{104}\text{Tc}$	-82503 25			18.3 m 0.3	$(3^+)$	07		1956	$\beta^- = 100$ *
$^{104}\text{Tc}^m$	-82433 25 69.7 0.2			3.5 $\mu\text{s}$ 0.3	$(5^+)$	07		1981	IT=100 *
$^{104}\text{Tc}^n$	-82397 25 106.1 0.3			400 ns 20	$(+)$	07		1999	IT=100 *
$^{104}\text{Ru}$	-88095.7 2.5			STABLE	$0^+$	07		1931	IS=18.62 27; $2\beta^- ?$ *
$^{104}\text{Rh}$	-86959.3 2.3			42.3 s 0.4	$1^+$	07		1939	$\beta^- \approx 100; \beta^+ = 0.45\ 10$ *
$^{104}\text{Rh}^m$	-86830.3 2.3 128.9679 0.0005			4.34 m 0.03	$5^+$	07		1939	IT $\approx$ 100; $\beta^- = 0.13\ 1$ *
$^{104}\text{Pd}$	-89395.1 1.3			STABLE	$0^+$	07		1935	IS=11.14 8 *
$^{104}\text{Ag}$	-85116 4			69.2 m 1.0	$5^+$	07		1955	$\beta^+ = 100$ *
$^{104}\text{Ag}^m$	-85109 4 6.90 0.22			33.5 m 2.0	$2^+$	07		1959	$\beta^+ \approx 100; \text{IT} < 0.07$ *
$^{104}\text{Cd}$	-83968.4 1.7			57.7 m 1.0	$0^+$	07		1955	$\beta^+ = 100$ *
$^{104}\text{In}$	-76183 6			1.80 m 0.03	$(5^+)$	07	13Ma15 J	1977	$\beta^+ = 100$ *
$^{104}\text{In}^m$	-76090 6 93.48 0.10			15.7 s 0.5	$(3^+)$	07		1988	IT=80; $\beta^+ = 20$ *
$^{104}\text{Sn}$	-71627 6			20.8 s 0.5	$0^+$	07		1985	$\beta^+ = 100$ *
$^{104}\text{Sb}$	-59170 120			470 ms 130		07	95Fa.A D	1995	$\beta^+ = ?; \beta^+ p < 7; p < 7; \alpha ?$ *
* $^{104}\text{Sr}$	T : average 15Lo04=53(5) 11Ni01=43(+9-7) **								
* $^{104}\text{Nb}$	D : $\beta^- n = 0.71\%$ of 83En03, conflicting, not used **								
* $^{104}\text{Tc}^m$	J : E2 $\gamma$ to $(3^+)$ level (from ENSDF) **								
* $^{104}\text{Ru}$	T : $0\nu\text{-}\beta\beta$ to 1st exc. state : 13Be09>650Ey 12An08>190Ey **								
* $^{104}\text{Sb}$	T : symmetrized from 440(+150-110) D : 95Fa.A supersedes 95Sc28 p<1% **								
$^{105}\text{Sr}$	-38610# 500#			39 ms 5	$5/2^+\#$	15		1997	$\beta^- = 100; \beta^- n = 10\#; \beta^- 2n = 1\#$ *
$^{105}\text{Y}$	-51270 1340			95 ms 9	$5/2^+\#$	15		1994	$\beta^- = 100; \beta^- n < 82; \beta^- 2n = 0\#$ *
$^{105}\text{Zr}$	-61465 12			670 ms 28	$(5/2^+)$	15		1992	$\beta^- = 100; \beta^- n < 2$ *
$^{105}\text{Nb}$	-69916 4			2.95 s 0.06	$5/2^+\#$	05		1984	$\beta^- = 100; \beta^- n = 1.7\ 9$ *
$^{105}\text{Mo}$	-77337 9			35.6 s 1.6	$(5/2^-)$	05		1962	$\beta^- = 100$ *
$^{105}\text{Tc}$	-82290 40			7.6 m 0.1	$(3/2^-)$	05		1955	$\beta^- = 100$ *
$^{105}\text{Ru}$	-85934.5 2.5			4.44 h 0.02	$3/2^+$	05		1945	$\beta^- = 100$ *
$^{105}\text{Ru}^m$	-85913.9 2.5 20.610 0.013			340 ns 15	$(5/2^+)$	05		1974	IT=100 *
$^{105}\text{Rh}$	-87851.2 2.5			35.357 h 0.037	$7/2^+$	05	09Go29 T	1945	$\beta^- = 100$ *
$^{105}\text{Rh}^m$	-87721.4 2.5 129.782 0.004			42.9 s 0.3	$1/2^-$	05		1950	IT=100 *
$^{105}\text{Pd}$	-88417.9 1.1			STABLE	$5/2^+$	05		1935	IS=22.33 8 *
$^{105}\text{Pd}^m$	-87928.8 1.1 489.14 0.04			36.1 $\mu\text{s}$ 0.4	$11/2^-$	05		1970	IT=100 *
$^{105}\text{Ag}$	-87071 5			41.29 d 0.07	$1/2^-$	05		1939	$\beta^+ = 100$ *
$^{105}\text{Ag}^m$	-87046 5 25.479 0.016			7.23 m 0.16	$7/2^+$	05		1969	IT $\approx$ 100; $\beta^+ = 0.34\ 7$ *
$^{105}\text{Cd}$	-84333.8 1.4			55.5 m 0.4	$5/2^+$	05		1950	$\beta^+ = 100$ *
$^{105}\text{In}$	-79641 10			5.07 m 0.07	$9/2^+$	05		1975	$\beta^+ = 100$ *
$^{105}\text{In}^m$	-78967 10 674.08 0.25			48 s 6	$(1/2^-)$	05		1975	IT=?; $\beta^+ = 25\#$ *
$^{105}\text{Sn}$	-73338 4			34 s 1	$(5/2^+)$	05	85De08 J	1981	$\beta^+ = 100; \beta^+ p = ?$ *
$^{105}\text{Sb}$	-64015 22			1.12 s 0.16	$(5/2^+)$	05	95Fa.A T	1994	$\beta^+ ?; p < 0.1; \beta^+ p ?$ *
$^{105}\text{Te}$	-52810 300			633 ns 66	$(7/2^+)$	06	06Se08 T	2006	$\alpha \approx 100$ *
* $^{105}\text{Sb}$	T : 95Fa.A supersedes 95Sc28=1.30(0.15), preliminary from same group **								
* $^{105}\text{Sb}$	D : p 05Li47<0.1% above 430 keV disagrees with 94Ti03 $\approx$ 1% **								
* $^{105}\text{Te}$	T : average 06Li41=620(70) 06Se08=700(+250-170) **								
* $^{105}\text{Te}$	J : same spin as 171.7 state in $^{101}\text{Sn}$ **								
$^{106}\text{Sr}$	-34790# 600#			21 ms 8	$0^+$	15	15Lo04 T	2010	$\beta^- = 100; \beta^- n = 10\#; \beta^- 2n = 0.03\#$ *
$^{106}\text{Y}$	-46050# 500#			74 ms 6	$2^+\#$	15	15Lo04 T	1997	$\beta^- = 100; \beta^- n = 20\#; \beta^- 2n = 0.5\#$ *
$^{106}\text{Zr}$	-58550 430			178.6 ms 5.8	$0^+$	15	15Lo04 T	1994	$\beta^- = 100; \beta^- n < 7$ *
$^{106}\text{Nb}$	-66203 4			1050 ms 100	$(1^-)$	15	14Lu07 J	1976	$\beta^- = 100; \beta^- n = 4.5\ 3$ *
$^{106}\text{Nb}^m$	-65998 4 204.8 0.1			800 ns 50	$(3^+)$	15		1999	IT=100 *
$^{106}\text{Nb}^n$	-65998 4 204.8 0.5			849 ns 45	$(3^+)$		14Lu07 EJ	1999	IT=100 *
$^{106}\text{Mo}$	-76135 9			8.73 s 0.12	$0^+$	08		1969	$\beta^- = 100$ *
$^{106}\text{Tc}$	-79776 12			35.6 s 0.6	$(1, 2)^{(\#)}$	08		1965	$\beta^- = 100$ *
$^{106}\text{Ru}$	-86323 5			371.8 d 0.18	$0^+$	08		1948	$\beta^- = 100$ *
$^{106}\text{Rh}$	-86363 5			30.07 s 0.35	$1^+$	08		1947	$\beta^- = 100$ *
$^{106}\text{Rh}^m$	-86231 10 132 11		BD	131 m 2	$(6)^+$	08		1955	$\beta^- = 100$ *
$^{106}\text{Pd}$	-89907.5 1.1			STABLE	$0^+$	08		1935	IS=27.33 3 *
$^{106}\text{Ag}$	-86942 3			23.96 m 0.04	$1^+$	08		1937	$\beta^+ = ?; \beta^- \approx 0.5$ *
$^{106}\text{Ag}^m$	-86852 3 89.66 0.07			8.28 d 0.02	$6^+$	08		1938	$\beta^+ = 100; \text{IT} \leq 4.2e-6$ *
$^{106}\text{Cd}$	-87132.1 1.1			STABLE ( $>1.1\ \text{Zy}$ )	$0^+$	08	16Be11 T	1935	IS=1.25 6; $2\beta^+ ?$ *
$^{106}\text{In}$	-80608 12			6.2 m 0.1	$7^+$	08		1962	$\beta^+ = 100$ *
$^{106}\text{In}^m$	-80579 12 28.6 0.3			5.2 m 0.1	$(2)^+$	08		1966	$\beta^+ = 100$ *

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>106</sup> Sn	-77354	5	1.92 m	0.08	0 <sup>+</sup>	08	1975	$\beta^+$ =100
<sup>106</sup> Sb	-66473	7	600 ms	200	(2 <sup>+</sup> )	08	1981	$\beta^+$ =100
<sup>106</sup> Sb <sup>m</sup>	-66370	7 103.5 0.3	226 ns	14	(4 <sup>+</sup> )	08 99So08 T	1998	IT=100
<sup>106</sup> Te	-58220	100	78 $\mu$ s	11	0 <sup>+</sup>	08 16Ca33 T	1981	$\alpha$ =100
* <sup>106</sup> Sr	T: symmetrized from 15Lo04=20(+8-7)							
* <sup>106</sup> Y	T: average 15Lo04=82(+10-5) 15NiZZ=62(9) 11Ni01=62(+25-14)							
* <sup>106</sup> Zr	T: average 15Lo04=175(7) 11Ni01=186(+11-10)							
* <sup>106</sup> Nb	T: unweighed average 09Pe06=1240(20) 96Me09=900(20) 83Sh06=1020(50)							
* <sup>106</sup> Nb <sup>n</sup>	T: average 12Ka36=660(+110-100) 99Ge01=890(50)							
* <sup>106</sup> Cd	T: for $\epsilon\beta^+$ , theoretically faster channel; others 12Be14>210Ey 02Tr04>410Ey							
* <sup>106</sup> Sb <sup>m</sup>	T: average 99So08=232(21) 98Li50=220(20)							
* <sup>106</sup> Te	T: average 16Ca33=70(+20-15) 05Ja03=85(+25-15) 94Pa11=60(+40-20) and							
* <sup>106</sup> Te	T: 81Sc17=60(+30-10)							
<sup>107</sup> Sr	-28900#	700#	10# ms	(>400 ns)	1/2 <sup>+</sup> #	10 100h02 I	2010	$\beta^-$ ?; $\beta^-$ -n=30#; $\beta^-$ -2n=3#
<sup>107</sup> Y	-42360#	500#	33.5 ms	0.3	5/2 <sup>+</sup> #	15 15Lo04 T	1997	$\beta^-$ =100; $\beta^-$ -n=30#; $\beta^-$ -2n=0.1#
<sup>107</sup> Zr	-54380	1120	145.7 ms	2.4	5/2 <sup>+</sup> #	15 15Lo04 T	1994	$\beta^-$ =100; $\beta^-$ -n<23
<sup>107</sup> Nb	-63724	8	289 ms	9	5/2 <sup>+</sup> #	08 15Lo04 T	1992	$\beta^-$ =100; $\beta^-$ -n=7.4 8
<sup>107</sup> Mo	-72552	9	3.5 s	0.5	(5/2 <sup>+</sup> )	08	1972	$\beta^-$ =100
<sup>107</sup> Mo <sup>m</sup>	-72487	9 65.4 0.2	420 ns	30	(1/2 <sup>+</sup> )	08	1976	IT=100
<sup>107</sup> Tc	-78750	9	21.2 s	0.2	(3/2 <sup>-</sup> )	08 09Gu11 J	1965	$\beta^-$ =100
<sup>107</sup> Tc <sup>m</sup>	-78720	9 30.1 0.1	3.85 $\mu$ s	0.05	(1/2 <sup>+</sup> )	08	2007	IT=100
<sup>107</sup> Tc <sup>n</sup>	-78684	9 65.72 0.14	184 ns	3	(5/2 <sup>+</sup> )	08	1974	IT=100
<sup>107</sup> Ru	-83863	9	3.75 m	0.05	(5/2 <sup>+</sup> )	08	1951	$\beta^-$ =100
<sup>107</sup> Rh	-86864	12	21.7 m	0.4	7/2 <sup>+</sup>	08	1951	$\beta^-$ =100
<sup>107</sup> Rh <sup>m</sup>	-86596	12 268.36 0.04	> 10 $\mu$ s		1/2 <sup>-</sup>	08	1986	IT=100
<sup>107</sup> Pd	-88372.6	1.2	6.5 My	0.3	5/2 <sup>+</sup>	08	1958	$\beta^-$ =100
<sup>107</sup> Pd <sup>m</sup>	-88256.9	1.2 115.74 0.12	850 ns	100	1/2 <sup>+</sup>	08	1969	IT=100
<sup>107</sup> Pd <sup>n</sup>	-88158.0	1.2 214.6 0.3	21.3 s	0.5	11/2 <sup>-</sup>	08	1952	IT=100
<sup>107</sup> Ag	-88406.7	2.4	STABLE		1/2 <sup>-</sup>	08 14Fe01 J	1924	IS=51.839 8
<sup>107</sup> Ag <sup>m</sup>	-88313.6	2.4 93.125 0.019	44.3 s	0.2	7/2 <sup>+</sup>	08	1940	IT=100
<sup>107</sup> Cd	-86990.3	1.7	6.50 h	0.02	5/2 <sup>+</sup>	08	1946	$\beta^+$ =100
<sup>107</sup> In	-83564	11	32.4 m	0.3	9/2 <sup>+</sup>	08	1949	$\beta^+$ =100
<sup>107</sup> In <sup>m</sup>	-82886	11 678.5 0.3	50.4 s	0.6	1/2 <sup>-</sup>	08	1973	IT=100
<sup>107</sup> Sn	-78512	5	2.90 m	0.05	(5/2 <sup>+</sup> )	08	1976	$\beta^+$ =100
<sup>107</sup> Sb	-70653	4	4.0 s	0.2	5/2 <sup>+</sup> #	08	1994	$\beta^+$ =100
<sup>107</sup> Te	-60540	70	3.1 ms	0.1	5/2 <sup>+</sup> #	08	1979	$\alpha$ =70 30; $\beta^+$ ?; $\beta^+$ p?
<sup>107</sup> I	-49430#	300#	20# $\mu$ s		5/2 <sup>+</sup> #			$\alpha$ ?
* <sup>107</sup> Zr	T: average 15Lo04=150(3) 11Ni01=138(4); not used 09Pe06=150(+40-30)							
* <sup>107</sup> Nb	T: average 15Lo04=280(20) 09Pe06=290(11) 96Me09=300(30)							
* <sup>107</sup> Nb	D: average 09Pe06=8(1)% 96Me09=6.0(1.5)%							
<sup>108</sup> Y	-37300#	600#	30 ms	5		15	2010	$\beta^-$ =100; $\beta^-$ -n=30#; $\beta^-$ -2n=2#
<sup>108</sup> Zr	-51350#	400#	78.5 ms	2.0	0 <sup>+</sup>	15	1997	$\beta^-$ =100; $\beta^-$ -n=2#
<sup>108</sup> Zr <sup>m</sup>	-49280#	400# 2074.5 0.8	540 ns	30	(6 <sup>+</sup> )	15 12Ka36 T	2011	IT $\approx$ 100
<sup>108</sup> Nb	-59546	8	198 ms	6	(2 <sup>+</sup> )	15	1994	$\beta^-$ =100; $\beta^-$ -n=6.3 5; $\beta^-$ -2n=0#
<sup>108</sup> Nb <sup>m</sup>	-59379	8 166.6 0.5	109 ns	2	(4 <sup>-</sup> , 5)	15 12Ka36 J	2012	IT=100
<sup>108</sup> Mo	-70756	9	1.105 s	0.010	0 <sup>+</sup>	08 09Pe06 TD	1972	$\beta^-$ =100; $\beta^-$ -n<0.5
<sup>108</sup> Tc	-75923	9	5.17 s	0.07	(2) <sup>+</sup>	08	1970	$\beta^-$ =100
<sup>108</sup> Ru	-83661	9	4.55 m	0.05	0 <sup>+</sup>	08	1955	$\beta^-$ =100
<sup>108</sup> Rh	-85032	14	16.8 s	0.5	1 <sup>+</sup>	08	1955	$\beta^-$ =100
<sup>108</sup> Rh <sup>m</sup>	-84917	12 115 18 MD	6.0 m	0.3	(5) <sup>(+)</sup> #	08	1969	$\beta^-$ =100
<sup>108</sup> Pd	-89524.2	1.1	STABLE		0 <sup>+</sup>	08	1935	IS=26.46 9
<sup>108</sup> Ag	-87606.8	2.4	2.382 m	0.011	1 <sup>+</sup>	08	1937	$\beta^-$ =97.15 20; $\beta^+$ =2.85 20
<sup>108</sup> Ag <sup>m</sup>	-87497.3	2.4 109.466 0.007	438 y	9	6 <sup>+</sup>	08	1969	$\beta^+$ =91.3 9; IT=8.7 9
<sup>108</sup> Cd	-89252.4	1.1	STABLE	(>410 Py)	0 <sup>+</sup>	08 95Ge14 T	1935	IS=0.89 3; 2 $\beta^+$ ?
<sup>108</sup> In	-84120	9	58.0 m	1.2	7 <sup>+</sup>	08	1949	$\beta^+$ =100
<sup>108</sup> In <sup>m</sup>	-84090	9 29.75 0.05	39.6 m	0.7	2 <sup>+</sup>	08	1955	$\beta^+$ =100
<sup>108</sup> Sn	-82070	5	10.30 m	0.08	0 <sup>+</sup>	08	1968	$\beta^+$ =100
<sup>108</sup> Sb	-72445	5	7.4 s	0.3	(4 <sup>+</sup> )	08	1976	$\beta^+$ =100
<sup>108</sup> Te	-65782	5	2.1 s	0.1	0 <sup>+</sup>	08 85Ti02 D	1974	$\beta^+$ =51 4; $\alpha$ =49 4; $\beta^+$ p=2.4 10; $\beta^+$ $\alpha$ <0.065
<sup>108</sup> I	-52650	130	36 ms	6	1 <sup>+</sup> #	08 94Pa12 D	1991	$\alpha$ ?; $\beta^+$ =9#; p<1; $\beta^+$ p?
* <sup>108</sup> Zr <sup>m</sup>	T: symmetrized from 12Ka36=536(+26-25); other 11Su11=620(150)							
* <sup>108</sup> Mo	T: average 09Pe06=1.110(0.011) 95Jo02=1.090(0.020) D: $\beta^-$ -n not allowed							
* <sup>108</sup> I	D: $\beta^+$ =9%# estimated in 94Pa12 using theoretical $\beta^+$ half-life $\approx$ 400 ms							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{109}\text{Y}$	-33200# 700#		25 ms 5	$5/2^+\#$	15		2010	$\beta^-\approx 100; \beta^-\text{n}=60\#; \beta^-\text{2n}=1.5\#$
$^{109}\text{Zr}$	-46190# 500#		56 ms 3	$1/2^+\#$	15		1997	$\beta^-\approx 100; \beta^-\text{n}=5\#; \beta^-\text{2n}=0\#$
$^{109}\text{Nb}$	-56690 260		106.9 ms 4.9	$5/2^+\#$	06	15Lo04 T	1994	$\beta^-\approx 100; \beta^-\text{n}=31.5$
$^{109}\text{Nb}^m$	-56380 260	312.2 0.5	115 ns 8			12Ka36 ET	2011	IT=100
$^{109}\text{Mo}$	-66666 11		700 ms 14	$5/2^+\#$	06	09Pe06 TD	1992	$\beta^-\approx 100; \beta^-\text{n}=1.3.6$
$^{109}\text{Mo}^m$	-66596 11	69.7 0.5	210 ns 60	$(1/2^+)$		12Ka36 ET	2012	IT=100
$^{109}\text{Tc}$	-74283 10		1.14 s 0.03	$5/2^+$	06	09Pe06 T	1976	$\beta^-\approx 100; \beta^-\text{n}=0.08.2$
$^{109}\text{Ru}$	-80738 9		34.5 s 1.0	$5/2^+\#$	06		1967	$\beta^-\approx 100$
$^{109}\text{Ru}^m$	-80642 9	96.2 0.3	680 ns 30	$(5/2^-)$	06		1976	IT=100
$^{109}\text{Rh}$	-84999 4		80 s 2	$7/2^+$	06		1972	$\beta^-\approx 100$
$^{109}\text{Rh}^m$	-84773 4	225.974 0.021	1.66 $\mu\text{s}$ 0.04	$3/2^+$	06	FGK127 J	1987	IT=100
$^{109}\text{Pd}$	-87606.5 1.1		13.7012 h 0.0024	$5/2^+$	06		1937	$\beta^-\approx 100$
$^{109}\text{Pd}^m$	-87493.1 1.1	113.400 0.010	380 ns 50.	$1/2^+$	06		1978	IT=100
$^{109}\text{Pd}^n$	-87417.5 1.1	188.990 0.010	4.696 m 0.003	$11/2^-$	06		1957	IT=100
$^{109}\text{Ag}$	-88719.4 1.3		STABLE	$1/2^-$	06		1924	IS=48.161.8
$^{109}\text{Ag}^m$	-88631.4 1.3	88.0341 0.0011	39.6 s 0.2	$7/2^+$	06		1967	IT=100
$^{109}\text{Cd}$	-88504.3 1.5		461.6 d 0.4	$5/2^+$	06	16Fe04 T	1950	$\epsilon=100$
$^{109}\text{Cd}^m$	-88444.8 1.5	59.49 0.11	12 $\mu\text{s}$ 2	$1/2^+$	06		1956	IT=100
$^{109}\text{Cd}^n$	-88040.8 1.5	463.5 0.3	10.9 $\mu\text{s}$ 0.5	$11/2^-$	06		1964	IT=100
$^{109}\text{In}$	-86490 4		4.167 h 0.018	$9/2^+$	06		1948	$\beta^+\approx 100$
$^{109}\text{In}^m$	-85840 4	650.1 0.3	1.34 m 0.07	$1/2^-$	06		1966	IT=100
$^{109}\text{In}^n$	-84388 4	2101.8 0.2	209 ms 6	$(19/2^+)$	06		1963	IT=100
$^{109}\text{Sn}$	-82630 8		18.0 m 0.2	$(5/2^+, 7/2^+)$	06	13Ma15 J	1966	$\beta^+\approx 100$
$^{109}\text{Sb}$	-76251 5		17.0 s 0.7	$5/2^+\#$	06		1976	$\beta^+\approx 100$
$^{109}\text{Te}$	-67715 4		4.6 s 0.3	$(5/2^+)$	06		1967	$\beta^+\approx 96.1.13; \alpha=3.9.13; \dots$
$^{109}\text{I}$	-57672 7		103 $\mu\text{s}$ 5	$1/2^+$	06	07Ma35 D	1984	$p=100; \alpha=0.014.4$
$^{109}\text{Xe}$	-46170 300		13 ms 2	$7/2^+\#$		06Li41 TDJ	2006	$\alpha\approx 100; \beta^+ ?; \beta^+ p ?$
$^{109}\text{Nb}$	T : average 15Lo04=110(6) 11Ni01=100(+9-8); other 09Pe06=130(20)							
$^{109}\text{Nb}$	D : 09Pe06 $\beta^-\text{n}<15\%$ conflicting							
$^{109}\text{Nb}^m$	E : other 11Wa03=313.1(0.5) keV							
$^{109}\text{Nb}^n$	T : symmetrized from 12Ka36=114(+8-7); other 11Wa03=150(30)							
$^{109}\text{Mo}$	T : also 15Lo04=700(+40-60)							
$^{109}\text{Mo}^m$	T : symmetrized from 12Ka36=194(+76-49)							
$^{109}\text{Tc}$	J : 12Ku28=5/2 <sup>+</sup>							
$^{109}\text{Rh}^m$	J : 225.9 keV E2 $\gamma$ ray to 7/2 <sup>+</sup>							
$^{109}\text{Cd}$	T : unweighed average 16Fe04=462.1(0.3) 14Un01=462.3(0.8)							
$^{109}\text{Cd}$	T : 11Va02=462.29(0.30) 04Sc04=459.6(1.7) 97Ma75=460.15(0.16)							
$^{109}\text{Cd}$	T : 82La25=463.1(0.8) 81Va11=461.9(0.3)							
$^{109}\text{Te}$	D : ... ; $\beta^+p=9.4.31; \beta^+\alpha<0.005$							
$^{109}\text{Xe}$	J : same as 150 level in $^{105}\text{Te}$							
$^{110}\text{Zr}$	-42890# 600#		37.5 ms 2.0	$0^+$	12	15Lo04 T	1997	$\beta^-\approx 100; \beta^-\text{n}=7\#; \beta^-\text{2n}=0\#$
$^{110}\text{Nb}$	-52310 840		82 ms 2	$(5)^{(+\#)}$	12	15Lo04 T	1994	$\beta^-\approx 100; \beta^-\text{n}=40.8; \beta^-\text{2n}=0.09\#$
$^{110}\text{Mo}$	-64543 24		292 ms 7	$0^+$	12	15Lo04 T	1992	$\beta^-\approx 100; \beta^-\text{n}=2.0.7$
$^{110}\text{Tc}$	-71035 9		900 ms 13	$(2^+, 3^+)$	12		1976	$\beta^-\approx 100; \beta^-\text{n}=0.04.2$
$^{110}\text{Ru}$	-80073 9		12.04 s 0.17	$0^+$	12		1970	$\beta^-\approx 100$
$^{110}\text{Rh}$	-82829 18		3.35 s 0.12	$(1^+)$	12		1963	$\beta^-\approx 100$
$^{110}\text{Rh}^m$	-82610# 150#	220# 150#	28.5 s 1.3	$(6^+)$	12		1969	$\beta^-\approx 100$
$^{110}\text{Pd}$	-88330.9 0.6		STABLE (>200 Ey)	$0^+$	12	13Le10 T	1935	IS=11.72.9; $2\beta^- ?$
$^{110}\text{Ag}$	-87457.3 1.3		24.56 s 0.11	$1^+$	12		1937	$\beta^-\approx 100; \epsilon=0.30.6$
$^{110}\text{Ag}^m$	-87456.2 1.3	1.112 0.016	660 ns 40	$2^-$	12		1975	IT=100
$^{110}\text{Ag}^n$	-87339.7 1.3	117.59 0.05	249.83 d 0.04	$6^+$	12		1938	$\beta^-\approx 98.67.8; \text{IT}=1.33.8$
$^{110}\text{Cd}$	-90348.0 0.4		STABLE	$0^+$	12		1925	IS=12.49.18
$^{110}\text{In}$	-86470 12		4.92 h 0.08	$7^+$	12		1939	$\beta^+\approx 100$
$^{110}\text{In}^m$	-86408 12	62.08 0.04	69.1 m 0.5	$2^+$	12		1962	$\beta^+\approx 100$
$^{110}\text{Sn}$	-85842 14		4.154 h 0.004	$0^+$	12		1965	$\epsilon=100$
$^{110}\text{Sb}$	-77450 6		23.6 s 0.3	$(3^+)$	12		1972	$\beta^+\approx 100$
$^{110}\text{Te}$	-72230 7		18.6 s 0.8	$0^+$	12		1977	$\beta^+\approx 100; \alpha=0.003\#$
$^{110}\text{I}$	-60460 50		664 ms 24	$(1^+)$	12		1977	$\beta^+=83.4; \alpha=17.4; \beta^+p=11.3; \beta^+\alpha=1.1.3$
$^{110}\text{Xe}$	-51920 100		93 ms 3	$0^+$	12		1981	$\alpha=64.35; \beta^+ ?; \beta^+ p ?$
$^{110}\text{Pd}$	T : >198Ey, >172Ey(95% CL) for first excited $0^+$ and $2^+$ ; 52Wi26>0.6Ey							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>111</sup> Zr	-37560#	700#		24.0 ms	0.5	3/2 <sup>+</sup> #	15 15Lo04	T 2010	$\beta^- = 100; \beta^-_n = 10\#; \beta^-_{2n} = 1\#$
<sup>111</sup> Nb	-48880#	300#		54 ms	2	5/2 <sup>+</sup> #	15	1997	$\beta^- = 100; \beta^-_n = 90\#; \beta^-_{2n} = 0\#$
<sup>111</sup> Mo	-59940	13		193.6 ms	4.4	1/2 <sup>+</sup> #	15 15Lo04	T 1994	$\beta^- = 100; \beta^-_n < 12$
<sup>111</sup> Mo <sup>m</sup>	-59840#	50#	100# 50#	200 ms		7/2 <sup>-</sup> #	15	2011	$\beta^- = 100; \beta^-_n = 0.02\#$
<sup>111</sup> Tc	-69025	11		350 ms	11	5/2 <sup>+</sup> #	09 09Pe06	T 1988	$\beta^- = 100; \beta^-_n = 0.85 20$
<sup>111</sup> Ru	-76785	10		2.12 s	0.07	5/2 <sup>+</sup>	09	1971	$\beta^- = 100$
<sup>111</sup> Rh	-82304	7		11 s	1	(7/2 <sup>+</sup> )	09	1975	$\beta^- = 100$
<sup>111</sup> Pd	-85985.9	0.7		23.4 m	0.2	5/2 <sup>+</sup>	09	1937	$\beta^- = 100$
<sup>111</sup> Pd <sup>m</sup>	-85813.7	0.7	172.18 0.08	5.5 h	0.1	11/2 <sup>-</sup>	09	1952	IT=73 3; $\beta^- = 27 3$
<sup>111</sup> Ag	-88215.4	1.5		7.433 d	0.010	1/2 <sup>-</sup>	09 16Co01	T 1937	$\beta^- = 100$
<sup>111</sup> Ag <sup>m</sup>	-88155.6	1.5	59.82 0.04	64.8 s	0.8	7/2 <sup>+</sup>	09	1957	IT=99.3 2; $\beta^- = 0.7 2$
<sup>111</sup> Cd	-89252.2	0.4		STABLE		1/2 <sup>+</sup>	09	1925	IS=12.80 12
<sup>111</sup> Cd <sup>m</sup>	-88856.0	0.4	396.214 0.021	48.50 m	0.09	11/2 <sup>-</sup>	09	1945	IT=100
<sup>111</sup> In	-88392	3		2.8063 d	0.0007	9/2 <sup>+</sup>	09 16Dz01	T 1947	$\epsilon = 100$
<sup>111</sup> In <sup>m</sup>	-87855	3	536.99 0.07	7.7 m	0.2	1/2 <sup>-</sup>	09	1966	IT=100
<sup>111</sup> Sn	-85939	5		35.3 m	0.6	7/2 <sup>+</sup>	09	1949	$\beta^+ = 100$
<sup>111</sup> Sn <sup>m</sup>	-85684	5	254.71 0.04	12.5 $\mu$ s	1.0	1/2 <sup>+</sup>	09	1972	IT=100
<sup>111</sup> Sb	-80837	9		75 s	1	(5/2 <sup>+</sup> )	09	1972	$\beta^+ = 100$
<sup>111</sup> Te	-73587	6		26.2 s	0.6	(5/2 <sup>+</sup> )	09 05Sh24	T 1967	$\beta^+ = 100; \beta^+ p = ?$
<sup>111</sup> I	-64954	5		2.5 s	0.2	5/2 <sup>+</sup> #	09	1977	$\beta^+ \approx 100; \alpha \approx 0.1; \beta^+ p ?$
<sup>111</sup> Xe	-54400	90		740 ms	200	5/2 <sup>+</sup> #	09 12Ca03	D 1979	$\beta^+ ?; \alpha = 10.4 1.9; \beta^+ p ?$
<sup>111</sup> Cs	-42820#	200#		1# $\mu$ s		3/2 <sup>+</sup> #			p ?
* <sup>111</sup> Mo	T : average 15Lo04=196(5) 11Ku16=186(9); other 09Pe06=200(+41-36)								
* <sup>111</sup> Ag	T : average 16Co01=7.423(0.013) 74Ro18=7.450(0.017)								
* <sup>111</sup> In	T : average 16Dz01=2.8067(0.0034) 14Un01=2.805(0.004) 04Sc04=2.8063(0.0007)								
* <sup>111</sup> Te	T : others 67Ka01=19.0(7) 67Bo41=19.5(5) conflicting, not used								
<sup>112</sup> Zr	-33810#	700#		43 ms	21	0 <sup>+</sup>	15 15Lo04	T 2010	$\beta^- = 100; \beta^-_n = 30\#; \beta^-_{2n} = 0.3\#$
<sup>112</sup> Nb	-44270#	300#		38 ms	2	2 <sup>+</sup> #	15 15Lo04	T 1997	$\beta^- = 100; \beta^-_n = 70\#; \beta^-_{2n} = 1\#$
<sup>112</sup> Mo	-57460#	200#		125 ms	5	0 <sup>+</sup>	15 15Lo04	T 1994	$\beta^- = 100; \beta^-_n = 0.3\#$
<sup>112</sup> Tc	-65259	6		323 ms	6	(2 <sup>+</sup> )	15 15Lo04	T 1990	$\beta^- = 100; \beta^-_n = 1.5 2$
<sup>112</sup> Tc <sup>m</sup>	-64907	6	352.3 0.7	150 ns	17		15 FGK127	E 2010	IT=100
<sup>112</sup> Ru	-75631	10		1.75 s	0.07	0 <sup>+</sup>	15	1970	$\beta^- = 100$
<sup>112</sup> Rh	-79730	40		3.4 s	0.4	(1 <sup>+</sup> )	15 99Lh01	T 1972	$\beta^- = 100$
<sup>112</sup> Rh <sup>m</sup>	-79390	60	340 70 BD	6.73 s	0.15	(6 <sup>+</sup> )	15 99Lh01	T 1987	$\beta^- = 100$
<sup>112</sup> Pd	-86322	7		21.04 h	0.17	0 <sup>+</sup>	15	1951	$\beta^- = 100$
<sup>112</sup> Ag	-86583.7	2.4		3.130 h	0.008	2(-)	15	1938	$\beta^- = 100$
<sup>112</sup> Cd	-90574.86	0.25		STABLE		0 <sup>+</sup>	15	1925	IS=24.13 21
<sup>112</sup> In	-87990	4		14.88 m	0.15	1 <sup>+</sup>	15	1947	$\beta^+ = 57.4 48; \beta^- = 42.6 48$
<sup>112</sup> In <sup>m</sup>	-87833	4	156.592 0.025	20.67 m	0.08	4 <sup>+</sup>	15	1953	IT=100
<sup>112</sup> In <sup>n</sup>	-87639	4	350.80 0.05	690 ns	50	(7 <sup>+</sup> )	15	1976	IT=100
<sup>112</sup> In <sup>p</sup>	-87376	4	613.82 0.06	2.81 $\mu$ s	0.03	8 <sup>-</sup>	15 87Eb02	J 1976	IT=100
<sup>112</sup> Sn	-88655.06	0.29		STABLE		0 <sup>+</sup>	15	1927	IS=0.97 1; $2\beta^+ ?$
<sup>112</sup> Sb	-81599	18		53.5 s	0.6	(3 <sup>+</sup> )	15	1959	$\beta^+ = 100$
<sup>112</sup> Sb <sup>m</sup>	-80773	18	825.9 0.4	536 ns	22	(8 <sup>-</sup> )	15	1976	IT=100
<sup>112</sup> Te	-77568	8		2.0 m	0.2	0 <sup>+</sup>	15	1976	$\beta^+ = 100$
<sup>112</sup> I	-67063	10		3.34 s	0.08	1 <sup>+</sup> #	15 78Ro19	D 1977	$\beta^+ \approx 100; \alpha = 0.0012; \beta^+ p = 0.88 10; \dots$
<sup>112</sup> Xe	-60026	8		2.7 s	0.8	0 <sup>+</sup>	15	1978	$\beta^+ \approx 100; \alpha = 1.2 8; \beta^+ p ?$
<sup>112</sup> Cs	-46290	90		490 $\mu$ s	30	1 <sup>+</sup> #	15	1994	$p \approx 100; \alpha < 0.26$
* <sup>112</sup> Zr	T : symmetrized from 15Lo04=30(+30-10)								
* <sup>112</sup> Tc <sup>m</sup>	E : 12Ka36=93.1(0.5) keV and 259.2(0.5) keV $\gamma$ rays in cascade to 2 <sup>+</sup> # ground-state								
* <sup>112</sup> Rh	T : 99Lh01=3.45(0.37) supersedes 91Jo11=2.1(0.3), 88Ay02=3.8(0.6) same group								
* <sup>112</sup> Rh <sup>m</sup>	T : supersedes 88Ay02=6.8(0.2) of same group								
* <sup>112</sup> Sn	T : >1.3 Zy for 0v- $\epsilon\epsilon$ transition to 0 <sub>3</sub> <sup>+</sup> state in <sup>112</sup> Cd								
* <sup>112</sup> I	D : ...; $\beta^+ \alpha = 0.104 12$								
<sup>113</sup> Nb	-40510#	400#		32 ms	4	5/2 <sup>+</sup> #	15	1997	$\beta^- = 100; \beta^-_n = 90\#; \beta^-_{2n} = 2\#$
<sup>113</sup> Mo	-52490#	300#		80 ms	2	3/2 <sup>+</sup> #	15	1994	$\beta^- = 100; \beta^-_n = 3\#$
<sup>113</sup> Tc	-62812	3		152 ms	8	5/2 <sup>+</sup> #	15	1992	$\beta^- = 100; \beta^-_n = 2.1 3$
<sup>113</sup> Tc <sup>m</sup>	-62698	3	114.4 0.5	527 ns	16	(5/2 <sup>-</sup> )	15 12Ka36	T 2010	IT=100
<sup>113</sup> Ru	-71870	40		800 ms	50	(1/2 <sup>+</sup> )	10	1988	$\beta^- = 100$
<sup>113</sup> Ru <sup>m</sup>	-71740	40	130 18	510 ms	30	(7/2 <sup>-</sup> )	10 98Ku17	E 1998	IT=?; $\beta^- = ?$
<sup>113</sup> Rh	-78768	7		2.80 s	0.12	(7/2 <sup>+</sup> )	10 93Pe11	J 1971	$\beta^- = 100$
<sup>113</sup> Pd	-83591	7		93 s	5	(5/2 <sup>+</sup> )	10	1954	$\beta^- = 100$
<sup>113</sup> Pd <sup>m</sup>	-83510	7	81.1 0.3	300 ms	100	(9/2 <sup>-</sup> )	10	1993	IT=100
<sup>113</sup> Ag	-87027	17		5.37 h	0.05	1/2 <sup>-</sup>	10	1949	$\beta^- = 100$
<sup>113</sup> Ag <sup>m</sup>	-86984	17	43.50 0.10	68.7 s	1.6	7/2 <sup>+</sup>	10	1958	IT=64 7; $\beta^- = 36 7$
<sup>113</sup> Cd	-89043.28	0.24		8.04 Py	0.05	1/2 <sup>+</sup>	10	1925	IS=12.22 12; $\beta^- = 100$
<sup>113</sup> Cd <sup>m</sup>	-88779.74	0.24	263.54 0.03	13.89 y	0.11	11/2 <sup>-</sup>	10 11Ko01	TD 1965	$\beta^- = 99.9036 19; IT = 0.0964 19$
<sup>113</sup> In	-89367.12	0.19		STABLE		9/2 <sup>+</sup>	10	1934	IS=4.29 5
<sup>113</sup> In <sup>m</sup>	-88975.42	0.19	391.699 0.003	1.6579 h	0.0004	1/2 <sup>-</sup>	10	1939	IT=100

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...									
<sup>113</sup> Sn	-88328.1	1.6		115.09 d	0.03			1939	$\beta^+=100$
<sup>113</sup> Sn <sup>m</sup>	-88250.7	1.6	77.389	0.019				1961	IT=91.1 23; $\beta^+=8.9$ 23
<sup>113</sup> Sb	-84417	17						1958	$\beta^+=100$
<sup>113</sup> Te	-78347	28						1974	$\beta^+=100$
<sup>113</sup> I	-71120	8						1977	$\beta^+=100$ ; $\alpha=3.310e-7$ ; $\beta^+\alpha$ ?
<sup>113</sup> Xe	-62204	7						1973	$\beta^+\approx 100$ ; $\alpha\approx 0.011$ ; $\beta^+p=7$ 4; $\beta^+\alpha\approx 0.007$ 4
<sup>113</sup> Xe <sup>m</sup>	-61800	7	403.6	1.4			13Pr01	ETJ 2013	IT=100
<sup>113</sup> Cs	-51765	9						1984	p=100
<sup>113</sup> Ba	-39780#	300#							p ?; $\alpha$ ?
* <sup>113</sup> Tc	J : 07Ku23 > 5/2								**
* <sup>113</sup> Tc <sup>m</sup>	T : symmetrized from 12Ka36=526(+16-15) ms								**
* <sup>113</sup> Ru <sup>m</sup>	E : above the 99 keV level and below 160 keV								**
* <sup>113</sup> Cd	T : from 07Be61=8.037(0.005)(0.05 systematics);								**
* <sup>113</sup> Cd	T : other 09Da03=8.00(0.11)(syt 0.24) outweighed								**
* <sup>113</sup> Cd <sup>m</sup>	T : average 11Ko01=13.97(0.13) 72Wa11=14.6(0.5) 65F102=13.6(0.2)								**
* <sup>113</sup> In <sup>m</sup>	T : 99.476 m 23								**
* <sup>113</sup> Xe	D : $\alpha=0.0024-0.0204\%$ from estimated limit for the reduced width, see 85Ti02								**
* <sup>113</sup> Xe	D : $\beta^+p$ and $\beta^+\alpha$ derived from $\beta^+p/\alpha=605(35)$ and $\beta^+p/\beta^+\alpha=500-1500$ in 85Ti02								**
<sup>114</sup> Nb	-35390#	500#						2010	$\beta^-=100$ ; $\beta^-n=50\#$ ; $\beta^-2n=6\#$
<sup>114</sup> Mo	-49810#	300#						1997	$\beta^-=100$ ; $\beta^-n=3\#$
<sup>114</sup> Tc	-58600	430						1994	$\beta^-=100$ ; $\beta^-n=6\#$
<sup>114</sup> Tc <sup>m</sup>	-58437	13	160	430				2011	$\beta^-$ ?; $\beta^-n=6\#$
<sup>114</sup> Ru	-70222	4						1991	$\beta^-=100$ ; $\beta^-n=0\#$ ; $\beta^-2n=0\#$
<sup>114</sup> Rh	-75710	70						1988	$\beta^-=100$
<sup>114</sup> Rh <sup>m</sup>	-75510#	170#	200#	150#				1987	$\beta^-=100$
<sup>114</sup> Pd	-83491	7						1958	$\beta^-=100$
<sup>114</sup> Ag	-84931	5						1958	$\beta^-=100$
<sup>114</sup> Ag <sup>m</sup>	-84732	7	199	5				1990	IT=100
<sup>114</sup> Cd	-90014.93	0.28						1925	IS=28.73 42; $2\beta^-$ ?
<sup>114</sup> In	-88569.8	0.3						1937	$\beta^-=99.50$ 15; $\beta^+=0.50$ 15
<sup>114</sup> In <sup>m</sup>	-88379.5	0.3	190.2682	0.0008				1939	IT=96.75 24; $\beta^+=3.25$ 24
<sup>114</sup> In	-88067.9	0.3	501.948	0.003				1958	IT=100
<sup>114</sup> In <sup>p</sup>	-87928.1	0.3	641.745	0.003				1975	IT=100
<sup>114</sup> Sn	-90559.723	0.029						1927	IS=0.66 1
<sup>114</sup> Sn <sup>m</sup>	-87472.35	0.08	3087.37	0.07				1980	IT=100
<sup>114</sup> Sb	-84497	22						1959	$\beta^+=100$
<sup>114</sup> Sb <sup>m</sup>	-84002	22	495.5	0.7				1973	IT=100
<sup>114</sup> Te	-81889	28						1968	$\beta^+=100$
<sup>114</sup> I	-72800#	150#						1977	$\beta^+=100$ ; $\beta^+p$ ?
<sup>114</sup> I <sup>m</sup>	-72530#	150#	265.9	0.5				1995	$\beta^+=91$ 2; IT=9 2
<sup>114</sup> Xe	-67086	11						1977	$\beta^+=100$
<sup>114</sup> Cs	-54680	70						1978	$\beta^+\approx 100$ ; $\alpha=0.018$ 6; $\beta^+p=8.7$ 13; $\beta^+\alpha=0.19$ 3
<sup>114</sup> Ba	-45910	100						1995	$\beta^+\approx 100$ ; $\beta^+p=20$ 10; $\alpha=0.9$ 3; $12C<0.0034$
* <sup>114</sup> Tc	T : others, might be mixture of ground-state and m : 15Lo04=120(10) 06Mo07=91(+62-35)								**
* <sup>114</sup> Tc	T : 99Wa09=150(30)								**
* <sup>114</sup> Ru	T : average 06Mo07=510(+69-65) 92Jo05=530(60) 91Le09=570(50)								**
* <sup>114</sup> In <sup>p</sup>	T : typo in NUBASE2012 : 4.3 $\mu$ s, should be 4.3 ns cf. 75Ra07 and ENSDF2012								**
* <sup>114</sup> I <sup>m</sup>	D : evaluated for NUBASE by J. Blachot, based on <sup>114</sup> I IT decay								**
* <sup>114</sup> Ba	T : average 16Ca33=380(+190-110) 97Ja12=430(+300-150)								**
<sup>115</sup> Nb	-31350#	500#						2010	$\beta^-=100$ ; $\beta^-n=60\#$ ; $\beta^-2n=1\#$
<sup>115</sup> Mo	-44750#	400#						2010	$\beta^-=100$ ; $\beta^-n=3\#$ ; $\beta^-2n=0.01\#$
<sup>115</sup> Tc	-56320	790						1994	$\beta^-=100$ ; $\beta^-n=20\#$
<sup>115</sup> Ru	-66190	90						1992	$\beta^-=100$ ; $\beta^-n=0.02\#$
<sup>115</sup> Ru <sup>m</sup>	-65940#	140#	250#	100#				2010	IT=100
<sup>115</sup> Rh	-74230	7						1988	$\beta^-=100$ ; $\beta^-n=0.05\#$
<sup>115</sup> Pd	-80426	14						1958	$\beta^-=100$
<sup>115</sup> Pd <sup>m</sup>	-80337	14	89.21	0.16				1987	$\beta^-=92.0$ 20; IT=8.0 20
<sup>115</sup> Ag	-84983	18						1949	$\beta^-=100$
<sup>115</sup> Ag <sup>m</sup>	-84942	18	41.16	0.10				1958	$\beta^-=79.0$ 3; IT=21.0 3
<sup>115</sup> Cd	-88084.5	0.7						1939	$\beta^-=100$
<sup>115</sup> Cd <sup>m</sup>	-87903.5	0.9	181.0	0.5				1959	$\beta^-\approx 100$ ; IT<0.003
<sup>115</sup> In	-89536.346	0.012						1924	IS=95.71 5; $\beta^-=100$
<sup>115</sup> In <sup>m</sup>	-89200.102	0.021	336.244	0.017				1961	IT=95.0 7; $\beta^-=5.0$ 7
<sup>115</sup> Sn	-90033.835	0.015						1927	IS=0.34 1
<sup>115</sup> Sn <sup>m</sup>	-89421.03	0.04	612.81	0.04				1967	IT=100
<sup>115</sup> Sn <sup>n</sup>	-89320.20	0.12	713.64	0.12				1958	IT=100
... A-group is continued on next page ...									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...										
<sup>115</sup> Sb	-87003	16	32.1 m	0.3	5/2 <sup>+</sup>	12	1958	$\beta^+=100$		
<sup>115</sup> Sb <sup>m</sup>	-84207	16	2796.26	0.09	159 ns	3	1977	IT=100		
<sup>115</sup> Te	-82063	28	*	5.8 m	0.2	7/2 <sup>+</sup>	12	1961	$\beta^+=100$	
<sup>115</sup> Te <sup>m</sup>	-82053	29	10	7	* 6.7 m	0.4	(1/2) <sup>+</sup>	12	GAu E 1974	$\beta^+\approx 100$ ; IT<0.06
<sup>115</sup> Te <sup>n</sup>	-81783	28	280.05	0.20	7.5 $\mu$ s	0.2	11/2 <sup>-</sup>	12	1972	IT=100
<sup>115</sup> I	-76338	29			1.3 m	0.2	5/2 <sup>+</sup> #	12	1969	$\beta^+=100$
<sup>115</sup> Xe	-68657	12			18 s	4	(5/2 <sup>+</sup> )	12	1969	$\beta^+=100$ ; $\beta^+p=0.34$ 6; $\alpha=0.0003$ 1
<sup>115</sup> Cs	-59700#	100#			1.4 s	0.8	9/2 <sup>+</sup> #	12	1978	$\beta^+=100$ ; $\beta^+p\approx 0.07$
<sup>115</sup> Ba	-49020#	200#			450 ms	50	5/2 <sup>+</sup> #	12	1997	$\beta^+=100$ ; $\beta^+p>15$
* <sup>115</sup> Ru	J : suggested in 111Ri07 from $\beta^-$ decay study								**	
* <sup>115</sup> Ru <sup>m</sup>	E : ENSDF2012 > 61.7 keV								**	
* <sup>115</sup> Cd <sup>m</sup>	J : measured magnetic moment and L(d,p)=5								**	
* <sup>115</sup> Te <sup>m</sup>	E : less than 20 keV, from ENSDF								**	
<sup>116</sup> Mo	-41500#	500#			32 ms	4	0 <sup>+</sup>	15	2010	$\beta^-=100$ ; $\beta^-n=10\#$ ; $\beta^-2n=0\#$
<sup>116</sup> Tc	-51460#	300#			57 ms	3	2 <sup>+</sup> #	15	1997	$\beta^-=100$ ; $\beta^-n=20\#$ ; $\beta^-2n=0.05\#$
<sup>116</sup> Ru	-64069	4			204 ms	6	0 <sup>+</sup>	15	1994	$\beta^-=100$ ; $\beta^-n=0.1\#$
<sup>116</sup> Rh	-70740	70			* 685 ms	39	1 <sup>+</sup>	10	06Mo07 TD 1970	$\beta^-=100$ ; $\beta^-n<2.1$
<sup>116</sup> Rh <sup>m</sup>	-70540#	170#	200#	150#	* 570 ms	50	(6 <sup>-</sup> )	10	1987	$\beta^-=100$
<sup>116</sup> Pd	-79832	7			11.8 s	0.4	0 <sup>+</sup>	10	1970	$\beta^-=100$
<sup>116</sup> Ag	-82543	3			3.83 m	0.08	(0 <sup>-</sup> )	10	1958	$\beta^-=100$
<sup>116</sup> Ag <sup>m</sup>	-82495	3	47.90	0.10	20 s	1	(3 <sup>+</sup> )	10	2005	$\beta^-=93.0$ ; IT=7.0
<sup>116</sup> Ag <sup>n</sup>	-82413	3	129.80	0.22	9.3 s	0.3	(6 <sup>-</sup> )	10	1970	$\beta^-=92.0$ ; IT=8.0
<sup>116</sup> Cd	-88712.48	0.16			28.7 Ey	1.3	0 <sup>+</sup>	10	15Ba11 T 1925	IS=7.49 18; $2\beta^-=100$
<sup>116</sup> In	-88249.75	0.22			14.10 s	0.03	1 <sup>+</sup>	10	13Wr01 D 1937	$\beta^-\approx 100$ ; $\epsilon=0.0237$ 43
<sup>116</sup> In <sup>m</sup>	-88122.48	0.22	127.267	0.006	54.29 m	0.17	5 <sup>+</sup>	10	1945	$\beta^-=100$
<sup>116</sup> In <sup>n</sup>	-87960.09	0.22	289.660	0.006	2.18 s	0.04	8 <sup>-</sup>	10	1950	IT=100
<sup>116</sup> Sn	-91525.97	0.10			STABLE		0 <sup>+</sup>	10	1922	IS=14.54 9
<sup>116</sup> Sn <sup>m</sup>	-89160.00	0.10	2365.975	0.021	348 ns	19	5 <sup>-</sup>	10	1964	IT=100
<sup>116</sup> Sn <sup>n</sup>	-87978.81	0.20	3547.16	0.17	833 ns	30	10 <sup>+</sup>	10	1978	IT=100
<sup>116</sup> Sb	-86822	5			15.8 m	0.8	3 <sup>+</sup>	10	1949	$\beta^+=100$
<sup>116</sup> Sb <sup>m</sup>	-86728	5	93.99	0.05	194 ns	4	1 <sup>+</sup>	10	1976	IT=100
<sup>116</sup> Sb <sup>n</sup>	-86440	40	390	40	BD 60.3 m	0.6	8 <sup>-</sup>	10	1949	$\beta^+=100$
<sup>116</sup> Te	-85269	28			2.49 h	0.04	0 <sup>+</sup>	10	1958	$\beta^+=100$
<sup>116</sup> I	-77490	100			2.91 s	0.15	1 <sup>+</sup>	10	1976	$\beta^+=100$
<sup>116</sup> I <sup>m</sup>	-77060	100	430.4	0.5	3.27 $\mu$ s	0.16	(7 <sup>-</sup> )	10	1990	IT=100
<sup>116</sup> Xe	-73047	13			59 s	2	0 <sup>+</sup>	10	1969	$\beta^+=100$
<sup>116</sup> Cs	-62040#	100#			* 700 ms	40	(1 <sup>+</sup> )	10	77Bo28 D 1975	$\beta^+=100$ ; $\beta^+p=0.28$ 7; $\beta^+\alpha=0.049$ 25
<sup>116</sup> Cs <sup>m</sup>	-61940#	120#	100#	60#	* 3.85 s	0.13	4 <sup>+</sup> , 5, 6	10	1975	$\beta^+=100$ ; $\beta^+p=0.51$ 15; $\beta^+\alpha=0.008$ 2
<sup>116</sup> Ba	-54580#	200#			1.3 s	0.2	0 <sup>+</sup>	10	1997	$\beta^+=100$ ; $\beta^+p=3$ 1
<sup>116</sup> La	-40650#	310#			10# ms			10		$\beta^+?$ ; $\beta^+p?$ ; $p?$
* <sup>116</sup> Rh	T : average 06Mo07=688(+52-50) 88Ay02=680(60) D : $\beta^-n$ limit from 06Mo07								**	
* <sup>116</sup> Ag	T : 230(5) s								**	
* <sup>116</sup> Cd	T : also 0v- $\beta\beta$ 96Ar36>5000Ey and Majoron 96Ar36>1200Ey 98Da23>1200Ey								**	
* <sup>116</sup> In	D : average 13Wr01=0.0246(44stat)(39syst) 98Bh04=0.0227 63								**	
* <sup>116</sup> In	T : also 13Wr01=14.9(0.8)								**	
* <sup>116</sup> Cs	D : from 77Bo28; ENSDF2010 erroneously gives $\beta^+p=2.8$ 7								**	
* <sup>116</sup> La	T : half-life estimate is for $\beta^+$ decay; no p-decay within 20 $\mu$ s-20ms								**	
<sup>117</sup> Mo	-36170#	500#			22 ms	5	3/2 <sup>+</sup> #	15	2010	$\beta^-=100$ ; $\beta^-n=10\#$ ; $\beta^-2n=0.2\#$
<sup>117</sup> Tc	-48380#	400#			44.5 ms	3.0	5/2 <sup>+</sup> #	15	1997	$\beta^-=100$ ; $\beta^-n=30\#$ ; $\beta^-2n=0\#$
<sup>117</sup> Ru	-59490	430			151 ms	3	3/2 <sup>+</sup> #	15	1994	$\beta^-=100$ ; $\beta^-n=0.5\#$
<sup>117</sup> Ru <sup>m</sup>	-59310	430	185.0	0.4	2.49 $\mu$ s	0.6		15	2012	IT=100
<sup>117</sup> Rh	-68897	9			421 ms	30	7/2 <sup>+</sup> #	11	06Mo07 TD 1991	$\beta^-=100$ ; $\beta^-n<7.6$
<sup>117</sup> Pd	-76424	7			4.3 s	0.3	(3/2 <sup>+</sup> )	11	04Ur04 J 1968	$\beta^-=100$
<sup>117</sup> Pd <sup>m</sup>	-76221	7	203.3	0.3	19.1 ms	0.7	(9/2 <sup>-</sup> )	11	04Ur04 J 1990	IT=100
<sup>117</sup> Ag	-82182	14			73.6 s	1.4	1/2 <sup>-</sup> #	11	1958	$\beta^-=100$
<sup>117</sup> Ag <sup>m</sup>	-82153	14	28.6	0.2	5.34 s	0.05	7/2 <sup>+</sup> #	11	1990	$\beta^-=94.0$ 15; IT=6.0 15
<sup>117</sup> Cd	-86418.4	1.0			2.49 h	0.04	1/2 <sup>+</sup>	11	1939	$\beta^-=100$
<sup>117</sup> Cd <sup>m</sup>	-86282.0	1.0	136.4	0.2	3.36 h	0.05	11/2 <sup>-</sup>	11	13Yo02 J 1966	$\beta^-\approx 100$ ; IT $\approx 0$
<sup>117</sup> In	-88943	5			43.2 m	0.3	9/2 <sup>+</sup>	11	1937	$\beta^-=100$
<sup>117</sup> In <sup>m</sup>	-88628	5	315.303	0.011	116.2 m	0.3	1/2 <sup>-</sup>	11	1940	$\beta^-=52.9$ 15; IT=47.1 15
<sup>117</sup> Sn	-90397.8	0.5			STABLE		1/2 <sup>+</sup>	11	1923	IS=7.68 7
<sup>117</sup> Sn <sup>m</sup>	-90083.2	0.5	314.58	0.04	14.00 d	0.05	11/2 <sup>-</sup>	12	1950	IT=100
<sup>117</sup> Sn <sup>n</sup>	-87991.4	0.6	2406.4	0.4	1.75 $\mu$ s	0.07	(19/2 <sup>+</sup> )	11	1979	IT=100
... A-group is continued on next page ...										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>117</sup> Sb	-88640	8	2.80 h	0.01	5/2 <sup>+</sup>	11	1947	$\beta^+$ =100
<sup>117</sup> Sb <sup>m</sup>	-85509	8	3130.76	0.19	355 $\mu$ s	17	1970	IT=100
<sup>117</sup> Sb <sup>n</sup>	-85409	8	3230.7	0.2	290 ns	5	1987	IT=100
<sup>117</sup> Te	-85095	13			62 m	2	1958	$\epsilon$ =75 1; $e^+$ =25 1
<sup>117</sup> Te <sup>m</sup>	-84799	13	296.1	0.5	103 ms	3	99Mo30 J	IT ?
<sup>117</sup> I	-80436	26			2.22 m	0.04	1969	$\beta^+$ =100; $e^+$ $\approx$ 77
<sup>117</sup> Xe	-74185	10			61 s	2	1969	$\beta^+$ =100; $\beta^+$ p=0.0029 6
<sup>117</sup> Cs	-66490	60			8.4 s	0.6	1972	$\beta^+$ =100
<sup>117</sup> Cs <sup>m</sup>	-66340#	100#	150#	80#	6.5 s	0.4	1978	$\beta^+$ =100
<sup>117</sup> Cs <sup>s</sup>	-66440	80	50	50	R=?	spmix		
<sup>117</sup> Ba	-57460	250			1.75 s	0.07	97Ja12 D	$\beta^+$ =100; $\beta^+$ p=13 3; $\beta^+$ $\alpha$ =0.024 8
<sup>117</sup> La	-46470#	200#			21.7 ms	1.8	11Li28 TJ	p=?; $\beta^+$ =6.1#; $\beta^+$ p ?
<sup>117</sup> La <sup>m</sup>			non existent	RN	10 ms	5	01So02 I	
* <sup>117</sup> Ru <sup>m</sup>	T: symmetrized from 12Ka36=2.487(+0.058-0.055)							
* <sup>117</sup> Rh	T: average 06Mo07=394(+47-43) 91Pe10=440(40)							
* <sup>117</sup> Ag	T: symmetrized from 72.8(+2.0-0.7)							
* <sup>117</sup> Ag <sup>m</sup>	J: E3 to ground-state 1/2 <sup>-</sup> #							
* <sup>117</sup> Ba	D: $\beta^+$ p from 97Ja12. $\beta^+$ p/ $\beta^+$ $\alpha$ =350-1200 from 85Ti02 yields $\beta^+$ $\alpha$ =0.011%-0.037%							
* <sup>117</sup> La	T: average 11Li28=20.1(2.5) 01Ma69=24(3) 01So02=22(5)							
* <sup>117</sup> La <sup>m</sup>	I: reported in 01So02 with E=121(10) keV. Not observed in 11Li28							
<sup>118</sup> Mo	-32630#	500#			21 ms	6	15Lo04 TD	$\beta^-$ =100; $\beta^-$ n=10#; $\beta^-$ 2n=0.08#
<sup>118</sup> Tc	-43790#	400#			30 ms	4	2010	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=0.6#
<sup>118</sup> Ru	-57260#	200#			99 ms	3	1994	$\beta^-$ =100; $\beta^-$ n=1#
<sup>118</sup> Rh	-64887	24			284 ms	9	06 15Lo04 T	$\beta^-$ =100; $\beta^-$ n=3.1 14
<sup>118</sup> Pd	-75388.7	2.5			1.9 s	0.1	1969	$\beta^-$ =100
<sup>118</sup> Ag	-79553.8	2.5			3.76 s	0.15	93Ja03 J	$\beta^-$ =100
<sup>118</sup> Ag <sup>m</sup>	-79508.0	2.5	45.79	0.09	0.1 $\mu$ s		1989	IT=100
<sup>118</sup> Ag <sup>n</sup>	-79426.2	2.5	127.63	0.10	2.0 s	0.2	1971	$\beta^-$ =59; IT=41
<sup>118</sup> Ag <sup>p</sup>	-79274.4	2.5	279.37	0.20	0.1 $\mu$ s		1989	IT=100
<sup>118</sup> Cd	-86702	20			50.3 m	0.2	1961	$\beta^-$ =100
<sup>118</sup> In	-87228	8			5.0 s	0.5	1949	$\beta^-$ =100
<sup>118</sup> In <sup>m</sup>	-87130#	50#	100#	50#	4.364 m	0.007	94It.A T	$\beta^-$ =100
<sup>118</sup> In <sup>n</sup>	-86990#	50#	240#	50#	8.5 s	0.3	1969	IT=98.6 3; $\beta^-$ =1.4 3
<sup>118</sup> Sn	-91652.9	0.5			STABLE		1924	IS=24.22 9
<sup>118</sup> Sn <sup>m</sup>	-89078.0	0.5	2574.91	0.04	230 ns	10	1961	IT=100
<sup>118</sup> Sn <sup>n</sup>	-88544.8	0.5	3108.06	0.22	2.52 $\mu$ s	0.06	11Fo15 J	IT=100
<sup>118</sup> Sb	-87996	3			3.6 m	0.1	1947	$\beta^+$ =100
<sup>118</sup> Sb <sup>m</sup>	-87945	3	50.814	0.021	20.6 $\mu$ s	0.6	1975	IT=100
<sup>118</sup> Sb <sup>n</sup>	-87746	5	250	6	5.00 h	0.02	1947	$\beta^+$ =100
<sup>118</sup> Te	-87697	18			6.00 d	0.02	1948	$\epsilon$ =100
<sup>118</sup> I	-80971	20			13.7 m	0.5	1957	$\beta^+$ =100
<sup>118</sup> I <sup>m</sup>	-80782	20	188.8	0.7	8.5 m	0.5	03Mo36 E	$\beta^+$ $\approx$ 100; IT=?
<sup>118</sup> Xe	-78079	10			3.8 m	0.9	1965	$\beta^+$ =100
<sup>118</sup> Cs	-68409	13			14 s	2	1969	$\beta^+$ =100; $\beta^+$ p=0.021 14; $\beta^+$ $\alpha$ =0.0012 5
<sup>118</sup> Cs <sup>m</sup>	-68310#	60#	100#	60#	17 s	3	93Be46 J	$\beta^+$ =100; $\beta^+$ p=0.021 14; $\beta^+$ $\alpha$ =0.0012 5
<sup>118</sup> Cs <sup>s</sup>	-68404	12	5	4	R < 0.1	spmix		
<sup>118</sup> Ba	-62350#	200#			5.2 s	0.2	06 97Ja12 T	$\beta^+$ =100
<sup>118</sup> La	-49560#	300#			200# ms			$\beta^+$ ?; $\beta^+$ p ?
* <sup>118</sup> Mo	T: symmetrized from 15Lo04=19(+7-4)							
* <sup>118</sup> Rh	T: average 15Lo04=285(10) 06Mo07=266(+22-21) 00Jo18=310(30)							
* <sup>118</sup> Rh	J: from 00Jo18							
* <sup>118</sup> In <sup>n</sup>	E: 138.2(0.5) keV above <sup>118</sup> In <sup>m</sup> , from ENSDF							
* <sup>118</sup> I <sup>m</sup>	E: from a least-squares fit to level scheme of 03Mo36							
* <sup>118</sup> Cs	D: derived from $\beta^+$ p=0.042(6)%, $\beta^+$ $\alpha$ =0.0024(4)% for mixture of ground-state and isomer.							
* <sup>118</sup> Cs	D: Replaced by uniform distributions from zero to values for each isomer							
<sup>119</sup> Tc	-40370#	500#			22 ms	3	2010	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=0.1#
<sup>119</sup> Ru	-52560#	300#			69.5 ms	2.0	1997	$\beta^-$ =100; $\beta^-$ n=3#; $\beta^-$ 2n=0#
<sup>119</sup> Ru <sup>m</sup>	-52330#	300#	227.1	0.7	384 ns	22	2012	IT=100
<sup>119</sup> Rh	-62823	9			190 ms	6	09 15Lo04 T	$\beta^-$ =100; $\beta^-$ n=6.4 16
<sup>119</sup> Pd	-71408	8			920 ms	80	06Mo07 TD	$\beta^-$ =100; $\beta^-$ n=0#
<sup>119</sup> Pd <sup>m</sup>	-71110#	150#	300#	150#	3# ms			IT ?; $\beta^-$ ?
<sup>119</sup> Ag	-78646	15			6.0 s	0.5	1975	$\beta^-$ =100
<sup>119</sup> Ag <sup>m</sup>	-78626#	25#	20#	20#	2.1 s	0.1	1975	$\beta^-$ =100
<sup>119</sup> Cd	-83980	40			2.69 m	0.02	09 13Yo02 J	$\beta^-$ =100
<sup>119</sup> Cd <sup>m</sup>	-83830	40	146.54	0.11	2.20 m	0.02	09 13Yo02 J	$\beta^-$ =100
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)				
... A-group continued ...													
<sup>119</sup> In	-87699	7		2.4 m	0.1	9/2 <sup>+</sup>	09	1949	$\beta^-$ =100				
<sup>119</sup> In <sup>m</sup>	-87388	7	311.37	0.03		18.0 m	0.3	1/2 <sup>-</sup>	09	$\beta^-$ =95.6; IT=4.4			
<sup>119</sup> In <sup>n</sup>	-87045	7	654.27	0.07		130 ns	15	(3/2) <sup>+</sup>	09	IT=100			
<sup>119</sup> In <sup>p</sup>	-85042	7	2656.9	1.8		240 ns	25	(25/2 <sup>+</sup> )	09	IT=100			
<sup>119</sup> Sn	-90065.0	0.7				STABLE		1/2 <sup>+</sup>	09	IS=8.59 4			
<sup>119</sup> Sn <sup>m</sup>	-89975.5	0.7	89.531	0.013		293.1 d	0.7	11/2 <sup>-</sup>	09	IT=100			
<sup>119</sup> Sn <sup>n</sup>	-87938.0	1.2	2127.0	1.0		9.6 $\mu$ s	1.2	(19/2 <sup>+</sup> )	09	IT=100			
<sup>119</sup> Sn <sup>p</sup>	-87696.0	0.8	2369.0	0.3		96 ns	9	23/2 <sup>+</sup>	09	IT=100			
<sup>119</sup> Sb	-89474	8				38.19 h	0.22	5/2 <sup>+</sup>	09	$\epsilon$ =100			
<sup>119</sup> Sb <sup>m</sup>	-86920	8	2553.6	0.3		130 ns	3	19/2 <sup>-</sup>	09	IT=100			
<sup>119</sup> Sb <sup>n</sup>	-86622	11	2852	7		850 ms	90	27/2 <sup>+</sup> #	09	IT=100			
<sup>119</sup> Te	-87181	8				16.05 h	0.05	1/2 <sup>+</sup>	09	$\epsilon$ =97.94 5; $\epsilon^+$ =2.06 5			
<sup>119</sup> Te <sup>m</sup>	-86920	8	260.96	0.05		4.70 d	0.04	11/2 <sup>-</sup>	09	$\epsilon$ =99.59 4; $\epsilon^+$ =0.41 4; IT<0.008			
<sup>119</sup> I	-83766	28				19.1 m	0.4	5/2 <sup>+</sup>	09	$\epsilon^+$ =51 4; $\epsilon$ =49 4			
<sup>119</sup> Xe	-78794	10				5.8 m	0.3	5/2 <sup>(+)</sup>	09	$\epsilon^+$ =79 5; $\epsilon$ =21 5			
<sup>119</sup> Cs	-72305	14		*		43.0 s	0.2	9/2 <sup>+</sup>	09	$\beta^+$ =100; $\beta^+$ $\alpha$ <2e-6			
<sup>119</sup> Cs <sup>m</sup>	-72260#	30#	50#	30#	*	30.4 s	0.1	3/2 <sup>(+)</sup>	09	$\beta^+$ =100			
<sup>119</sup> Cs <sup>s</sup>	-72289	9	16	11		R = .5	.25	spmix					
<sup>119</sup> Ba	-64590	200				5.4 s	0.3	(5/2 <sup>+</sup> )	09	$\beta^+$ =100; $\beta^+$ p=25 2			
<sup>119</sup> La	-54790#	300#				1#	s	11/2 <sup>-</sup> #		$\beta^+$ ?			
<sup>119</sup> Ce	-43940#	500#				200#	ms	5/2 <sup>+</sup> #		$\beta^+$ ?; $\beta^+$ p ?			
* <sup>119</sup> Ru <sup>m</sup>	T : symmetrized from 12Ka36=383(+22-21)												
* <sup>119</sup> Pd	T : average 06Mo07=918(111) 91Pe04=920(130)												
* <sup>119</sup> Ag <sup>m</sup>	E : estimated from 7/2 <sup>+</sup> level in isotopes <sup>113</sup> Ag=43 <sup>115</sup> Ag=41 <sup>117</sup> Ag=28												
* <sup>119</sup> Sb <sup>n</sup>	E : estimated less than 20 keV above 2841.7 level												
<sup>120</sup> Tc	-35520#	500#				21 ms	5		15	2010	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=2#		
<sup>120</sup> Ru	-50010#	400#				45 ms	2	0 <sup>+</sup>	15	2010	$\beta^-$ =100; $\beta^-$ n=4#		
<sup>120</sup> Rh	-58820#	200#				129.6 ms	4.2		13	15Lo04 T	1994	$\beta^-$ =100; $\beta^-$ n<5.4; $\beta^-$ 2n=0#	
<sup>120</sup> Rh <sup>m</sup>	-58660#	200#	157.2	0.7		295 ns	16		13	12Ka36 ETD	2012	IT=100	
<sup>120</sup> Pd	-70280.1	2.3				492 ms	33	0 <sup>+</sup>	14	06Mo07 TD	1993	$\beta^-$ =100; $\beta^-$ n<0.7	
<sup>120</sup> Ag	-75652	4				1.52 s	0.07	4 <sup>(+)</sup>	02	12Ba58 TJ	1971	$\beta^-$ =100; $\beta^-$ n<0.003	
<sup>120</sup> Ag <sup>m</sup>	-75650#	50#	0#	50#		940 ms	100	(0 <sup>-</sup> , 1 <sup>-</sup> )		12Ba58 TJD	2012	$\beta^-$ =?; IT ?	
<sup>120</sup> Ag <sup>n</sup>	-75449	4	203.0	0.2		384 ms	22	7 <sup>(-)</sup>	02	12Ba58 EJD	1971	IT=68 10; $\beta^-$ =32 5	
<sup>120</sup> Cd	-83957	4				50.80 s	0.21	0 <sup>+</sup>	02		1973	$\beta^-$ =100	
<sup>120</sup> In	-85730	40			*	3.08 s	0.08	1 <sup>+</sup>	02		1958	$\beta^-$ =100	
<sup>120</sup> In <sup>m</sup>	-85680#	50#	50#	60#	*	&	46.2 s	0.8	(5) <sup>+</sup>	02	13Ma15 J	1960	$\beta^-$ =100
<sup>120</sup> In <sup>n</sup>	-85430#	200#	300#	200#	*	&	47.3 s	0.5	8 <sup>(-)</sup>	02	79Fo10 J	1960	$\beta^-$ =100
<sup>120</sup> Sn	-91098.4	0.9				STABLE		0 <sup>+</sup>	02		1926	IS=32.58 9	
<sup>120</sup> Sn <sup>m</sup>	-88616.8	0.9	2481.63	0.06		11.8 $\mu$ s	0.5	7 <sup>-</sup>	02		1960	IT=100	
<sup>120</sup> Sn <sup>n</sup>	-88196.2	0.9	2902.22	0.22		6.26 $\mu$ s	0.11	10 <sup>+</sup>	02	FGK128 J	1987	IT=100	
<sup>120</sup> Sb	-88418	7			*	15.89 m	0.04	1 <sup>+</sup>	02		1937	$\beta^+$ =100	
<sup>120</sup> Sb <sup>m</sup>	-88420#	100#	0#	100#	*	5.76 d	0.02	8 <sup>-</sup>	02		1958	$\beta^+$ =100	
<sup>120</sup> Sb <sup>n</sup>	-88340	7	78.16	0.05		246 ns	2	(3 <sup>+</sup> )	02		1976	IT=100	
<sup>120</sup> Sb <sup>p</sup>	-86090	7	2328.3	0.6		400 ns	8		02		1983	IT=100	
<sup>120</sup> Te	-89368	3				STABLE		0 <sup>+</sup>	02		1936	IS=0.09 1; 2 $\beta^+$ ?	
<sup>120</sup> I	-83753	15				81.67 m	0.18	2 <sup>-</sup>	02	06Ph01 T	1957	$\beta^+$ =100	
<sup>120</sup> I <sup>m</sup>	-83680	15	72.61	0.09		242 ns	5	3 <sup>+</sup>	02	11Mo27 TJ	1974	IT=100	
<sup>120</sup> I <sup>n</sup>	-83433	21	320	15		53 m	4	(7 <sup>-</sup> )	02		1967	$\beta^+$ =100	
<sup>120</sup> Xe	-82172	12				46.0 m	0.6	0 <sup>+</sup>	02	06Ph01 T	1965	$\beta^+$ =100	
<sup>120</sup> Cs	-73889	10			*	60.4 s	0.6	2 <sup>(+)</sup>	02	06Ph01 T	1969	$\beta^+$ =100; $\beta^+$ $\alpha$ <2.0e-5 4; $\beta^+$ p<7e-6 3	
<sup>120</sup> Cs <sup>m</sup>	-73790#	60#	100#	60#	*	57 s	6	(7 <sup>-</sup> )	02	75Ho09 D	1977	$\beta^+$ =100; $\beta^+$ $\alpha$ <2.0e-5 4; $\beta^+$ p<7e-6 3	
<sup>120</sup> Cs <sup>s</sup>	-73884	9	5	4		R < 0.1		spmix					
<sup>120</sup> Ba	-68890	300				24 s	2	0 <sup>+</sup>	02	92Xu04 T	1974	$\beta^+$ =100	
<sup>120</sup> La	-57570#	300#				2.8 s	0.2		02		1984	$\beta^+$ =100; $\beta^+$ p=?	
<sup>120</sup> Ce	-49600#	500#				250#	ms	0 <sup>+</sup>				$\beta^+$ ?; $\beta^+$ p ?	
* <sup>120</sup> Rh	T : average 15Lo04=131(5) 06Mo07=136(+14-13)) 04Wa26=120(10)												
* <sup>120</sup> Rh <sup>m</sup>	E : 12Ka36=59.1(0.5) and 98.1(0.5) $\gamma$ rays in cascade to ground-state												
* <sup>120</sup> Rh <sup>n</sup>	T : symmetrized from 12Ka36=294(+16-15)												
* <sup>120</sup> Pd	D : 2v- $\beta\beta$ decay estimated 150(60)Ey												
* <sup>120</sup> Ag	T : not used 83Re05=1.25(0.03) 71Fo22=1.17(0.05) D : from 93Ru01												
* <sup>120</sup> Ag <sup>n</sup>	T : average 12Ba58=440(50) 03Wa13=400(30) 71Fo22=320(40)												
* <sup>120</sup> Sn <sup>n</sup>	J : E2 (from intensity balance) to 8 <sup>+</sup> I(354.9)/I(65.7)=8.7(1.0)												
* <sup>120</sup> I	T : average 06Ph01=82.1(0.6) 00Ho19=81.7(0.2) 65An05=81.0(0.6)												
* <sup>120</sup> I <sup>m</sup>	T : average 11Mo27=244(5) 74Mu10=228(15)												
* <sup>120</sup> Cs	T : average 06Ph01=60.0(7) 93Al03=60(2) 77Ge03=64(3) 69Ch18=61.3(1.4)												
* <sup>120</sup> Cs	D : isomers not distinguished by 75Ho09 in $\beta^+$ $\alpha$ and $\beta^+$ p. Values replaced												
* <sup>120</sup> Cs	D : by upper limits for both (see ENSDF evaluation of <sup>118</sup> Cs)												



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>121</sup> Tc	-31780#	500#	22 ms	6	5/2 <sup>+</sup> #	15	2015	$\beta^- = 100; \beta^- n = 50\#; \beta^- 2n = 0.7\#$		
<sup>121</sup> Ru	-45050#	400#	29 ms	2	3/2 <sup>+</sup> #	15	2010	$\beta^- = 100; \beta^- n = 6\#; \beta^- 2n = 0\#$		
<sup>121</sup> Rh	-56250	620	76 ms	5	7/2 <sup>+</sup> #	10	15Lo04 T 1994	$\beta^- = 100; \beta^- n = 7\#$		
<sup>121</sup> Pd	-66182	3	290 ms	1	3/2 <sup>+</sup> #	10	15Lo04 T 1994	$\beta^- = 100; \beta^- n < 0.8$		
<sup>121</sup> Pd <sup>m</sup>	-66047	3	135.5	0.5	460 ns	90	12Ka36 ETD 2007	IT=100; $\beta^- ?; \beta^- n = 0\#$		
<sup>121</sup> Pd <sup>n</sup>	-66022	15	160	15	460 ns	90	12Ka36 ETD 2007	IT=100; $\beta^- ?; \beta^- n = 0\#$		
<sup>121</sup> Ag	-74403	12	780 ms	20	7/2 <sup>+</sup> #	10	1982	$\beta^- = 100; \beta^- n = 0.080$ 13		
<sup>121</sup> Ag <sup>m</sup>	-74383#	23#	20#	20#	*	200# ms	1/2 <sup>-</sup> #	$\beta^- ?; IT ?$		
<sup>121</sup> Cd	-81073.8	1.9	13.5 s	0.3	3/2 <sup>+</sup>	10	13Yo02 J 1965	$\beta^- = 100$		
<sup>121</sup> Cd <sup>m</sup>	-80858.9	1.9	214.86	0.15	8.3 s	0.8	11/2 <sup>-</sup>	10 13Yo02 J 1982	$\beta^- = 100$	
<sup>121</sup> In	-85836	27	23.1 s	0.6	9/2 <sup>+</sup>	10	1960	$\beta^- = 100$		
<sup>121</sup> In <sup>m</sup>	-85522	27	313.68	0.07	3.88 m	0.10	1/2 <sup>-</sup>	10 1974	$\beta^- = 98.8$ 2; IT=1.2 2	
<sup>121</sup> In <sup>n</sup>	-83388	27	2448	1	17 $\mu$ s	2	(21/2 <sup>-</sup> )	10 10Re01 ETJ 2010	IT=100	
<sup>121</sup> Sn	-89197.3	1.0	27.03 h	0.04	3/2 <sup>+</sup>	10	1948	$\beta^- = 100$		
<sup>121</sup> Sn <sup>m</sup>	-89191.0	1.0	6.31	0.06	43.9 y	0.5	11/2 <sup>-</sup>	10 1962	IT=77.6 20; $\beta^- = 22.4$ 20	
<sup>121</sup> Sn <sup>n</sup>	-87198.5	1.3	1998.8	0.9	5.3 $\mu$ s	0.5	(19/2 <sup>+</sup> )	10 1995	IT=100	
<sup>121</sup> Sn <sup>p</sup>	-86976.2	1.1	2221.1	0.4	520 ns	50	(23/2 <sup>+</sup> )	10 16Is03 EJT 2012	IT=100	
<sup>121</sup> Sn <sup>q</sup>	-86362.7	2.1	2834.6	1.8	167 ns	25	(27/2 <sup>-</sup> )	10 1995	IT=100	
<sup>121</sup> Sb	-89600.3	2.6	STABLE		5/2 <sup>+</sup>	10	1922	IS=57.21 5		
<sup>121</sup> Sb <sup>m</sup>	-86859	12	2741	12	179 $\mu$ s	6	(25/2 <sup>+</sup> )	10 09Wa02 EJ 2008	IT=100	
<sup>121</sup> Te	-88546	26	19.17 d	0.04	1/2 <sup>+</sup>	10	1939	$\beta^+ = 100$		
<sup>121</sup> Te <sup>m</sup>	-88252	26	293.974	0.022	164.2 d	0.8	11/2 <sup>-</sup>	10 1940	IT=88.6 11; $\beta^+ = 11.4$ 11	
<sup>121</sup> I	-86251	5	2.12 h	0.01	5/2 <sup>+</sup>	10	1950	$\beta^+ = 100$		
<sup>121</sup> I <sup>m</sup>	-83874	5	2376.9	0.4	9.0 $\mu$ s	1.4	10	1982	IT=100	
<sup>121</sup> Xe	-82481	10	40.1 m	2.0	5/2 <sup>(+)</sup>	10	1952	$\beta^+ = 100$		
<sup>121</sup> Cs	-77102	14	155 s	4	3/2 <sup>(+)</sup>	10	1969	$\beta^+ = 100$		
<sup>121</sup> Cs <sup>m</sup>	-77034	14	68.5	0.3	122 s	3	9/2 <sup>(+)</sup>	10 1981	$\beta^+ = 83; IT = 17$	
<sup>121</sup> Cs <sup>x</sup>	-77056	16	46	8	R = 21	spmix				
<sup>121</sup> Ba	-70740	140	29.7 s	1.5	5/2 <sup>(+)</sup>	10	75Bo11 D 1975	$\beta^+ = 100; \beta^+ p = 0.02$ 1		
<sup>121</sup> La	-62190#	300#	5.3 s	0.2	11/2 <sup>-</sup> #	10	1988	$\beta^+ = 100; \beta^+ p ?$		
<sup>121</sup> Ce	-52690#	400#	1.1 s	0.1	(5/2) <sup>(+)</sup> #	10	1997	$\beta^+ = 100; \beta^+ p \approx 1$		
<sup>121</sup> Pr	-41420#	500#	12 ms	5	(3/2)	10	2005	p $\approx$ 100		
* <sup>121</sup> Pd <sup>m</sup>	T : symmetrized from 12Ka36=460(+85-92) E : other 07To23=135(3) keV									
* <sup>121</sup> Pd <sup>n</sup>	T : symmetrized from 12Ka36=463(+83-94) and assuming two cascading isomers									
* <sup>121</sup> Pd <sup>n</sup>	E : x keV above 121Pd <sup>m</sup> , x below energy threshold 50 keV									
* <sup>121</sup> In <sup>n</sup>	T : other 02Lu15=350(50) ns, assigned J=(25/2 <sup>+</sup> ); further studies are needed									
* <sup>121</sup> Sn <sup>n</sup>	E : <sup>121</sup> Sn <sup>n</sup> =1998.8(0.9) and <sup>121</sup> Sn <sup>p</sup> =2834.6(1.8) are from ENSDF2000, not in 2010									
* <sup>121</sup> Sb <sup>m</sup>	E : above 2720.9 level and <2761; other 08Jo03=2721.1 + x with x < 60 or x < 80									
* <sup>121</sup> Pr	T : symmetrized from 10(+6-3)									
<sup>122</sup> Ru	-42150#	500#	25 ms	1	0 <sup>+</sup>	15	2010	$\beta^- = 100; \beta^- n = 7\#; \beta^- 2n = 0\#$		
<sup>122</sup> Rh	-52080#	300#	51 ms	6		13	15Lo04 TD 1997	$\beta^- = 100; \beta^- n = 10\#; \beta^- 2n = 0.01\#$		
<sup>122</sup> Rh <sup>m</sup>	-51810#	300#	271.0	0.7	830 ns	120	12Ka36 ETD 2012	IT=100		
<sup>122</sup> Pd	-64616	20	195 ms	5	0 <sup>+</sup>	14	15Lo04 T 1994	$\beta^- = 100; \beta^- n < 2.5$		
<sup>122</sup> Ag	-71110	40	529 ms	13	(3 <sup>+</sup> )	07	1978	$\beta^- = 100; \beta^- n = 0.186$ 10		
<sup>122</sup> Ag <sup>m</sup>	-71030#	60#	80#	50#	*	550 ms	50	(1 <sup>-</sup> ) 07 2000	$\beta^- = 100; IT ?; \beta^- n = 0.2\#$	
<sup>122</sup> Ag <sup>n</sup>	-71030#	60#	80#	50#	*	200 ms	50	(9 <sup>-</sup> ) 07 2000	$\beta^- = 100; IT ?; \beta^- n = 0.2\#$	
<sup>122</sup> Ag <sup>p</sup>	-71030#	60#	80#	50#	*	6.3 $\mu$ s	1.0	13La11 T 2013	IT=100	
<sup>122</sup> Cd	-80612.4	2.3	5.24 s	0.03	0 <sup>+</sup>	07	1973	$\beta^- = 100$		
<sup>122</sup> In	-83570	50	1.5 s	0.3	1 <sup>+</sup>	07	1963	$\beta^- = 100$		
<sup>122</sup> In <sup>m</sup>	-83530#	80#	40#	60#	*	10.3 s	0.6	5 <sup>+</sup> 07 1979	$\beta^- = 100$	
<sup>122</sup> In <sup>n</sup>	-83290	130	290	140	BD	10.8 s	0.4	(8 <sup>-</sup> ) 07 1979	$\beta^- = 100$	
<sup>122</sup> Sn	-89941.3	2.4	STABLE		0 <sup>+</sup>	07	1928	IS=4.63 3; 2 $\beta^- ?$		
<sup>122</sup> Sn <sup>m</sup>	-87532.3	2.4	2409.03	0.04	7.5 $\mu$ s	0.9	7 <sup>-</sup>	07 1979	IT=100	
<sup>122</sup> Sn <sup>n</sup>	-87175.7	2.6	2765.6	1.0	62 $\mu$ s	3	(10 <sup>+</sup> )	07 1992	IT=100	
<sup>122</sup> Sn <sup>p</sup>	-85221.1	2.5	4720.2	0.5	146 ns	15	(15 <sup>-</sup> )	12As05 EJT 2012	IT=100	
<sup>122</sup> Sb	-88335.4	2.6	2.7238 d	0.0002	2 <sup>-</sup>	07	1939	$\beta^- = 97.59$ 12; $\beta^+ = 2.41$ 12		
<sup>122</sup> Sb <sup>m</sup>	-88274.0	2.6	61.4131	0.0005	1.86 $\mu$ s	0.08	3 <sup>+</sup>	07 1962	IT=100	
<sup>122</sup> Sb <sup>n</sup>	-88197.9	2.6	137.4726	0.0008	530 $\mu$ s	30	(5 <sup>+</sup> )	07 1963	IT=100	
<sup>122</sup> Sb <sup>p</sup>	-88171.8	2.6	163.5591	0.0017	4.191 m	0.003	(8 <sup>-</sup> )	07 1947	IT=100	
<sup>122</sup> Te	-90314.5	1.5	STABLE		0 <sup>+</sup>	07	1932	IS=2.55 12		
<sup>122</sup> I	-86080	5	3.63 m	0.06	1 <sup>+</sup>	07	12At01 D 1950	e <sup>+</sup> =78 2; e=22 2		
<sup>122</sup> I <sup>m</sup>	-85765	5	314.9	0.4	190 ns	10	(7 <sup>-</sup> )	07 12Mo.A T 2004	IT=100	
<sup>122</sup> I <sup>n</sup>	-85701	5	379.4	0.5	81 $\mu$ s	3	(7 <sup>-</sup> )	07 12Mo.A T 2004	IT=100	
<sup>122</sup> I <sup>p</sup>	-85686	5	394.1	0.5	81 $\mu$ s	3	(8 <sup>+</sup> )	07 12Mo.A T 2004	IT=100	
<sup>122</sup> I <sup>q</sup>	-85636	5	444.1	0.5	148 ns	5	(8 <sup>-</sup> )	07 12Mo.A T 2004	IT=100	
<sup>122</sup> Xe	-85355	11	20.1 h	0.1	0 <sup>+</sup>	07	1952	e=100		
<sup>122</sup> Cs	-78140	30	21.18 s	0.19	1 <sup>+</sup>	07	75Ho09 D 1969	$\beta^+ = 100; \beta^+ \alpha < 2e-7$		
<sup>122</sup> Cs <sup>m</sup>	-78090	30	45.87	0.12	> 1 $\mu$ s		(3 <sup>+</sup> )	07 1987	IT=100	
<sup>122</sup> Cs <sup>n</sup>	-78005	9	140	30	MD	3.70 m	0.11	8 <sup>(-)</sup>	07 1969	$\beta^+ = 100$
<sup>122</sup> Cs <sup>p</sup>	-78010	30	127.07	0.16	360 ms	20	(5 <sup>-</sup> )	07 1969	IT=100	
<sup>122</sup> Cs <sup>x</sup>	-78130	30	14	7	R = 0.1.05	spmix				

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...												
<sup>122</sup> Ba	-74609	28		1.95 m	0.15	0 <sup>+</sup>	07	1974	$\beta^+=100$			
<sup>122</sup> La	-64540#	300#		8.6 s	0.5		07	1984	$\beta^+=100; \beta^+p=?$			
<sup>122</sup> Ce	-57870#	400#		2# s		0 <sup>+</sup>	07	2005	$\beta^+ ?; \beta^+p ?$			
<sup>122</sup> Pr	-44780#	500#		500# ms					$\beta^+ ?; \beta^+p ?$			
* <sup>122</sup> Rh <sup>m</sup>	E : 12Ka36=63.9(0.5) and 207.1(0.5) $\gamma$ rays in cascade to ground-state											
* <sup>122</sup> Rh <sup>m</sup>	T : symmetrized from 12Ka36=820(+130-110)											
* <sup>122</sup> Ag <sup>m</sup>	D : $\beta^-n$ has been observed by 00Kr18 but not quantified											
* <sup>122</sup> Ag <sup>n</sup>	D : $\beta^-n$ has been observed by 00Kr18 but not quantified											
* <sup>122</sup> Cs	D : $\beta^+\alpha$ intensity upper limit is from 75Ho09											
<sup>123</sup> Ru	-37080#	500#		19 ms	2	3/2 <sup>+</sup> #	15	2010	$\beta^-=100; \beta^-n=20\#; \beta^-2n=0.2\#$			
<sup>123</sup> Rh	-49360#	400#		42 ms	4	7/2 <sup>+</sup> #	15	2010	$\beta^-=100; \beta^-n=20\#; \beta^-2n=0\#$			
<sup>123</sup> Pd	-60430	790		108 ms	2	3/2 <sup>+</sup> #	15	1994	$\beta^-=100; \beta^-n=0.4\#$			
<sup>123</sup> Ag	-69550	30		300 ms	5	7/2 <sup>+</sup> #	04	06Mo07	D	1976	$\beta^-=100; \beta^-n=1.0$ 5	
<sup>123</sup> Ag <sup>m</sup>	-69530#	40#	20#	100# ms		1/2 <sup>-</sup> #					$\beta^- ?; IT ?$	
<sup>123</sup> Ag <sup>n</sup>	-68150#	60#	1400#	202 ns	20			13La11	ETD	2013	IT=100	
<sup>123</sup> Ag <sup>p</sup>	-68080	30	1473	393 ns	15	(17/2 <sup>-</sup> )		13La11	ET	2009	IT=100	
<sup>123</sup> Cd	-77414.2	2.7		2.10 s	0.02	3/2 <sup>+</sup>	04	13Yo02	J	1983	$\beta^-=100$	
<sup>123</sup> Cd <sup>m</sup>	-77271	3	143	1.82 s	0.03	11/2 <sup>-</sup>	04	13Yo02	J	1986	$\beta^- ?; IT ?$	
<sup>123</sup> In	-83430	20		6.17 s	0.05	(9/2 <sup>+</sup> )	04			1960	$\beta^-=100$	
<sup>123</sup> In <sup>m</sup>	-83103	20	327.21	47.4 s	0.4	(1/2 <sup>-</sup> )	04			1960	$\beta^-=100$	
<sup>123</sup> In <sup>n</sup>	-81352	20	2078.1	1.4 $\mu$ s	0.2	(17/2 <sup>-</sup> )		04Sc42	ETJ	2004	IT=100	
<sup>123</sup> In <sup>p</sup>	-81300	50	2128.1	> 100 $\mu$ s		(21/2 <sup>-</sup> )	10	10Re01	EJT	2010	IT=100	
<sup>123</sup> Sn	-87816.2	2.4		129.2 d	0.4	11/2 <sup>-</sup>	04			1948	$\beta^-=100$	
<sup>123</sup> Sn <sup>m</sup>	-87791.6	2.4	24.6	40.06 m	0.01	3/2 <sup>+</sup>	04			1948	$\beta^-=100$	
<sup>123</sup> Sn <sup>n</sup>	-85871.2	2.6	1945.0	7.4 $\mu$ s	2.6	(19/2 <sup>+</sup> )	04			1992	IT=100	
<sup>123</sup> Sn <sup>p</sup>	-85663.2	2.7	2153.0	6 $\mu$ s		(23/2 <sup>+</sup> )	04			1994	IT=100	
<sup>123</sup> Sn <sup>q</sup>	-85103.2	2.8	2713.0	34 $\mu$ s		(27/2 <sup>-</sup> )	04			1994	IT=100	
<sup>123</sup> Sb	-89224.1	1.5		STABLE		7/2 <sup>+</sup>	04			1922	IS=42.79 5	
<sup>123</sup> Sb <sup>m</sup>	-86986.3	1.5	2237.8	214 ns	3	19/2 <sup>-</sup>		09Wa02	ETJ	2005	IT=100	
<sup>123</sup> Sb <sup>n</sup>	-86610.7	1.6	2613.4	65 $\mu$ s	1	23/2 <sup>+</sup>		09Wa02	ETJ	2007	IT=100	
<sup>123</sup> Te	-89172.2	1.5		STABLE	(>2 Py)	1/2 <sup>+</sup>	04	03Al02	T	1932	IS=0.89 3; $\epsilon=100$	
<sup>123</sup> Te <sup>m</sup>	-88924.7	1.5	247.47	119.2 d	0.1	11/2 <sup>-</sup>	04			1951	IT=100	
<sup>123</sup> I	-87944	4		13.2235 h	0.0019	5/2 <sup>+</sup>	04			1949	$\beta^+=100$	
<sup>123</sup> Xe	-85249	10		2.08 h	0.02	1/2 <sup>+</sup>	04	90Ne.A	J	1952	$\beta^+=100$	
<sup>123</sup> Xe <sup>m</sup>	-85064	10	185.18	5.49 $\mu$ s	0.26	7/2 <sup>(-)</sup>	04			1981	IT=100	
<sup>123</sup> Cs	-81044	12		5.88 ms	0.03	1/2 <sup>+</sup>	04			1954	$\beta^+=100$	
<sup>123</sup> Cs <sup>m</sup>	-80888	12	156.27	1.64 s	0.12	(11/2 <sup>-</sup> )	04			1972	IT=100	
<sup>123</sup> Cs <sup>n</sup>	-80792	23	252	114 ns	5	(9/2 <sup>+</sup> )	04	GAu127	E	2000	IT=100	
<sup>123</sup> Cs <sup>s</sup>	-81037	13	7	R < 0.1		spmix					*	
<sup>123</sup> Ba	-75655	12		2.7 m	0.4	5/2 <sup>(+)</sup>	04			1962	$\beta^+=100$	
<sup>123</sup> Ba <sup>m</sup>	-75534	12	120.95	830 ns	60	1/2 <sup>+</sup> #	04			1991	IT=100	
<sup>123</sup> La	-68650#	200#		17 s	3	11/2 <sup>-</sup> #	04			1978	$\beta^+=100$	
<sup>123</sup> Ce	-60290#	300#		3.8 s	0.2	(5/2 <sup>(+)</sup> )	04			1984	$\beta^+=100; \beta^+p=?$	
<sup>123</sup> Pr	-50230#	400#		800# ms		3/2 <sup>+</sup> #					$\beta^+ ?; \beta^+p ?$	
* <sup>123</sup> Ag <sup>n</sup>	E : 13La11=1365keV above <sup>123</sup> Ag <sup>m</sup>											
* <sup>123</sup> Ag <sup>p</sup>	T : average 13La11=393(16) 09St28=396(37) J : 09St28=(17/2 <sup>-</sup> )											
* <sup>123</sup> In <sup>n</sup>	E : derived by NUBASE from least-squares fit to $\gamma$ -ray energies											
* <sup>123</sup> In <sup>p</sup>	E : no direct depopulating $\gamma$ seen, assumed less than 50 keV											
* <sup>123</sup> Sb <sup>m</sup>	E : derived from least-squares fit to $\gamma$ -ray energies											
* <sup>123</sup> Sb <sup>m</sup>	ETJ : also 07Ju06 2239.1(1.0) keV, 190(30) ns, 19/2 <sup>-</sup> ; and											
* <sup>123</sup> Sb <sup>m</sup>	ETJ : 05Po03 2247.1(0.4) keV, 110(10) ns (conflicting), (19/2 <sup>-</sup> )											
* <sup>123</sup> Sb <sup>n</sup>	E : derived from least-squares fit to $\gamma$ -ray energies											
* <sup>123</sup> Sb <sup>n</sup>	ETJ : also 07Ju06 2614.1(1.0) keV, 66(4) $\mu$ s, 23/2 <sup>+</sup> ; and											
* <sup>123</sup> Sb <sup>n</sup>	ETJ : 08Jo03 2614.2(0.6) keV, 52(3) $\mu$ s (conflicting), 23/2 <sup>+</sup>											
* <sup>123</sup> Cs <sup>n</sup>	E : 231.63 + x; x estimated 20#20											
<sup>124</sup> Ru	-33960#	600#		15 ms	3	0 <sup>+</sup>	15	2010	$\beta^-=100; \beta^-n=10\#; \beta^-2n=0\#$			
<sup>124</sup> Rh	-44890#	400#		30 ms	2		15	2010	$\beta^-=100; \beta^-n=20\#; \beta^-2n=0.3\#$			
<sup>124</sup> Pd	-58390#	300#		88 ms	15	0 <sup>+</sup>	14	15Lo04	T	1997	$\beta^-=100; \beta^-n=0.03\#$	
<sup>124</sup> Pd <sup>m</sup>	-58330#	300#	62.2	> 20 $\mu$ s				14	12Ka36	ET	2012	IT=100; $\beta^- ?$
<sup>124</sup> Ag	-66200	250		177.9 ms	2.6	(2 <sup>-</sup> )	15	14Ba18	J	1984	$\beta^-=100; \beta^-n=1.3$ 9	
<sup>124</sup> Ag <sup>m</sup>	-66200#	270#	0#	144 ms	20	(8 <sup>-</sup> )	15	14Ba18	TJ	1995	$\beta^-=100; \beta^-n=1\#$	
<sup>124</sup> Ag <sup>n</sup>	-65970	250	231.1	1.7 $\mu$ s	0.3		15	12Ka36	E	2012	IT=100	
<sup>124</sup> Cd	-76701.7	3.0		1.25 s	0.02	0 <sup>+</sup>	08			1974	$\beta^-=100$	
<sup>124</sup> In	-80870	30		3.12 s	0.09	3 <sup>+</sup>	08	13Ma15	J	1964	$\beta^-=100$	
<sup>124</sup> In <sup>m</sup>	-80890	50	-20	3.7 s	0.2	(8 <sup>(-)</sup> )	08			1974	$\beta^- \approx 100; IT ?$	
... A-group is continued on next page ...												

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>124</sup> Sn	-88234.2	1.0	STABLE (>100 Py)	0+	08	52Ka41	T 1922	IS=5.79 5; 2β <sup>-</sup> ?	
<sup>124</sup> Sn <sup>m</sup>	-86029.6	1.0	270 ns	60	5-	08 FGK127	J 1979	IT=100	
<sup>124</sup> Sn <sup>n</sup>	-85909.2	1.0	3.1 μs	0.5	7-	08 FGK127	J 1979	IT=100	
<sup>124</sup> Sn <sup>p</sup>	-85577.6	1.1	45 μs	5	10+	08 FGK127	J 1992	IT=100	
<sup>124</sup> Sn <sup>q</sup>	-83682.8	1.2	260 ns	25	15-	12As05	EJT 2012	IT=100	
<sup>124</sup> Sb	-87620.2	1.5	60.20 d	0.03	3-	08		β <sup>-</sup> =100	
<sup>124</sup> Sb <sup>m</sup>	-87609.3	1.5	93 s	5	5+	08		IT=75 5; β <sup>-</sup> =25 5	
<sup>124</sup> Sb <sup>n</sup>	-87583.4	1.5	20.2 m	0.2	(8)-	08		IT=100	
<sup>124</sup> Sb <sup>p</sup>	-87579.4	1.5	3.2 μs	0.3	(3+,4+)	08		IT=100	
<sup>124</sup> Te	-90525.3	1.5	STABLE		0+	08		IS=4.74 14	
<sup>124</sup> I	-87365.7	2.4	4.1760 d	0.0003	2-	08		β <sup>+</sup> =100	
<sup>124</sup> Xe	-87661.4	1.8	STABLE (>200 Ty)		0+	08		IS=0.0952 3; 2β <sup>+</sup> ?	
<sup>124</sup> Cs	-81731	8	30.9 s	0.4	1+	08		β <sup>+</sup> =100	
<sup>124</sup> Cs <sup>m</sup>	-81268	8	462.63	0.14	6.3 s	0.2	(7)+	08 1983	IT=100
<sup>124</sup> Cs <sup>x</sup>	-81701	22	30	20	R=?			spmix	
<sup>124</sup> Ba	-79090	12	11.0 m	0.5	0+	08		β <sup>+</sup> =100	
<sup>124</sup> La	-70260	60	* & 29.21 s	0.17	(7-,8-)	08	92Id01 J 1978	β <sup>+</sup> =100	
<sup>124</sup> La <sup>m</sup>	-70160#	120#	100#	100#	* & 21 s	4	low(+#)	08 92Id01 J 1992	β <sup>+</sup> =100
<sup>124</sup> Ce	-64920#	300#			9.1 s	1.2	0+	08 97As05 T 1978	β <sup>+</sup> =100
<sup>124</sup> Pr	-53150#	400#			1.2 s	0.2	08		β <sup>+</sup> =100; β <sup>+</sup> p=?
<sup>124</sup> Nd	-44530#	500#			500# ms				β <sup>+</sup> ?; β <sup>+</sup> p ?
* <sup>124</sup> Ag	T : average 15Lo04=180(3) 95Fe12=172(5); other 14Ba18=191(28)								**
* <sup>124</sup> Ag <sup>n</sup>	E : 12Ka36=75.5(0.5) and 155.6(0.5) γ rays in cascade to ground-state								**
* <sup>124</sup> Sn <sup>m</sup>	J : E1 to 4+; L(p,p)=5 for <sup>124</sup> Sn <sup>m</sup> ; E2 to 5- for <sup>124</sup> Sn <sup>n</sup> ; E2 to 8+ for <sup>124</sup> Sn <sup>p</sup>								**
* <sup>124</sup> Xe	T : 2v-εε: 16Ab03>4.7 Zy (at 90% C.L.)								**
* <sup>124</sup> Ce	T : average 97As05=10.8(1.5) 78Bo32=6(2)								**
<sup>125</sup> Rh	-42000#	500#			26.5 ms	2.0	7/2+#	15 2010	β <sup>-</sup> =100; β <sup>-</sup> n=20#; β <sup>-</sup> 2n=0.03#
<sup>125</sup> Pd	-54120#	400#			57 ms	10	3/2+#	15 2008	β <sup>-</sup> =100 ?; β <sup>-</sup> n=6#
<sup>125</sup> Ag	-64520	430			159 ms	8	7/2+#	15 1994	β <sup>-</sup> =100; β <sup>-</sup> n=5#
<sup>125</sup> Ag <sup>m</sup>	-64500#	430#	20#	20#	50# ms		1/2-#		β <sup>-</sup> ?; IT ?
<sup>125</sup> Ag <sup>n</sup>	-63640#	430#	880#	20#	80 ns	17		15 13La11 ET 2013	IT=100
<sup>125</sup> Ag <sup>p</sup>	-63020	430	1501.2	0.6	491 ns	20	(17/2-)	15 2009	IT=100
<sup>125</sup> Cd	-73348.1	2.9			680 ms	40	3/2+	11 13Yo02 J 1986	β <sup>-</sup> =100
<sup>125</sup> Cd <sup>m</sup>	-73162	3	186	4	MD 480 ms	30	11/2-	11 13Yo02 J 1986	β <sup>-</sup> =100
<sup>125</sup> Cd <sup>n</sup>	-71840	70	1512	70	19 μs	3	(19/2+)	11 11Si32 EJT 2011	IT=100
<sup>125</sup> In	-80477	27			2.36 s	0.04	9/2+	11 1967	β <sup>-</sup> =100
<sup>125</sup> In <sup>m</sup>	-80117	27	360.12	0.09	12.2 s	0.2	1/2(-)	11 1974	β <sup>-</sup> =100
<sup>125</sup> In <sup>n</sup>	-78468	27	2009.4	0.7	9.4 μs	0.6	(19/2+)	11 1998	IT=100
<sup>125</sup> In <sup>p</sup>	-78316	27	2161.2	0.9	5.0 ms	1.5	(23/2-)	11 1998	IT=100
<sup>125</sup> Sn	-85896.4	1.0			9.64 d	0.03	11/2-	11 1939	β <sup>-</sup> =100
<sup>125</sup> Sn <sup>m</sup>	-85868.9	1.0	27.50	0.14	9.52 m	0.05	3/2+	11 1939	β <sup>-</sup> =100
<sup>125</sup> Sn <sup>n</sup>	-84003.6	1.0	1892.8	0.3	6.2 μs	0.2	19/2+	11 08Lo07 J 2000	IT=100
<sup>125</sup> Sn <sup>p</sup>	-83836.9	1.1	2059.5	0.4	650 ns	60	23/2+	11 16Is03 T 2008	IT=100
<sup>125</sup> Sn <sup>q</sup>	-83272.9	1.1	2623.5	0.5	230 ns	17	(27/2-)	11 08Lo07 T 2000	IT=100
<sup>125</sup> Sb	-88256.3	2.6			2.7586 y	0.0003	7/2+	11 1951	β <sup>-</sup> =100
<sup>125</sup> Sb <sup>m</sup>	-86285.1	2.6	1971.25	0.20	4.1 μs	0.2	15/2-	11 2007	IT=100
<sup>125</sup> Sb <sup>n</sup>	-86144.2	2.6	2112.1	0.3	28.0 μs	0.7	19/2-	11 FGK128 J 2007	IT=100
<sup>125</sup> Sb <sup>q</sup>	-85785.3	2.6	2471.0	0.4	272 ns	16	(23/2)+	11 2007	IT=100
<sup>125</sup> Te	-89023.0	1.5			STABLE		1/2+	11 1931	IS=7.07 15
<sup>125</sup> Te <sup>m</sup>	-88878.2	1.5	144.775	0.008	57.40 d	0.15	11/2-	11 1949	IT=100
<sup>125</sup> I	-88837.2	1.5			59.407 d	0.010	5/2+	11 1947	ε=100
<sup>125</sup> Xe	-87193.4	1.8			16.9 h	0.2	1/2(+)	11 1950	β <sup>+</sup> =100
<sup>125</sup> Xe <sup>m</sup>	-86940.8	1.8	252.61	0.14	56.9 s	0.9	9/2(-)	11 1954	IT=100
<sup>125</sup> Xe <sup>n</sup>	-86897.5	1.8	295.89	0.15	140 ns	30	7/2(+)	11 1979	IT=100
<sup>125</sup> Cs	-84088	8			46.7 m	0.1	1/2(+)	11 1954	β <sup>+</sup> =100
<sup>125</sup> Cs <sup>m</sup>	-83822	8	266.1	1.1	900 μs	30	(11/2-)	11 98Su16 J 1998	IT=100
<sup>125</sup> Ba	-79669	11			3.3 m	0.3	1/2(+#)	11 1968	β <sup>+</sup> =100
<sup>125</sup> Ba <sup>m</sup>	-79559	23	110	20	2.76 μs	0.14	(7/2-)	11 FGK128 J 1989	IT=100
<sup>125</sup> La	-73759	26			64.8 s	1.2	11/2-#	11 1973	β <sup>+</sup> =100
<sup>125</sup> La <sup>m</sup>	-73652	26	107.00	0.10	390 ms	40	(3/2+)	11 99Ca21 J 1998	IT=100
<sup>125</sup> Ce	-66660#	200#			9.7 s	0.3	(7/2-)	11 02Pe15 J 1978	β <sup>+</sup> =100; β <sup>+</sup> p=?
<sup>125</sup> Ce <sup>m</sup>	-66570#	200#	93.6	0.4	13 s	10	(1/2+)	11 07Su07 ETJ 2007	IT=100
... A-group is continued on next page ...									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>125</sup> Pr	-57940# 300#		3.3 s 0.7	3/2 <sup>+</sup> #	11		2002	$\beta^+$ =100; $\beta^+$ p ?
<sup>125</sup> Nd	-47600# 400#		650 ms 150	(5/2) <sup>(+)</sup> #	11		1999	$\beta^+$ =100; $\beta^+$ p>0
* <sup>125</sup> Ag <sup>m</sup>		E: 13La11: 859.9keV above <sup>125</sup> Ag <sup>m</sup>						**
* <sup>125</sup> Cd <sup>m</sup>		E: 11Si32=1461.8(0.5) keV above the 11/2 <sup>-</sup> isomer						**
* <sup>125</sup> Sn <sup>p</sup>		J: E2 to 19/2 <sup>+</sup> for <sup>125</sup> Sn <sup>p</sup> ; E2 to 23/2 <sup>-</sup> for <sup>125</sup> Sn <sup>q</sup>						**
* <sup>125</sup> Sb		T: rounded from ENSDF2011=2.75856(0.00025); other 16Li01=2.75817(0.00082)						**
* <sup>125</sup> Sb <sup>n</sup>		J: E2 to 15/2 <sup>-</sup> T: others recent 10Re01=25(4) 07Ju06=25(4)						**
* <sup>125</sup> Ba <sup>m</sup>		E: 67.7(0.4) above 5/2 <sup>+</sup> # level at estimated 30#20 J: E1 to 5/2 <sup>+</sup>						**
* <sup>125</sup> La <sup>m</sup>		J: 3/2 <sup>+</sup> # from trends in La isotopes; low spin and even-parity from 99Ca21						**
* <sup>125</sup> Ce <sup>m</sup>		T: symmetrized from 134(+641-61) s for fully ionized ion; icc=38.1 for a						**
* <sup>125</sup> Ce <sup>m</sup>		T: 93.6(0.4) keV, E3 transition; ENSDF quotes 3.4(2.7) s						**
<sup>126</sup> Rh	-37300# 500#		19 ms 3		15		2010	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=0.2#
<sup>126</sup> Pd	-51860# 400#		48.6 ms 1.2	0 <sup>+</sup>	15		2008	$\beta^-$ =100; $\beta^-$ n=0.4#
<sup>126</sup> Pd <sup>m</sup>	-49840# 400#	2023.5 0.7	330 ns 40	(5 <sup>-</sup> )	15		2013	IT=100
<sup>126</sup> Pd <sup>n</sup>	-49750# 400#	2109.7 0.9	440 ns 30	(7 <sup>-</sup> )	15		2013	IT=100
<sup>126</sup> Pd <sup>p</sup>	-49450# 400#	2406.0 1.0	23.0 ms 0.9	(10 <sup>+</sup> )	15		2014	IT=?; B ?
<sup>126</sup> Ag	-60680# 200#		99.3 ms 4.6	2 <sup>+</sup> #	15	15Lo04 T	1994	$\beta^-$ =100; $\beta^-$ n=6# *
<sup>126</sup> Ag <sup>m</sup>	-60580# 220#	100# 100#	92 ms 9	8 <sup>-</sup> #	15		1995	$\beta^-$ =100; IT ?; $\beta^-$ n=6#
<sup>126</sup> Ag <sup>n</sup>	-60430# 200#	254.8 0.5	27 $\mu$ s 6	1 <sup>-</sup> #	15		2012	IT=100
<sup>126</sup> Cd	-72256.8 2.5		513 ms 6	0 <sup>+</sup>	03	15Lo04 T	1978	$\beta^-$ =100
<sup>126</sup> In	-77773 27		1.53 s 0.01	3 <sup>(+)</sup> #	03		1974	$\beta^-$ =100
<sup>126</sup> In <sup>m</sup>	-77710 50	70 60	1.64 s 0.05	8 <sup>(-)</sup> #	03	79Fo10 J	1970	$\beta^-$ =100
<sup>126</sup> In <sup>n</sup>	-77530 27	243.3 0.2	22 $\mu$ s 2	1 <sup>(-)</sup>		04Sc42 ETJ	2003	IT=100
<sup>126</sup> Sn	-86015 10		230 ky 14	0 <sup>+</sup>	03		1962	$\beta^-$ =100
<sup>126</sup> Sn <sup>m</sup>	-83796 10	2218.99 0.08	5.8 $\mu$ s 0.7	7 <sup>-</sup>	03	12As05 T	1979	IT=100 *
<sup>126</sup> Sn <sup>n</sup>	-83451 10	2564.5 0.5	7.6 $\mu$ s 0.3	10 <sup>+</sup>	03	12As05 TJ	2000	IT=100 *
<sup>126</sup> Sn <sup>p</sup>	-81669 10	4345.7 0.8	160 ns 20	15 <sup>-</sup>		12As05 EJT	2012	IT=100
<sup>126</sup> Sb	-86390 30		12.35 d 0.06	(8 <sup>-</sup> )	03		1956	$\beta^-$ =100
<sup>126</sup> Sb <sup>m</sup>	-86370 30	17.7 0.3	19.15 m 0.08	(5 <sup>+</sup> )	03		1956	$\beta^-$ =86.4; IT=14.4
<sup>126</sup> Sb <sup>n</sup>	-86350 30	40.4 0.3	11 s	(3 <sup>-</sup> )	03		1976	IT=100
<sup>126</sup> Sb <sup>p</sup>	-86290 30	104.6 0.3	553 ns 5	(3 <sup>+</sup> )	03		1976	IT=100
<sup>126</sup> Te	-90065.3 1.5		STABLE	0 <sup>+</sup>	03		1924	IS=18.84 25
<sup>126</sup> I	-87911 4		12.93 d 0.05	2 <sup>-</sup>	03		1938	$\beta^+$ =52.7 5; $\beta^-$ =47.3 5
<sup>126</sup> I <sup>m</sup>	-87800 4	111.00 0.23	128 ns	3 <sup>+</sup>		12Mo.A EJT	2012	IT=100
<sup>126</sup> Xe	-89147 3		STABLE	0 <sup>+</sup>	03		1922	IS=0.0890 2; 2 $\beta^+$ ?
<sup>126</sup> Cs	-84351 10		1.64 m 0.02	1 <sup>+</sup>	03		1954	$\beta^+$ =100
<sup>126</sup> Cs <sup>m</sup>	-84078 10	273.0 0.7	> 1 $\mu$ s		03		1993	IT=100
<sup>126</sup> Cs <sup>n</sup>	-83755 10	596.1 1.1	171 $\mu$ s 14		03		1993	IT=100
<sup>126</sup> Ba	-82670 12		100 m 2	0 <sup>+</sup>	03		1954	$\beta^+$ =100
<sup>126</sup> La	-74970 90		54 s 2	(5) <sup>(+)</sup> #	03		1961	$\beta^+$ =100
<sup>126</sup> La <sup>m</sup>	-74760 400	210 410	20 s 20	(0 <sup>-</sup> , 1 <sup>-</sup> , 2 <sup>-</sup> )	03		1997	$\beta^+$ =100 *
<sup>126</sup> Ce	-70821 28		51.0 s 0.3	0 <sup>+</sup>	03		1978	$\beta^+$ =100
<sup>126</sup> Pr	-60320# 200#		3.12 s 0.18	(4, 5, 6)	03	88Ba42 T	1983	$\beta^+$ =100; $\beta^+$ p=? *
<sup>126</sup> Nd	-52990# 300#		1# s (>200 ns)	0 <sup>+</sup>	03	00So11 I	2000	$\beta^+$ ?; $\beta^+$ p ?
<sup>126</sup> Pm	-39350# 500#		500# ms					$\beta^+$ ?; $\beta^+$ p ?
* <sup>126</sup> Ag		T: average 15Lo04=98(5) 95Fe12=107(12); other 14Ba18=52(10) at variance						**
* <sup>126</sup> Sn <sup>m</sup>		T: average 12As05=6.6(1.4) 10Ti01=5.6(0.8)						**
* <sup>126</sup> Sn <sup>n</sup>		T: average 12As05=7.7(0.5) 10Ti01=7.5(0.3)						**
* <sup>126</sup> La <sup>m</sup>		T: 97As05: "by far shorter than 50 s"						**
* <sup>126</sup> Pr		T: average 95Os03=3.14(0.22) 88Ba42=3.0(0.4) 83Ni05=3.2(0.6)						**
<sup>127</sup> Rh	-34030# 600#		28 ms 14	7/2 <sup>+</sup> #	15	15Lo04 TD	2015	$\beta^-$ =100; $\beta^-$ n=30#; $\beta^-$ 2n=0.2# *
<sup>127</sup> Pd	-47180# 500#		38 ms 2	3/2 <sup>+</sup> #	15		2010	$\beta^-$ =100; $\beta^-$ n=10#; $\beta^-$ 2n=0#
<sup>127</sup> Ag	-58440# 200#		89 ms 2	7/2 <sup>+</sup> #	11	15Lo04 T	1995	$\beta^-$ =100; $\beta^-$ n=7# *
<sup>127</sup> Ag <sup>m</sup>	-58420# 200#	20# 20#	20# ms	1/2 <sup>-</sup> #				$\beta^-$ ?; IT ?
<sup>127</sup> Cd	-68747 12		330 ms 20	3/2 <sup>+</sup>	11	13Yo02 J	1986	$\beta^-$ =100; $\beta^-$ n=0.03# *
<sup>127</sup> Cd <sup>m</sup>	-68472 8	276 15	200# ms	11/2 <sup>-</sup>		13Yo02 J		$\beta^-$ ?; IT ?
<sup>127</sup> Cd <sup>n</sup>	-66930 30	1813 32	17.5 $\mu$ s 0.3	(19/2 <sup>+</sup> )		10Na17 ETJ	2010	IT=100 *
<sup>127</sup> In	-76896 21		1.09 s 0.01	(9/2 <sup>+</sup> )	11		1975	$\beta^-$ =100; $\beta^-$ n<0.03
<sup>127</sup> In <sup>m</sup>	-76487 21	408.9 0.3	3.67 s 0.04	1/2 <sup>-</sup> #	11		1974	$\beta^-$ =100; $\beta^-$ n=0.69 4
<sup>127</sup> In <sup>n</sup>	-75030 60	1870 60	1.04 s 0.10	(21/2 <sup>-</sup> )	11		2004	$\beta^-$ =100; $\beta^-$ n=1#
<sup>127</sup> In <sup>p</sup>	-74531 21	2364.7 0.9	9 $\mu$ s 2	(29/2 <sup>+</sup> )	11	04Sc42 ETJ	2004	IT=100 *
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...												
<sup>127</sup> Sn	-83471	10		2.10	h	0.04	11/2 <sup>-</sup>	11	1951	$\beta^-$ =100		
<sup>127</sup> Sn <sup>m</sup>	-83466	10	5.07	0.06	4.13	m	0.03	3/2 <sup>+</sup>	11	1962	$\beta^-$ =100	
<sup>127</sup> Sn <sup>n</sup>	-81644	10	1826.67	0.16	4.52	$\mu$ s	0.15	19/2 <sup>+</sup>	11	08Lo07 J	2000	IT=100
<sup>127</sup> Sn <sup>p</sup>	-81540	10	1930.97	0.17	1.26	$\mu$ s	0.15	(23/2 <sup>+</sup> )	11	2004	IT=100	
<sup>127</sup> Sn <sup>q</sup>	-80919	10	2552.4	1.0	250	ns	30	(27/2 <sup>-</sup> )	11	08Lo07 J	2008	IT=100
<sup>127</sup> Sb	-86699	5			3.85	d	0.05	7/2 <sup>+</sup>	11	1939	$\beta^-$ =100	
<sup>127</sup> Sb <sup>m</sup>	-84779	5	1920.19	0.21	11	$\mu$ s	1	15/2 <sup>-</sup>	11	09Wa24 J	1974	IT=100
<sup>127</sup> Sb <sup>n</sup>	-84374	5	2324.7	0.4	234	ns	12	23/2 <sup>+</sup>	11	09Wa24 TJ	2005	IT=100
<sup>127</sup> Te	-88281.7	1.5			9.35	h	0.07	3/2 <sup>+</sup>	11	1938	$\beta^-$ =100	
<sup>127</sup> Te <sup>m</sup>	-88193.5	1.5	88.23	0.07	106.1	d	0.7	11/2 <sup>-</sup>	11	1940	IT=97.6 2; $\beta^-$ =2.4 2	
<sup>127</sup> I	-88984	4			STABLE			5/2 <sup>+</sup>	11	1920	IS=100.	
<sup>127</sup> Xe	-88322	4			36.346	d	0.003	1/2 <sup>+</sup>	11	1950	$\epsilon$ =100	
<sup>127</sup> Xe <sup>m</sup>	-88025	4	297.10	0.08	69.2	s	0.9	9/2 <sup>-</sup>	11	1940	IT=100	
<sup>127</sup> Cs	-86240	6			6.25	h	0.10	1/2 <sup>+</sup>	11	1950	$\beta^+$ =100	
<sup>127</sup> Cs <sup>m</sup>	-85788	6	452.23	0.21	55	$\mu$ s	3	(11/2 <sup>-</sup> )	11	1980	IT=100	
<sup>127</sup> Ba	-82818	11			12.7	m	0.4	1/2 <sup>+</sup>	11	1952	$\beta^+$ =100	
<sup>127</sup> Ba <sup>m</sup>	-82738	11	80.32	0.11	1.93	s	0.07	7/2 <sup>-</sup>	11	1992	IT=100	
<sup>127</sup> La	-77896	26			5.1	m	0.1	(11/2 <sup>-</sup> )	11	1963	$\beta^+$ =100	
<sup>127</sup> La <sup>m</sup>	-77882	26	14.2	0.4	3.7	m	0.4	(3/2 <sup>+</sup> )	11	1963	$\beta^+$ $\approx$ 100	
<sup>127</sup> Ce	-71979	29			34	s	2	(1/2 <sup>+</sup> )	11	1978	$\beta^+$ =100	
<sup>127</sup> Ce <sup>m</sup>	-71972	29	7.3	1.1	28.6	s	0.7	5/2 <sup>+</sup> #	11	1978	$\beta^+$ =100	
<sup>127</sup> Ce <sup>n</sup>	-71942	29	36.8	1.2	> 10	$\mu$ s		(7/2 <sup>-</sup> )	11	1995	IT=100	
<sup>127</sup> Pr	-64540#	200#			4.2	s	0.3	3/2 <sup>+</sup> #	11	1995	$\beta^+$ =100	
<sup>127</sup> Pr <sup>m</sup>	-63940#	280#	600#	200#	50#	ms		11/2 <sup>-</sup>	11	98Mo30 J	1998	$\beta^+$ ?; IT ?
<sup>127</sup> Nd	-55540#	300#			1.8	s	0.4	5/2 <sup>+</sup> #	11	1983	$\beta^+$ =100; $\beta^+$ p=?	
<sup>127</sup> Pm	-44790#	400#			1#	s		5/2 <sup>+</sup> #			$\beta^+$ ?; p ?	
* <sup>127</sup> Rh	T : symmetrized from 15Lo04=20(+20-7)											
* <sup>127</sup> Ag	T : other 96Wo.A=79(3) supersedes 95Fe12=109(25) from same group											
* <sup>127</sup> Cd	T : from 15Lo04=330(20)											
* <sup>127</sup> Cd <sup>n</sup>	E : 1560.1(0.5) keV above <sup>127</sup> Cd <sup>m</sup> T : other 12Ka36=11.0(+9.2-3.5)											
* <sup>127</sup> In <sup>p</sup>	E : derived by NUBASE from least-squares fit to $\gamma$ -ray energies											
* <sup>127</sup> Sb <sup>n</sup>	T : also 05Po03=165(20) conflicting, not used											
... A-group is continued on next page ...												
<sup>128</sup> Pd	-44490#	500#			35	ms	3	0 <sup>+</sup>	16	2010	$\beta^-$ =100; $\beta^-$ n=20#	
<sup>128</sup> Pd <sup>m</sup>	-42340#	500#	2151.0	1.0	5.8	$\mu$ s	0.8	(8 <sup>+</sup> )	16	2013	IT=100	
<sup>128</sup> Ag	-54620#	300#			59	ms	5	0 <sup>+</sup>	15	15Lo04 T	2000	$\beta^-$ =100; $\beta^-$ n=8#; $\beta^-$ 2n=0.01#
<sup>128</sup> Cd	-67242	7			246	ms	2	0 <sup>+</sup>	15	16Du13 T	1986	$\beta^-$ =100; $\beta^-$ n=0.7#
<sup>128</sup> Cd <sup>m</sup>	-65372	7	1870.5	0.3	270	ms	7	(5 <sup>-</sup> )	15	2009	IT=100	
<sup>128</sup> Cd <sup>n</sup>	-64527	7	2714.6	0.4	3.56	$\mu$ s	0.06	(10 <sup>+</sup> )	15	2009	IT=100	
<sup>128</sup> Cd <sup>p</sup>	-62955	7	4286.6	1.5	6.3	ms	0.8	(15 <sup>-</sup> )	15	16Ju.A ETJ	2016	IT=100
<sup>128</sup> In	-74150	150			816	ms	27	(3 <sup>+</sup> )	15	93Ru01 D	1975	$\beta^-$ =100; $\beta^-$ n=0.038 3
<sup>128</sup> In <sup>m</sup>	-74060	30	80	160	720	ms	100	(8 <sup>-</sup> )	15	1986	$\beta^-$ =100	
<sup>128</sup> In <sup>n</sup>	-73900	150	247.87	0.10	23	$\mu$ s	2	(1 <sup>-</sup> )	15	04Sc42 J	1988	IT=100
<sup>128</sup> Sn	-83362	18			59.07	m	0.14	0 <sup>+</sup>	15	1956	$\beta^-$ =100	
<sup>128</sup> Sn <sup>m</sup>	-81271	18	2091.50	0.11	6.5	s	0.5	(7 <sup>-</sup> )	15	1979	IT=100	
<sup>128</sup> Sn <sup>n</sup>	-80870	18	2491.91	0.17	2.91	$\mu$ s	0.14	(10 <sup>+</sup> )	15	1981	IT=100	
<sup>128</sup> Sn <sup>p</sup>	-79263	18	4099.5	0.4	220	ns	30	(15 <sup>-</sup> )	15	2011	IT=100	
<sup>128</sup> Sb	-84630	19			9.05	h	0.04	8 <sup>-</sup>	15	1956	$\beta^-$ =100	
<sup>128</sup> Sb <sup>m</sup>	-84620	18	10	7	10.41	m	0.18	5 <sup>+</sup>	15	1955	$\beta^-$ =96.4 10; IT=3.6 10	
<sup>128</sup> Te	-88993.7	0.9			2.0	Yy	0.3	0 <sup>+</sup>	15	15Ba11 T	1924	IS=31.74 8; 2 $\beta^-$ =100
<sup>128</sup> Te <sup>m</sup>	-86202.9	0.9	2790.8	0.3	363	ns	27	(10 <sup>+</sup> )	15	04Va03 T	1998	IT=100
<sup>128</sup> I	-87739	4			24.99	m	0.02	1 <sup>+</sup>	15	1934	$\beta^-$ =93.1 8; $\beta^+$ =6.9 8	
<sup>128</sup> I <sup>m</sup>	-87601	4	137.851	0.003	845	ns	20	4 <sup>-</sup>	15	1982	IT=100	
<sup>128</sup> I <sup>n</sup>	-87572	4	167.368	0.004	175	ns	15	(6 <sup>-</sup> )	15	1991	IT=100	
<sup>128</sup> Xe	-89860.3	1.1			STABLE			0 <sup>+</sup>	15	1922	IS=1.9102 8	
<sup>128</sup> Xe <sup>m</sup>	-87073.1	1.1	2787.2	0.3	83	ns	2	8 <sup>-</sup>	15	1981	IT=100	
<sup>128</sup> Cs	-85932	5			3.640	m	0.014	1 <sup>+</sup>	15	93Al03 T	1951	$\beta^+$ =100
<sup>128</sup> Ba	-85378	5			2.43	d	0.05	0 <sup>+</sup>	15	1950	$\epsilon$ =100	
<sup>128</sup> La	-78630	50			5.18	m	0.14	(5 <sup>+</sup> )	15	97Ha30 T	1961	$\beta^+$ =100
<sup>128</sup> La <sup>m</sup>	-78530#	110#	100#	100#	< 1.4	m		(1 <sup>+</sup> , 2 <sup>-</sup> )	15	1995	$\beta^+$ =100	
<sup>128</sup> Ce	-75534	28			3.93	m	0.02	0 <sup>+</sup>	15	00Li08 T	1968	$\beta^+$ =100
<sup>128</sup> Pr	-66331	30			2.85	s	0.09	(3 <sup>+</sup> )	15	99Xi03 J	1985	$\beta^+$ =100; $\beta^+$ p=?
<sup>128</sup> Nd	-60310#	200#			5#	s		0 <sup>+</sup>	15	1985	$\beta^+$ ?	

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
$^{128}\text{Pm}$	-47790#	300#	1.0 s	0.3	(5,6,7) <sup>(+)</sup>	15 93Li40 D	1999	$\beta^+ \approx 100$ ; $\beta^+ p ?$ ; $p=0$
$^{128}\text{Sm}$	-38670#	500#	500# ms		$0^+$			$\beta^+ ?$ ; $\beta^+ p ?$
$^{128}\text{Cd}$	T : average 16Du13=246.2(2.1) 15Lo04t=245(5)							
$^{128}\text{In}$	T : average 15Lo04=810(30) 86Go10=840(60)							
$^{128}\text{Sb}^m$	E : less than 20 keV above ground state, see ENSDF							
$^{128}\text{Te}^m$	T : average 04Va03=337(59) 98Zh09=370(30)							
$^{128}\text{Cs}$	T : average 93Al03=3.66(0.02) 76He04=3.62(0.02)							
$^{128}\text{La}$	T : average 97Ha30=5.4(0.2) 77Zo02=5.2(0.4) 66Pa06=4.9(0.4) 66Li04=4.9(0.4)							
$^{128}\text{Ce}$	T : average 00Li08=4.0(0.1) 97Ha30=4.1(0.3) 97As05=3.925(0.021)							
$^{128}\text{Pr}$	T : average 99Xi03=2.8(0.1) 88Ba42=3.1(0.3) 85Wi07=3.2(+0.5-0.4)							
$^{128}\text{Pr}$	D : from 85Wi07							
$^{128}\text{Nd}$	T : 83Ni05 gave 4(2) s. Proved, in 85Wi07, to be due to $^{128}\text{Pr}$ , not to $^{128}\text{Nd}$							
$^{128}\text{Pm}$	D : p=0% from 93Li40 J : from 02Xu11 and calculated $6^+$							
$^{129}\text{Pd}$	-37610#	600#	31 ms	7	$7/2^-$	15	2015	$\beta^- = 100$ ; $\beta^- n = 90\%$ ; $\beta^- 2n = 2\%$
$^{129}\text{Ag}$	-51980#	400#	49.9 ms	3.5	$7/2^+$	14 15Lo04 T	2000	$\beta^- = 100$ ; $\beta^- n = 10\%$
$^{129}\text{Ag}^m$	-51960#	400#	10# ms		$1/2^-$	14		$\beta^- ?$ ; $\beta^- n = 10\%$
$^{129}\text{Cd}$	-63058	17	151.5 ms	5.7	$3/2^+$	14 16Du13 T	1986	$\beta^- = 100$ ; $\beta^- n = 0.1\%$
$^{129}\text{Cd}^m$	-62910#	150#	147 ms	3	$11/2^-$	14 16Du13 T	2003	$\beta^- = 100$ ; $\beta^- n = 0.1\%$
$^{129}\text{Cd}^n$	-60970#	150#	3.6 ms	0.2	$(21/2^+)$	14 14Ta29 TJ	2014	IT=100
$^{129}\text{In}$	-72837.7	2.7	570 ms	10	$(9/2^+)$	14 15Lo04 T	1975	$\beta^- = 100$ ; $\beta^- n = 0.237$
$^{129}\text{In}^m$	-72380	3	1.23 s	0.03	$(1/2^-)$	14 04Ga24 J	1976	$\beta^- \approx 100$ ; IT<0.3; $\beta^- n = 3.64$
$^{129}\text{In}^n$	-71149.7	2.7	11.2 $\mu\text{s}$	0.2	$(17/2^-)$	14 14Ta.A T	2003	IT=100
$^{129}\text{In}^p$	-71180	50	670 ms	100	$(23/2^-)$	14 04Ga24 ETJ	2004	$\beta^- \approx 100$ ; IT ?
$^{129}\text{In}^q$	-70920	50	110 ms	15	$(29/2^+)$	14	2004	IT $\approx$ 100; $\beta^- ?$
$^{129}\text{Sn}$	-80591	17	2.23 m	0.04	$3/2^+$	14	1962	$\beta^- = 100$
$^{129}\text{Sn}^m$	-80556	17	6.9 m	0.1	$11/2^-$	14	1962	$\beta^- \approx 100$ ; IT<0.002
$^{129}\text{Sn}^n$	-78829	17	3.49 $\mu\text{s}$	0.11	$(19/2^+)$	14 08Lo07 T	2000	IT=100
$^{129}\text{Sn}^p$	-78788	17	2.22 $\mu\text{s}$	0.13	$23/2^+$	14 08Lo07 TJ	2000	IT=100
$^{129}\text{Sn}^q$	-78038	17	2.21 ns	18	$(27/2^-)$	14 08Lo07 J	2008	IT=100
$^{129}\text{Sb}$	-84629	21	4.366 h	0.026	$7/2^+$	14	1939	$\beta^- = 100$
$^{129}\text{Sb}^m$	-82778	21	17.7 m	0.1	$(19/2^-)$	14	1982	$\beta^- = 85$ ; IT=15
$^{129}\text{Sb}^n$	-82768	21	2.2 $\mu\text{s}$	0.2	$(15/2^-)$	14	1987	IT=100
$^{129}\text{Sb}^p$	-82490	21	1.1 $\mu\text{s}$	0.1	$(23/2^+)$	14	2003	IT=100
$^{129}\text{Te}$	-87004.8	0.9	69.6 m	0.3	$3/2^+$	14	1939	$\beta^- = 100$
$^{129}\text{Te}^m$	-86899.3	0.9	33.6 d	0.1	$11/2^-$	14	1940	IT=64.7; $\beta^- = 36.7$
$^{129}\text{I}$	-88507	3	15.7 My	0.4	$7/2^+$	14	1951	$\beta^- = 100$
$^{129}\text{Xe}$	-88696.059	0.005	STABLE		$1/2^+$	14	1920	IS=26.4006 82
$^{129}\text{Xe}^m$	-88459.92	0.03	8.88 d	0.02	$11/2^-$	14	1951	IT=100
$^{129}\text{Cs}$	-87499	5	32.06 h	0.06	$1/2^+$	14	1950	$\beta^+ = 100$
$^{129}\text{Cs}^m$	-86924	5	718 ns	21	$(11/2^-)$	14	1977	IT=100
$^{129}\text{Ba}$	-85063	11	2.23 h	0.11	$1/2^+$	14	1950	$\beta^+ = 100$
$^{129}\text{Ba}^m$	-85055	11	2.135 h	0.010	$7/2^+$	14	1950	$\beta^+ \approx 100$ ; IT=?
$^{129}\text{La}$	-81325	21	11.6 m	0.2	$(3/2^+)$	14	1963	$\beta^+ = 100$
$^{129}\text{La}^m$	-81153	21	560 ms	50	$(11/2^-)$	14	1969	IT=100
$^{129}\text{Ce}$	-76287	28	3.5 m	0.3	$(5/2^+)$	14	1977	$\beta^+ = 100$
$^{129}\text{Pr}$	-69774	30	30 s	4	$(3/2^+)$	14 96Gi08 J	1977	$\beta^+ = 100$
$^{129}\text{Pr}^m$	-69390	30	1# ms		$(11/2^-)$	14	1997	IT=100
$^{129}\text{Nd}$	-62320#	200#	6.8 s	0.6	$5/2^+$	14 10Xu12 T	1977	$\beta^+ = 100$ ; $\beta^+ p=?$
$^{129}\text{Nd}^m$	-62270#	220#	2.6 s	0.4	$1/2^+$	14	2010	$\beta^+ = 100$ ; $\beta^+ p=?$
$^{129}\text{Pm}$	-52880#	300#	2.4 s	0.9	$(5/2^-)$	14	2004	$\beta^+ = 100$ ; $\beta^+ p ?$ ; $p ?$
$^{129}\text{Sm}$	-42000#	500#	550 ms	100	$(3/2^+, 1/2^+)$	14	1999	$\beta^+ = 100$ ; $\beta^+ p=?$
$^{129}\text{Ag}$	T : average 15Lo04=52(4) 00Kr18=46(+5-9)							
$^{129}\text{Ag}$	D : $\beta^- n$ has been observed by 00Kr18 but not quantified							
$^{129}\text{Ag}^m$	T : 00Kr18 $\approx$ 160 ms is not convincing							
$^{129}\text{Cd}$	D : $\beta^- n$ has been observed by 05Kr20 but not quantified							
$^{129}\text{Cd}$	T : average 16Du13=157(8) 15Ta13=146(8)							
$^{129}\text{Cd}$	T : other 15Lo04=154.5(2.0) for mixture of two states							
$^{129}\text{Cd}^m$	T : other 15Ta13=151(15) ms							
$^{129}\text{Cd}^n$	D : $\beta^- n$ has been observed by 05Kr20 but not quantified							
$^{129}\text{Cd}^n$	E : 1940 keV above the $11/2^-$ isomer							
$^{129}\text{In}$	J : from 04Ga24							
$^{129}\text{In}^q$	E : 281.0 (0.2) keV $\gamma$ above the $23/2^-$ isomer							
$^{129}\text{Sn}^n$	T : average 08Lo07=3.4(0.4) 04Ga24=3.2(0.2) 00Pi03=3.7(0.2) 00Ge07=3.6(0.2)							
$^{129}\text{Sn}^p$	T : average 08Lo07=2.4(4) 04Ga24=2.0(2) 00Ge07=2.4(2)							
$^{129}\text{Nd}$	T : average 10Xu12=6.7(0.7) 97Gi07=7(1); 85Wi07=4.9(0.2) is for gs+m mixture							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>130</sup> Ag	-45700#	500#	40.6 ms	4.5		15 15Lo04	T	2000 $\beta^-$ =100; $\beta^-$ n=90#; $\beta^-$ 2n=2# *
<sup>130</sup> Cd	-61118	22	126.8 ms	1.8		08 16Du13	T	1986 $\beta^-$ =100; $\beta^-$ n=3.5 10 *
<sup>130</sup> Cd <sup>m</sup>	-58988	22	240 ns	16		08 12Ka36	ET	2007 IT=100 *
<sup>130</sup> In	-69880	40	284 ms	10		08 15Lo04	T	1973 $\beta^-$ =100; $\beta^-$ n=0.93 13 *
<sup>130</sup> In <sup>m</sup>	-69830	40	50	50	BD *			1973 $\beta^-$ =100; $\beta^-$ n=1.65 15 *
<sup>130</sup> In <sup>n</sup>	-69480	50	400	60	BD			1986 $\beta^-$ =100; $\beta^-$ n=1.65 15 *
<sup>130</sup> In <sup>p</sup>	-69490	40	388.3	0.2		08 12Ka36	T	2003 IT=100 *
<sup>130</sup> Sn	-80132.2	1.9	3.72 m	0.07				1972 $\beta^-$ =100
<sup>130</sup> Sn <sup>m</sup>	-78185.3	1.9	1946.88	0.10				$\beta^-$ =100
<sup>130</sup> Sn <sup>n</sup>	-77697.4	1.9	2434.79	0.12		01 11Pi05	T	1981 IT=100
<sup>130</sup> Sb	-82286	14	39.5 m	0.8		01 02Ge07	J	1962 $\beta^-$ =100
<sup>130</sup> Sb <sup>m</sup>	-82281	14	4.80	0.20				$\beta^-$ =100
<sup>130</sup> Sb <sup>n</sup>	-82201	14	84.67	0.04		01 02Ge07	TJ	2002 IT=100
<sup>130</sup> Sb <sup>p</sup>	-80741	14	1544.7	0.5		02Ge07	ETJ	2002 IT=100
<sup>130</sup> Te	-87352.949	0.011	690 Ey	130		01 15Ba11	T	1924 IS=34.08 62; 2 $\beta^-$ =100 *
<sup>130</sup> Te <sup>m</sup>	-85206.54	0.04	186 ns	11		01 04Va03	T	1972 IT=100 *
<sup>130</sup> Te <sup>n</sup>	-84685.7	0.8	1.90 $\mu$ s	0.08		01 04Br19	E	1998 IT=100 *
<sup>130</sup> Te <sup>p</sup>	-82977.5	1.8	261 ns	33				1998 IT=100
<sup>130</sup> I	-86936	3	12.36 h	0.01				1938 $\beta^-$ =100
<sup>130</sup> I <sup>m</sup>	-86896	3	39.9525	0.0013				1966 IT=84 2; $\beta^-$ =16 2
<sup>130</sup> I <sup>n</sup>	-86866	3	69.5865	0.0007				1989 IT=100
<sup>130</sup> I <sup>p</sup>	-86854	3	82.3960	0.0019				1989 IT=100
<sup>130</sup> I <sup>q</sup>	-86851	3	85.1099	0.0010				1975 IT=100
<sup>130</sup> Xe	-89880.463	0.009	STABLE					1922 IS=4.0710 13
<sup>130</sup> Cs	-86900	8	29.21 m	0.04				$\beta^+$ =98.4; $\beta^-$ =1.6
<sup>130</sup> Cs <sup>m</sup>	-86737	8	3.46 m	0.06				1977 IT $\approx$ 100; $\beta^+$ =0.16 2
<sup>130</sup> Cs <sup>s</sup>	-86873	17	27	15	R = .2 .1			fsmix
<sup>130</sup> Ba	-87261.5	2.6	STABLE	1Zy				01 15Ba11 T 1936 IS=0.106 1; 2 $\beta^+$ ?
<sup>130</sup> Ba <sup>m</sup>	-84786.4	2.6	9.54 ms	0.14				01 02Mo31 T 1969 IT=100 *
<sup>130</sup> La	-81627	26	8.7 m	0.1				1961 $\beta^+$ =100
<sup>130</sup> La <sup>m</sup>	-81413	26	760 ns	90		14Io01	ETJ	2012 IT=100
<sup>130</sup> La <sup>n</sup>	-81308	26	33 ns	1		14Io01	ETJ	2014 IT=100
<sup>130</sup> Ce	-79423	28	22.9 m	0.5				1965 $\beta^+$ =100
<sup>130</sup> Ce <sup>m</sup>	-76969	28	100 ns	8				1999 IT=100
<sup>130</sup> Pr	-71180	60	40.0 s	0.4				(6,7) <sup>(+)</sup>
<sup>130</sup> Pr <sup>m</sup>	-71080#	120#	10# s			01 88Ba42	J	1977 $\beta^+$ =100
<sup>130</sup> Nd	-66596	28	21 s	3		01 88Ba42	J	1988 $\beta^+$ ? *
<sup>130</sup> Pm	-55400#	200#	2.6 s	0.2		01 01Gi17	T	1977 $\beta^+$ =100 *
<sup>130</sup> Sm	-47510#	400#	1# s			01 99Xi03	J	1985 $\beta^+$ =100; $\beta^+$ p=?
<sup>130</sup> Eu	-33680#	500#	1.0 ms	0.4				08 $\beta^+$ ?
* <sup>130</sup> Ag	T : average 15Lo04=42(5) 05Kr20=35(10) **							
* <sup>130</sup> Cd	T : average 16Du13=126(4) 15Lo04=127(2) **							
* <sup>130</sup> Cd <sup>m</sup>	T : average 12Ka36=248(+21-19) 07Ju05=220(30) **							
* <sup>130</sup> Cd <sup>n</sup>	E : 12Ka36=128.0(0.5), 138.0(0.5), 538.2(0.5) and 1325.4(0.5) $\gamma$ rays in cascade to ground-state **							
* <sup>130</sup> In <sup>p</sup>	E : other 12Ka36=388.5(0.5) **							
* <sup>130</sup> In <sup>p</sup>	T : symmetrized from 12Ka36=5.25(+0.40-0.35); other 04Sc42=3.1(0.3) **							
* <sup>130</sup> Te	T : 15Al20 : 0v- $\beta\beta$ >2700 Zy **							
* <sup>130</sup> Te <sup>m</sup>	T : other conflicting data: 72Ke28=115(11) J : E1 to 6 <sup>+</sup> , E2 to 4 <sup>+</sup> **							
* <sup>130</sup> Te <sup>n</sup>	E : other: less than 25 keV above 2648.57(0.22) (8 <sup>+</sup> ) level, see ENSDF'01 **							
* <sup>130</sup> Te <sup>n</sup>	T : other conflicting data, not used: 98Zh09=4.2(0.9) $\mu$ s **							
* <sup>130</sup> Ba <sup>m</sup>	T : others 66Br14=8.8(0.2) 69Wa.A=13.5(1.0) not used **							
* <sup>130</sup> Pr <sup>m</sup>	J : 88Ba42: there is also a low-spin component in <sup>130</sup> Pr activity **							
* <sup>130</sup> Pr <sup>m</sup>	J : see also the discussion in 01Gi17 on three isomeric states in <sup>130</sup> Pr **							
* <sup>130</sup> Nd	T : other 00Xu08=13(3) 77Bo02=28(3) conflicting, not used **							
* <sup>130</sup> Eu	T : symmetrized from 0.90(+0.49-0.29) D : estim from $\beta^+$ half-live=49#ms **							
<sup>131</sup> Ag	-40380#	500#	35 ms	8				2013 $\beta^-$ =100; $\beta^-$ n=90#; $\beta^-$ 2n=10
<sup>131</sup> Cd	-55220	100	98 ms	2		06 15Lo04	T	2000 $\beta^-$ =100; $\beta^-$ n=3.5 10; $\beta^-$ 2n=0# *
<sup>131</sup> In	-68025.0	2.7	261 ms	3		06 15Lo04	T	1976 $\beta^-$ =100; $\beta^-$ n=2.2 3 *
<sup>131</sup> In <sup>m</sup>	-67660	7	365	8	MD			1984 $\beta^-$ $\approx$ 100; $\beta^-$ n<2.0 3; IT<0.018
<sup>131</sup> In <sup>n</sup>	-64280	90	320 ms	60	BD			1984 $\beta^-$ >99; $\beta^-$ n=0.028 5; IT<1
<sup>131</sup> In <sup>p</sup>	-64241.4	2.7	3783.6	0.5				09Go40 TJ 2009 IT=100 *
<sup>131</sup> Sn	-77265	4	56.0 s	0.5				06 05Le34 J 1963 $\beta^-$ =100
<sup>131</sup> Sn <sup>m</sup>	-77200	4	65.1	0.3				06 04Fo06 E 1977 $\beta^-$ =100; IT<0.0004# *
<sup>131</sup> Sn <sup>n</sup>	-72595	4	4670.0	0.3				06 12Ka36 T 2001 IT=100 *
<sup>131</sup> Sb	-81981.4	2.1	23.03 m	0.04				06 1956 $\beta^-$ =100
<sup>131</sup> Sb <sup>m</sup>	-80305.3	2.1	91 $\mu$ s	4				06 15/2 <sup>-</sup> # 1969 IT=100
<sup>131</sup> Sb <sup>n</sup>	-80294.2	2.3	4.3 $\mu$ s	0.8				06 (19/2 <sup>-</sup> ) 2000 IT=100
<sup>131</sup> Sb <sup>p</sup>	-79815.8	2.6	1.1 $\mu$ s	0.2				06 (23/2 <sup>+</sup> ) 2000 IT=100

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...												
<sup>131</sup> Te	-85211.01	0.06			25.0 m	0.1	3/2 <sup>+</sup>	06		1939	$\beta^-$ =100	
<sup>131</sup> Te <sup>m</sup>	-85028.75	0.06	182.258	0.018	32.48 h	0.11	11/2 <sup>-</sup>	06	08Ea01	T	1940	$\beta^-$ =74.1 5; IT=25.9 5
<sup>131</sup> Te <sup>n</sup>	-83271.0	0.4	1940.0	0.4	93 ms	12	(23/2 <sup>+</sup> )	06			1998	IT=100
<sup>131</sup> I	-87442.7	0.6			8.0252 d	0.0006	7/2 <sup>+</sup>	06			1939	$\beta^-$ =100
<sup>131</sup> I <sup>m</sup>	-85524.3	0.7	1918.4	0.42	24 $\mu$ s	1	19/2 <sup>-</sup>	06	09Wa11	EJT	2009	IT=100
<sup>131</sup> Xe	-88413.558	0.009			STABLE		3/2 <sup>+</sup>	06			1920	IS=21.2324 30
<sup>131</sup> Xe <sup>m</sup>	-88249.628	0.012	163.930	0.008	11.84 d	0.04	11/2 <sup>-</sup>	06			1966	IT=100
<sup>131</sup> Cs	-88059	5			9.689 d	0.016	5/2 <sup>+</sup>	06			1947	$\epsilon$ =100
<sup>131</sup> Ba	-86683.7	2.6			11.52 d	0.01	1/2 <sup>+</sup>	06	12Da04	T	1947	$\beta^+$ =100
<sup>131</sup> Ba <sup>m</sup>	-86495.7	2.6	187.995	0.009	14.26 m	0.09	9/2 <sup>-</sup>	06	12Da04	T	1963	IT=100
<sup>131</sup> La	-83769	28			59 m	2	3/2 <sup>+</sup>	06			1951	$\beta^+$ =100
<sup>131</sup> La <sup>m</sup>	-83464	28	304.60	0.24	170 $\mu$ s	7	11/2 <sup>-</sup>	06			1966	IT=100
<sup>131</sup> Ce	-79710	30			10.3 m	0.3	7/2 <sup>+</sup>	06			1966	$\beta^+$ =100
<sup>131</sup> Ce <sup>m</sup>	-79650	30	63.09	0.09	5.4 m	0.4	(1/2 <sup>+</sup> )	06	96Gi08	E	1966	$\beta^+$ =100
<sup>131</sup> Pr	-74300	50			1.50 m	0.03	3/2 <sup>+</sup> #	06	96Gi08	T	1977	$\beta^+$ =100
<sup>131</sup> Pr <sup>m</sup>	-74150	50	152.4	0.3	5.73 s	0.20	(11/2 <sup>-</sup> )	06			1996	IT=96.4 12; $\beta^+$ =3.6 12
<sup>131</sup> Nd	-67768	28			25.4 s	0.9	(5/2 <sup>+</sup> ) <sup>(+)</sup> #	06			1977	$\beta^+$ =100; $\beta^+$ p=?
<sup>131</sup> Pm	-59660#	200#			6.3 s	0.8	(11/2 <sup>-</sup> )	06	99Ga41	T	1998	$\beta^+$ =100
<sup>131</sup> Sm	-50130#	400#			1.2 s	0.2	5/2 <sup>+</sup> #	06			1986	$\beta^+$ =100; $\beta^+$ p=?
<sup>131</sup> Eu	-39270#	400#			17.8 ms	1.9	3/2 <sup>+</sup>	06			1998	p=89 9; $\beta^+$ ?; $\beta^+$ p ?
* <sup>131</sup> In	D: from 93Ru01										**	
* <sup>131</sup> In <sup>p</sup>	E: other 12Ka36=3783.6(0.5)										**	
* <sup>131</sup> In <sup>p</sup>	T: average 12Ka36=685(+42-39) 09Go40=630(60)				J: from 09Go40							**
* <sup>131</sup> Sn <sup>m</sup>	J: from 05Le34										**	
* <sup>131</sup> Sn <sup>n</sup>	E: 4605.02(0.21) above the 58.4 s 11/2 <sup>-</sup> level										**	
* <sup>131</sup> Sn <sup>n</sup>	T: average 12Ka36=309(+24-23) 84Fo19=300(20)										**	
* <sup>131</sup> I <sup>m</sup>	E: derived from least-squares fit to $\gamma$ -ray energies										**	
* <sup>131</sup> Pr	T: average 96Gi08=1.57(0.07) 93Al03=1.48(0.02) 83Ga.A=1.58(0.05)										**	
<sup>132</sup> Ag	-33790#	500#			30 ms	14		15	15Lo04	TD	2015	$\beta^-$ =100; $\beta^-$ n=0#; $\beta^-$ 2n=90#
<sup>132</sup> Cd	-50260#	200#			82 ms	4	0 <sup>+</sup>	05	15Lo04	T	2000	$\beta^-$ =100; $\beta^-$ n=60 15; $\beta^-$ 2n=0.2#
<sup>132</sup> In	-62410	60			198 ms	2	(7 <sup>-</sup> )	05	15Lo04	T	1973	$\beta^-$ =100; $\beta^-$ n=6.3 9; $\beta^-$ 2n=0#
<sup>132</sup> Sn	-76546.5	2.0			39.7 s	0.8	0 <sup>+</sup>	05			1963	$\beta^-$ =100
<sup>132</sup> Sn <sup>m</sup>	-71698.0	2.0	4848.52	0.20	2.079 $\mu$ s	0.016	(8 <sup>+</sup> )	05	12Ka36	T	1986	IT=100
<sup>132</sup> Sb	-79635.3	2.5			2.79 m	0.07	(4 <sup>+</sup> )	05			1956	$\beta^-$ =100
<sup>132</sup> Sb <sup>m</sup>	-79440	30	200	30	4.10 m	0.05	(8 <sup>-</sup> )	05	89St06	E	1956	$\beta^-$ =100
<sup>132</sup> Sb <sup>n</sup>	-79380.8	2.5	254.5	0.3	102 ns	4	(6 <sup>-</sup> )	05			1974	IT=100
<sup>132</sup> Te	-85188	3			3.204 d	0.013	0 <sup>+</sup>	05			1948	$\beta^-$ =100
<sup>132</sup> Te <sup>m</sup>	-83413	3	1774.80	0.09	145 ns	8	6 <sup>+</sup>	05			1973	IT=100
<sup>132</sup> Te <sup>n</sup>	-83263	3	1925.47	0.09	28.1 $\mu$ s	1.5	7 <sup>-</sup>	05	FGK128	J	1979	IT=100
<sup>132</sup> Te <sup>p</sup>	-82465	3	2723.3	0.8	3.70 $\mu$ s	0.09	(10 <sup>+</sup> )	05			1979	IT=100
<sup>132</sup> I	-85703	4			2.295 h	0.013	4 <sup>+</sup>	05			1948	$\beta^-$ =100
<sup>132</sup> I <sup>m</sup>	-85594	10	110	11	1.387 h	0.015	(8 <sup>-</sup> )	05			1973	IT=86 2; $\beta^-$ =14 2
<sup>132</sup> Xe	-89278.962	0.005			STABLE		0 <sup>+</sup>	05			1920	IS=26.9086 33
<sup>132</sup> Xe <sup>m</sup>	-86526.75	0.17	2752.21	0.17	8.39 ms	0.11	(10 <sup>+</sup> )	05			1976	IT=100
<sup>132</sup> Cs	-87152.7	1.0			6.480 d	0.006	2 <sup>+</sup>	05			1953	$\beta^+$ =98.13 9; $\beta^+$ =1.87 9
<sup>132</sup> Ba	-88435.0	1.1			STABLE	(>300 Ey)	0 <sup>+</sup>	05	96Ba24	T	1936	IS=0.101 1; 2 $\beta^+$ ?
<sup>132</sup> La	-83720	40			4.8 h	0.2	2 <sup>-</sup>	05			1951	$\beta^+$ =100
<sup>132</sup> La <sup>m</sup>	-83530	40	188.20	0.11	24.3 m	0.5	6 <sup>-</sup>	05			1969	IT=76; $\beta^+$ =24
<sup>132</sup> Ce	-82471	20			3.51 h	0.11	0 <sup>+</sup>	05			1960	$\beta^+$ =100
<sup>132</sup> Ce <sup>m</sup>	-80130	20	2341.15	0.21	9.4 ms	0.3	8 <sup>-</sup>	05	09Pe31	J	1969	IT=100
<sup>132</sup> Pr	-75227	29			1.49 m	0.11	(2 <sup>+</sup> )	05	94Bu18	TJ	1974	$\beta^+$ =100
<sup>132</sup> Pr <sup>m</sup>	-75200#	40#	30#	30#	1#	s	(5 <sup>+</sup> )	05	90Ko25	J	1990	$\beta^+$ ?
<sup>132</sup> Pr <sup>n</sup>	-74980#	40#	250#	30#	2.46 $\mu$ s	0.04	(8 <sup>+</sup> )	05	12Ta18	TJD	2012	IT=100
<sup>132</sup> Pr <sup>p</sup>	-74960#	100#	270#	100#	486 ns	70	(8 <sup>-</sup> )	05	12Ta18	TJD	2012	IT=100
<sup>132</sup> Nd	-71426	24			1.56 m	0.10	0 <sup>+</sup>	05	95Bu11	T	1977	$\beta^+$ =100
<sup>132</sup> Pm	-61630#	150#			6.2 s	0.6	(3 <sup>+</sup> )	05			1977	$\beta^+$ =100; $\beta^+$ p $\approx$ 5e-5
<sup>132</sup> Sm	-55080#	300#			4.0 s	0.3	0 <sup>+</sup>	05			1989	$\beta^+$ =100; $\beta^+$ p ?
<sup>132</sup> Eu	-42200#	400#			100# ms			05	93Li40	D		$\beta^+$ ?; $\beta^+$ p ?; p=0
* <sup>132</sup> Ag	T: symmetrized from 15Lo04=28(+15-12)										**	
* <sup>132</sup> Sn <sup>m</sup>	T: average 12Ka36=2.088(0.017) 94Fo14=2.03(4); other 82Ka25=1.7(2)										**	
* <sup>132</sup> Te <sup>n</sup>	J: E1 to 6 <sup>+</sup>										**	
* <sup>132</sup> Pr	T: average 94Bu18=1.47(0.12) 74Ar27=1.6(0.3)										**	
* <sup>132</sup> Pr <sup>n</sup>	E: 12Ta18=219.9(0.14) keV above (5 <sup>+</sup> ) isomer										**	
* <sup>132</sup> Pr <sup>p</sup>	E: 12Ta18=273.0(0.14) keV above (5 <sup>+</sup> ) isomer										**	
* <sup>132</sup> Nd	T: average 95Bu11=1.47(0.12) 77Bo02=1.75(0.17)										**	



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>133</sup> Cd	-43920#	300#	61 ms	6	7/2 <sup>-</sup> #	11 15Lo04 T	2010	$\beta^-$ =100; $\beta^-$ n=0.5#; $\beta^-$ 2n=90# *	
<sup>133</sup> In	-57460#	200#	165 ms	3	(9/2 <sup>+</sup> )	11 96Ho16 J	1996	$\beta^-$ =100; $\beta^-$ n=85 10; $\beta^-$ 2n=0.4# *	
<sup>133</sup> In <sup>m</sup>	-57130#	200#	180# ms		(1/2 <sup>-</sup> )	11 96Ho16 J	1996	IT ?; $\beta^-$ ? *	
<sup>133</sup> Sn	-70873.9	1.9	1.46 s	0.03	(7/2 <sup>-</sup> )	11	1973	$\beta^-$ =100; $\beta^-$ n=0.0294 24	
<sup>133</sup> Sb	-78924	3	2.34 m	0.05	7/2 <sup>+</sup> #	11	1966	$\beta^-$ =100	
<sup>133</sup> Sb <sup>m</sup>	-74360	100	16.54 $\mu$ s	0.19	(21/2 <sup>+</sup> )	11	1978	IT=100	
<sup>133</sup> Te	-82937.1	2.1	12.5 m	0.3	3/2 <sup>+</sup> #	11	1940	$\beta^-$ =100	
<sup>133</sup> Te <sup>m</sup>	-82602.8	2.1	334.26	0.04	(11/2 <sup>-</sup> )	11	1957	$\beta^-$ =83.5 20; IT=16.5 20	
<sup>133</sup> Te <sup>n</sup>	-81326.7	2.2	1610.4	0.5	(19/2 <sup>-</sup> )	11	2001	IT=100	
<sup>133</sup> I	-85858	6	20.83 h	0.08	7/2 <sup>+</sup>	11	1940	$\beta^-$ =100	
<sup>133</sup> I <sup>m</sup>	-84224	6	1634.148	0.010	9 s	2	(19/2 <sup>-</sup> )	11 1970 IT=100	
<sup>133</sup> I <sup>n</sup>	-84129	6	1729.137	0.010	170 ns	(15/2 <sup>-</sup> )	11 1984 IT=100		
<sup>133</sup> I <sup>p</sup>	-83423	6	2435.00	0.23	780 ns	1606	(19/2 <sup>+</sup> )	11 2004 IT=100	
<sup>133</sup> I <sup>q</sup>	-83364	6	2493.7	0.4	469 ns	15	(23/2 <sup>+</sup> )	11 2009 IT=100	
<sup>133</sup> Xe	-87643.6	2.4	5.2475 d	0.0005	3/2 <sup>+</sup>	11 02Un02 T	1940	$\beta^-$ =100	
<sup>133</sup> Xe <sup>m</sup>	-87410.4	2.4	2.198 d	0.013	11/2 <sup>-</sup>	11	1951	IT=100	
<sup>133</sup> Cs	-88070.931	0.008	STABLE		7/2 <sup>+</sup>	11	1921	IS=100.	
<sup>133</sup> Ba	-87553.6	1.0	10.551 y	0.011	1/2 <sup>+</sup>	11	1941	$\epsilon$ =100	
<sup>133</sup> Ba <sup>m</sup>	-87265.3	1.0	288.252	0.009	38.90 h	0.06	11/2 <sup>-</sup>	11 1941 IT $\approx$ 100; $\epsilon$ =0.0104 5 *	
<sup>133</sup> La	-85494	28	3.912 h	0.008	5/2 <sup>+</sup>	11	1950	$\beta^+$ =100	
<sup>133</sup> Ce	-82418	16	97 m	4	1/2 <sup>+</sup>	11	1951	$\beta^+$ =100	
<sup>133</sup> Ce <sup>m</sup>	-82381	16	37.2	0.7	5.1 h	0.3	9/2 <sup>-</sup>	11 1951 $\beta^+$ =100	
<sup>133</sup> Pr	-77938	12	6.5 m	0.3	(3/2 <sup>+</sup> )	11	1970	$\beta^+$ =100	
<sup>133</sup> Pr <sup>m</sup>	-77746	12	192.12	0.14	1.1 s	0.2	(11/2 <sup>-</sup> )	11 1995 IT=100	
<sup>133</sup> Nd	-72330	50	70 s	10	(7/2 <sup>+</sup> )	11	1977	$\beta^+$ =100	
<sup>133</sup> Nd <sup>m</sup>	-72200	50	127.97	0.12	70 s	(1/2 <sup>+</sup> )	11 95Br24 D	1993 $\beta^+$ $\approx$ 100; IT=?	
<sup>133</sup> Nd <sup>n</sup>	-72150	50	176.10	0.10	301 ns	18	(9/2 <sup>-</sup> )	11 1993 IT=100	
<sup>133</sup> Pm	-65410	50	13.5 s	2.1	(3/2 <sup>+</sup> )	11	1977	$\beta^+$ =100	
<sup>133</sup> Pm <sup>m</sup>	-65280	50	129.7	0.7	8# s	(11/2 <sup>-</sup> )	11 1996 $\beta^+$ ?; IT ?		
<sup>133</sup> Sm	-57230#	300#	2.89 s	0.16	(5/2 <sup>+</sup> )	11	1977	$\beta^+$ =100; $\beta^+$ p=?	
<sup>133</sup> Sm <sup>m</sup>	-57110#	310#	120#	60#	3.5 s	0.4	(1/2 <sup>-</sup> )	11 1993 $\beta^+$ ?; IT ?; $\beta^+$ p ?	
<sup>133</sup> Eu	-47240#	300#	200# ms		11/2 <sup>-</sup> #			$\beta^+$ ?; $\beta^+$ p ?	
<sup>133</sup> Gd	-35860#	500#	10# ms		5/2 <sup>+</sup> #			$\beta^+$ ?; $\beta^+$ p ?	
* <sup>133</sup> Cd	T : average 15Lo04=64(8) 05Kr20=57(10)								**
* <sup>133</sup> Cd	D : delayed neutrons were observed in 05Kr20								**
* <sup>133</sup> In	T : average 15Lo04=163(7) 02D112=165(3)								**
* <sup>133</sup> Ba <sup>m</sup>	T : average 12Da04=38.88(0.08) 11Gr01=38.92(0.09)								**
<sup>134</sup> Cd	-38920#	400#	65 ms	15	0 <sup>+</sup>	15	2015	$\beta^-$ =100; $\beta^-$ n=0.2#; $\beta^-$ 2n=90#	
<sup>134</sup> In	-51660#	300#	140 ms	4	high	04 95Jo.A D	1996	$\beta^-$ =100; $\beta^-$ n=65; $\beta^-$ 2n<4 *	
<sup>134</sup> Sn	-66434	3	890 ms	20	0 <sup>+</sup>	04 15Lo04 T	1974	$\beta^-$ =100; $\beta^-$ n=17 13	
<sup>134</sup> Sn <sup>m</sup>	-65187	3	87 ns	8	6 <sup>+</sup>	04 12Ka36 T	2000	IT=100 *	
<sup>134</sup> Sb	-74020.5	1.7	780 ms	60	(0 <sup>-</sup> )	11	1967	$\beta^-$ =100; $\beta^-$ n=5#	
<sup>134</sup> Sb <sup>m</sup>	-73741.5	2.0	10.07 s	0.05	(7 <sup>-</sup> )	11	1968	$\beta^-$ =100; $\beta^-$ n=0.088 4	
<sup>134</sup> Te	-82533.7	2.7	41.8 m	0.8	0 <sup>+</sup>	04	1948	$\beta^-$ =100	
<sup>134</sup> Te <sup>m</sup>	-80842.4	2.7	1691.34	0.16	164.1 ns	0.9	6 <sup>+</sup>	04 1970 IT=100	
<sup>134</sup> I	-84043	5	52.5 m	0.2	(4 <sup>+</sup> )	04	1948	$\beta^-$ =100	
<sup>134</sup> I <sup>m</sup>	-83727	5	3.52 m	0.04	(8 <sup>-</sup> )	04	1970	IT=97.7 10; $\beta^-$ =2.3 10	
<sup>134</sup> Xe	-88125.822	0.009	STABLE	(>11 Py)	0 <sup>+</sup>	04 89Ba22 T	1920	IS=10.4357 21; 2 $\beta^-$ ? *	
<sup>134</sup> Xe <sup>m</sup>	-86160.3	0.5	1965.5	0.5	290 ms	17	(7 <sup>-</sup> )	04 1968 IT=100	
<sup>134</sup> Xe <sup>n</sup>	-85100.6	1.5	3025.2	1.5	5 $\mu$ s	1	(10 <sup>+</sup> )	04 2001 IT=100	
<sup>134</sup> Cs	-86891.154	0.016	2.0652 y	0.0004	4 <sup>+</sup>	04	1940	$\beta^-$ =100; $\epsilon$ =0.0003 1	
<sup>134</sup> Cs <sup>m</sup>	-86752.410	0.016	138.7441	0.0026	2.912 h	0.002	8 <sup>-</sup>	04 1975 IT=100	
<sup>134</sup> Ba	-88949.9	0.3	STABLE		0 <sup>+</sup>	04	1936	IS=2.417 18	
<sup>134</sup> Ba <sup>m</sup>	-85992.7	0.6	2957.2	0.5	2.63 $\mu$ s	0.14	(10 <sup>+</sup> )	04 1982 IT=100	
<sup>134</sup> La	-85219	20	6.45 m	0.16	1 <sup>+</sup>	04	1951	$\beta^+$ =100	
<sup>134</sup> La <sup>m</sup>	-84780#	100#	29 $\mu$ s	4		04	1985	IT=100 *	
<sup>134</sup> Ce	-84833	20	3.16 d	0.04	0 <sup>+</sup>	04	1951	$\epsilon$ =100	
<sup>134</sup> Ce <sup>m</sup>	-81624	20	3208.6	0.4	308 ms	5	10 <sup>+</sup>	04 1980 IT=100	
<sup>134</sup> Pr	-78528	20	17 m	2	2 <sup>-</sup>	04	1967	$\beta^+$ =100	
<sup>134</sup> Pr <sup>m</sup>	-78460	20	68	1	11 m	(6 <sup>-</sup> )	04 11Ti10 E	1973 $\beta^+$ =100; IT $\approx$ 0	
<sup>134</sup> Nd	-75646	12	8.5 m	1.5	0 <sup>+</sup>	04	1970	$\beta^+$ =100	
<sup>134</sup> Nd <sup>m</sup>	-73353	12	2293.0	0.4	410 $\mu$ s	30	(8 <sup>-</sup> )	04 1969 IT=100	
<sup>134</sup> Pm	-66740	60	* &	22 s	1	(5 <sup>+</sup> )	04	1977 $\beta^+$ =100	
<sup>134</sup> Pm <sup>m</sup>	-66740#	120#	0#	100#	5 s	(2 <sup>+</sup> )	04	1988 $\beta^+$ =100	
<sup>134</sup> Pm <sup>n</sup>	-66620#	80#	120#	50#	20 $\mu$ s	1	(7 <sup>-</sup> )	04 09Cu02 TJ	2009 IT=100 *
<sup>134</sup> Sm	-61380#	200#	9.5 s	0.8	0 <sup>+</sup>	04	1977	$\beta^+$ =100	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>134</sup> Eu	-49930# 300#		500 ms 200			04	1989	$\beta^+=100; \beta^+p=?$
<sup>134</sup> Gd	-41300# 400#		400# ms	0+		04		$\beta^+ ?; \beta^+p ?$
* <sup>134</sup> In	T: other 15Lo04=126(7)							
* <sup>134</sup> In	D: $\beta^-2n$ intensity limits is from 95Jo.A							
* <sup>134</sup> Sn <sup>m</sup>	T: symmetrized from 12Ka36=86(+8-7); other 00Ko15=80(15)							
* <sup>134</sup> Xe	D: and 89Ba22: 0v- $\beta\beta>58Zy$ and $>26Zy$ for $0^+ \rightarrow 0^+$ and $0^+ \rightarrow 2^+$ respectively							
* <sup>134</sup> La <sup>m</sup>	E: 100#100 keV above 336.44(17) level							
* <sup>134</sup> Pm <sup>m</sup>	E: 70.7(0.2) keV above a $6^+$ state that decays via a low-energy $\gamma$ to $5^+$							
<sup>135</sup> In	-46530# 400#		101 ms 4	9/2 <sup>+</sup> #		08 15Lo04 T	2002	$\beta^- ?; \beta^-n=90\#; \beta^-2n=8\#$
<sup>135</sup> Sn	-60632 3		515 ms 5	7/2 <sup>-</sup> #		08 15Lo04 T	1994	$\beta^- =100; \beta^-n=21 3; \beta^-2n=6\#$
<sup>135</sup> Sb	-69690.3 2.6		1.679 s 0.015	(7/2 <sup>+</sup> )		08	1964	$\beta^- =100; \beta^-n=22 3$
<sup>135</sup> Te	-77728.8 1.7		19.0 s 0.2	(7/2 <sup>-</sup> )		08	1969	$\beta^- =100$
<sup>135</sup> Te <sup>m</sup>	-76173.9 1.7	1554.89 0.16	511 ns 20	(19/2 <sup>-</sup> )		08	1980	IT=100
<sup>135</sup> I	-83779.1 2.1		6.58 h 0.03	7/2 <sup>+</sup>		08	1940	$\beta^- =100$
<sup>135</sup> Xe	-86413 4		9.14 h 0.02	3/2 <sup>+</sup>		08	1940	$\beta^- =100$
<sup>135</sup> Xe <sup>m</sup>	-85886 4	526.551 0.013	15.29 m 0.05	11/2 <sup>-</sup>		08	1960	IT $\approx$ 100; $\beta^- =0.30 17$
<sup>135</sup> Cs	-87581.6 1.0		1.33 My 0.19	7/2 <sup>+</sup>		08 16Ma05 T	1949	$\beta^- =100$
<sup>135</sup> Cs <sup>m</sup>	-85948.7 1.8	1632.9 1.5	53 m 2	19/2 <sup>-</sup>		08	1962	IT=100
<sup>135</sup> Ba	-87850.5 0.3		STABLE	3/2 <sup>+</sup>		08	1932	IS=6.592 12
<sup>135</sup> Ba <sup>m</sup>	-87582.3 0.3	268.218 0.020	28.11 h 0.02	11/2 <sup>-</sup>		08 12Da04 T	1948	IT=100
<sup>135</sup> La	-86643 9		19.5 h 0.2	5/2 <sup>+</sup>		08	1948	$\beta^+ =100$
<sup>135</sup> Ce	-84616 10		17.7 h 0.3	1/2 <sup>(+)</sup>		08	1948	$\beta^+ =100$
<sup>135</sup> Ce <sup>m</sup>	-84170 10	445.81 0.21	20 s 1	(11/2 <sup>-</sup> )		08	1963	IT=100
<sup>135</sup> Pr	-80936 12		24 m 1	3/2 <sup>(+)</sup>		08	1954	$\beta^+ =100$
<sup>135</sup> Pr <sup>m</sup>	-80578 12	358.06 0.06	105 $\mu$ s 10	(11/2 <sup>-</sup> )		08	1973	IT=100
<sup>135</sup> Nd	-76214 19		12.4 m 0.6	9/2 <sup>(-)</sup>		08	1970	$\beta^+ =100$
<sup>135</sup> Nd <sup>m</sup>	-76149 19	64.95 0.24	5.5 m 0.5	(1/2 <sup>+</sup> )		08	1970	$\beta^+ >99.97; IT <0.03$
<sup>135</sup> Pm	-70050 80		49 s 3	(5/2 <sup>+</sup> , 3/2 <sup>+</sup> )		08	1975	$\beta^+ =100$
<sup>135</sup> Pm <sup>m</sup>	-69830# 50#	220# 90#	40 s 3	(11/2 <sup>-</sup> )		08 89Ko07 TJ	1989	$\beta^+ =100$
<sup>135</sup> Sm	-62860 150		10.3 s 0.5	(7/2 <sup>+</sup> )		08 77Bo02 J	1977	$\beta^+ >99.97; \beta^+p=0.02 1$
<sup>135</sup> Sm <sup>m</sup>	-62860# 340#	0# 300#	2.4 s 0.9	(3/2 <sup>+</sup> , 5/2 <sup>+</sup> )		08 89Vi04 TJD	1989	$\beta^+ =100$
<sup>135</sup> Eu	-54150# 200#		1.5 s 0.2	11/2 <sup>-</sup> #		08	1989	$\beta^+ =100; \beta^+p ?$
<sup>135</sup> Gd	-44390# 400#		1.1 s 0.2	(5/2 <sup>+</sup> )		08	1996	$\beta^+ =100; \beta^+p=18$
<sup>135</sup> Tb	-32830# 400#		1.01 ms 0.28	(7/2 <sup>-</sup> )		08	2004	$p \approx 100; \beta^+ ?$
* <sup>135</sup> In	T: average 15Lo04=103(5) 02Di12=92(10)							
* <sup>135</sup> In	D: delayed neutrons were observed in 02Di12							
* <sup>135</sup> Xe <sup>m</sup>	D: $\beta^-$ ranging from 0.004% to 0.6%							
* <sup>135</sup> Cs	T: average 16Ma05=1.6(0.6) by AMS and 1.3(0.2) ICPMS							
* <sup>135</sup> Pm <sup>m</sup>	E: Trends of 11/2 <sup>-</sup> level in Pm isotopes: <sup>133</sup> Pm: 129.7(0.7) <sup>135</sup> Pm: 150#50							
* <sup>135</sup> Pm <sup>m</sup>	E: <sup>137</sup> Pm: 150(50) <sup>139</sup> Pm: 188.7(0.3) <sup>141</sup> Pm: 628.40(0.10) <sup>143</sup> Pm: 959.7(0.1)							
* <sup>135</sup> Pm <sup>m</sup>	E: ( $N > 82$ ) <sup>145</sup> Pm: 794.6(0.4) <sup>147</sup> Pm: 649.3(0.4) <sup>149</sup> Pm: 240.215(0.007)							
* <sup>135</sup> Pm <sup>m</sup>	E: ENSDF2008 : 68.7 + y							
* <sup>135</sup> Sm <sup>m</sup>	I: existence of <sup>135</sup> Sm <sup>m</sup> and spins of both states are discussed in ENSDF							
* <sup>135</sup> Tb	T: symmetrized from 940(+330-220) $\mu$ s							
<sup>136</sup> In	-40510# 400#		86 ms 9			15 15Lo04 TD	2015	$\beta^- =100; \beta^-n=0\#; \beta^-2n=90\#$
<sup>136</sup> Sn	-55900# 300#		350 ms 5	0+		14 15Lo04 T	1994	$\beta^- =100; \beta^-n=28 3; \beta^-2n=2\#$
<sup>136</sup> Sb	-64507 6		923 ms 14	(1 <sup>-</sup> )		02 15Lo08 J	1976	$\beta^- =100; \beta^-n=16.3 32; \beta^-2n=10\#$
<sup>136</sup> Sb <sup>m</sup>	-64230 6	277.0 0.7	570 ns 5	(6 <sup>-</sup> )		02 12Ka36 ET	2001	IT=100
<sup>136</sup> Te	-74425.3 2.3		17.63 s 0.08	0+		02	1974	$\beta^- =100; \beta^-n=1.31 5$
<sup>136</sup> I	-79545 14		83.4 s 1.0	(1 <sup>-</sup> )		02	1949	$\beta^- =100$
<sup>136</sup> I <sup>m</sup>	-79339 5	206 15 BD	46.9 s 1.0	(6 <sup>-</sup> )		02	1959	$\beta^- =100; IT=0$
<sup>136</sup> Xe	-86429.159 0.007		2.19 Zy 0.06	0+		02 15Ba11 T	1920	IS=8.8573 44; $2\beta^- =100$
<sup>136</sup> Xe <sup>m</sup>	-84537.456 0.016	1891.703 0.014	2.95 $\mu$ s 0.09	6 <sup>+</sup>		02	1969	IT=100
<sup>136</sup> Cs	-86338.7 1.9		13.16 d 0.03	5 <sup>+</sup>		02	1951	$\beta^- =100$
<sup>136</sup> Cs <sup>m</sup>	-85820.8 1.9	517.9 0.1	17.5 s 0.2	8 <sup>-</sup>		02 11Wi09 ET	1981	IT=?; $\beta^- ?$
<sup>136</sup> Ba	-88886.9 0.3		STABLE	0+		02	1932	IS=7.854 24
<sup>136</sup> Ba <sup>m</sup>	-86856.4 0.3	2030.466 0.018	308.4 ms 1.9	7 <sup>-</sup>		02	1965	IT=100
<sup>136</sup> Ba <sup>n</sup>	-85529.5 0.5	3357.4 0.4	91 ns 2	(10 <sup>+</sup> )		04Va03 TD	2004	IT=100
<sup>136</sup> La	-86040 50		9.87 m 0.03	1 <sup>+</sup>		02	1950	$\beta^+ =100$
<sup>136</sup> La <sup>m</sup>	-85780 50	259.3 0.4	114 ms 3	(7) <sup>(-#)</sup>		02 05Bh06 EJ	1966	IT=100
<sup>136</sup> Ce	-86508.4 0.4		STABLE (>38 Py)	0+		02 01Da22 T	1936	IS=0.185 2; $2\beta^+ ?$
<sup>136</sup> Ce <sup>m</sup>	-83412.9 0.6	3095.5 0.4	1.96 $\mu$ s 0.09	10 <sup>+</sup>		02 13Va10 T	1991	IT=100
<sup>136</sup> Pr	-81340 11		13.1 m 0.1	2 <sup>+</sup>		02	1968	$\beta^+ =100$
<sup>136</sup> Nd	-79199 12		50.7 m 0.3	0+		02	1968	$\beta^+ =100$
<sup>136</sup> Pm	-71170 70		107 s 6	(5) <sup>(+#)</sup>		02 FGK12a J	1982	$\beta^+ =100$
<sup>136</sup> Pm <sup>m</sup>	-71070 90	100 120 MD * &	300 s 50	(2) <sup>(+#)</sup>		02 88Ke03 T	1988	$\beta^+ =100$
<sup>136</sup> Pm <sup>n</sup>	-71100 70	68 25	1.5 $\mu$ s 0.1	8 <sup>+#</sup>		02 08Ri05 ET	1987	IT=100
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)					
... A-group continued ...													
<sup>136</sup> Sm	-66811	12	47 s	2	0 <sup>+</sup>	02	1982	$\beta^+=100$					
<sup>136</sup> Sm <sup>m</sup>	-64546	12	2264.7	1.1	15 $\mu$ s	1	1994	IT=100					
<sup>136</sup> Eu	-56240#	200#			3.3 s	0.3	02	89Vi04 D 1987	$\beta^+=100; \beta^+p=0.09$ 3				
<sup>136</sup> Eu <sup>m</sup>	-56240#	540#	0#	500#	*	3.8 s	0.3	(3 <sup>+</sup> ) 02 89Vi04 D 1987	$\beta^+=100; \beta^+p=0.09$ 3				
<sup>136</sup> Gd	-49090#	300#			1# s	(>200 ns)	0 <sup>+</sup>	02 00So11 I 2000	$\beta^+ ?; \beta^+p ?$				
<sup>136</sup> Tb	-36130#	500#			200# ms			02	$\beta^+ ?; \beta^+p ?$				
* <sup>136</sup> In	T: symmetrized from 15Lo04=85(+10-8)								**				
* <sup>136</sup> Sn	D: $\beta^-n$ average 11Ar18=27(4)% 02Sh08=30(5)%								**				
* <sup>136</sup> Sb <sup>m</sup>	E: 12Ka36=53.9(0.5) and 173.1(0.5) $\gamma$ in cascade to ground-state; 15Lo08=269.3								**				
* <sup>136</sup> Sb <sup>m</sup>	T: others 15Lo08=489(40) 07Si27=480(100) 01Mi22=570(50)								**				
* <sup>136</sup> Xe	T: others 14Al03=2.165(0.061)Zy 12Ga17=2.38(0.14)Zy 02Be74>10Zy								**				
* <sup>136</sup> Xe	T: 0v- $\beta\beta$ : 16As01>2500 Zy 13Ga07>19000 Zy 12Au03>16000 Zy (all at 90% C.L.)								**				
* <sup>136</sup> Cs <sup>m</sup>	E: also 83We07=518(5)								**				
* <sup>136</sup> Ba <sup>n</sup>	T: other 04Sh15=94(10) outweighed								**				
* <sup>136</sup> Ce	T: also 11Be02>18Py; both for 2v- $\beta\beta$ and 1 $\sigma$								**				
* <sup>136</sup> Ce <sup>m</sup>	T: average 13Va10=1.9(0.1) 75Yo01=2.2(0.2)								**				
* <sup>136</sup> Pm	J: expected 5 <sup>+</sup> n9/2[514]+p1/2[550]; supported by observed direct feeding								**				
* <sup>136</sup> Pm	J: to I=4,5,6 levels following <sup>136</sup> Pm $\beta^+$ decay								**				
* <sup>136</sup> Pm <sup>m</sup>	J: expected 2 <sup>+</sup> n9/2[514]+p1/2[550]; supported by observed direct feeding								**				
* <sup>136</sup> Pm <sup>m</sup>	J: to I=2 <sup>+</sup> and 3 <sup>+</sup> levels following $\beta^+$ decay								**				
* <sup>136</sup> Pm <sup>n</sup>	E: 08Ri05=42.7(0.2) keV above a long lived state that could be either the								**				
* <sup>136</sup> Pm <sup>n</sup>	E: ground-state or an excited level located <50 keV above the ground-state owing to non-								**				
* <sup>136</sup> Pm <sup>n</sup>	E: observation of any decay radiation								**				
<sup>137</sup> In	-35040#	500#	70 ms	40	9/2 <sup>+</sup> #	15	15Lo04	TD 2015	$\beta^-=100; \beta^-n=0\#; \beta^-2n=90\#$	*			
<sup>137</sup> Sn	-49790#	400#	273 ms	7	5/2 <sup>-</sup> #	07	11Ar18	TD 1994	$\beta^-=100; \beta^-n=50$ 8; $\beta^-2n=40\#$	*			
<sup>137</sup> Sb	-60060	50	484 ms	22	7/2 <sup>+</sup> #	07	11Ar18	TD 1994	$\beta^-=100; \beta^-n=49$ 6; $\beta^-2n=0.3\#$	*			
<sup>137</sup> Te	-69303.8	2.1	2.49 s	0.05	3/2 <sup>-</sup> #	07		1975	$\beta^-=100; \beta^-n=2.99$ 16	*			
<sup>137</sup> I	-76356	8	24.13 s	0.12	7/2 <sup>+</sup> #	07	16Ag03	D 1943	$\beta^-=100; \beta^-n=7.76$ 14	*			
<sup>137</sup> Xe	-82383.40	0.10	3.818 m	0.013	7/2 <sup>-</sup>	07		1943	$\beta^-=100$	*			
<sup>137</sup> Cs	-86545.6	0.4	30.08 y	0.09	7/2 <sup>+</sup>	07		1951	$\beta^-=100$	*			
<sup>137</sup> Ba	-87721.2	0.3	STABLE		3/2 <sup>+</sup>	07		1932	IS=11.232 24	*			
<sup>137</sup> Ba <sup>m</sup>	-87059.5	0.3	661.659	0.003	2.552 m	0.001		11/2 <sup>-</sup>	07	1965	IT=100		
<sup>137</sup> Ba <sup>n</sup>	-85372.1	0.6	2349.1	0.5	590 ns	100		(17/2 <sup>-</sup> )	07	1973	IT=100		
<sup>137</sup> La	-87140.7	1.7	60 ky	20	7/2 <sup>+</sup>	07		1948	$\epsilon=100$	*			
<sup>137</sup> La <sup>m</sup>	-85271.2	1.7	1869.50	0.21	342 ns	25		19/2 <sup>-</sup>	07	1982	IT=100		
<sup>137</sup> Ce	-85918.6	0.4	9.0 h	0.3	3/2 <sup>+</sup>	07		1948	$\beta^+=100$	*			
<sup>137</sup> Ce <sup>m</sup>	-85664.3	0.4	254.29	0.05	34.4 h	0.3		11/2 <sup>-</sup>	07	1958	IT=99.21 4; $\beta^+=0.79$ 4		
<sup>137</sup> Pr	-83202	8	1.28 h	0.03	5/2 <sup>+</sup>	07		1958	$\beta^+=100$	*			
<sup>137</sup> Pr <sup>m</sup>	-82641	8	561.22	0.23	2.66 $\mu$ s	0.07		11/2 <sup>-</sup>	07	1987	IT=100		
<sup>137</sup> Nd	-79585	12	38.5 m	1.5	1/2 <sup>+</sup>	07		1970	$\beta^+=100$	*			
<sup>137</sup> Nd <sup>m</sup>	-79066	12	519.43	0.20	1.60 s	0.15		11/2 <sup>-</sup>	07	1970	IT=100		
<sup>137</sup> Pm	-74073	13			2# m			5/2 <sup>+</sup> #		1975	$\beta^+ ?$		
<sup>137</sup> Pm <sup>m</sup>	-73930	50	150	50	BD	&		2.4 m	0.1	11/2 <sup>-</sup>	07	1973	$\beta^+=100$
<sup>137</sup> Sm	-68030	40			45 s	1		(9/2 <sup>-</sup> )	07	1986	$\beta^+=100$		
<sup>137</sup> Sm <sup>m</sup>	-67850#	60#	180#	50#	20# s			1/2 <sup>+</sup> #			$\beta^+ ?$		
<sup>137</sup> Eu	-60146	4			8.4 s	0.5		11/2 <sup>-</sup> #	07	88Be.A T 1982	$\beta^+=100$		
<sup>137</sup> Gd	-51210#	300#			2.2 s	0.2		(7/2) <sup>(+)</sup> #	07	1999	$\beta^+=100; \beta^+p=?$		
<sup>137</sup> Tb	-40970#	400#			600# ms			11/2 <sup>-</sup> #			p ?; $\beta^+ ?$		
* <sup>137</sup> In	T: symmetrized from 15Lo04=65(+40-30)								**				
* <sup>137</sup> Sb	T: average 11Ar18=492(25) 02Sh08=450(50)								**				
* <sup>137</sup> Sb	D: $\beta^-n$ average 11Ar18=49(8)% 02Sh08=49(10)%								**				
* <sup>137</sup> Te	J: TNN of N=85 isotones. ENSDF'07 gives (7/2 <sup>-</sup> ) from shell-model prediction								**				
* <sup>137</sup> Te	D: from 93Ru01 evaluation								**				
* <sup>137</sup> I	T: 93Ru01=24.13(0.12) supersedes 74Ru08=24.5(0.2) from same group								**				
<sup>138</sup> Sn	-44860#	500#	150 ms	30	0 <sup>+</sup>	16		2010	$\beta^-=100; \beta^-n\approx 36; \beta^-2n=5\#$	*			
<sup>138</sup> Sn <sup>m</sup>	-43520#	500#	1344	2	210 ns	45		(6 <sup>+</sup> )	16	2014	IT=100		
<sup>138</sup> Sb	-54220	1060	348 ms	15	(0 <sup>-</sup> )	16	FGK16a	J 1994	$\beta^-=100; \beta^-n=72$ 8; $\beta^-2n=2\#$	*			
<sup>138</sup> Te	-65696	4	1.4 s	0.4	0 <sup>+</sup>	03		1975	$\beta^-=100; \beta^-n=6.3$ 21	*			
<sup>138</sup> I	-71980	6	6.23 s	0.03	(1 <sup>-</sup> )	16	93Ru01	D 1949	$\beta^-=100; \beta^-n=5.46$ 18	*			
<sup>138</sup> I <sup>m</sup>	-71912	6	67.9	0.5	1.26 $\mu$ s	0.16		(3 <sup>-</sup> )	16	2007	IT=100		
<sup>138</sup> Xe	-79972.2	2.8	14.14 m	0.05	0 <sup>+</sup>	03	12Wa21	T 1943	$\beta^-=100$	*			
<sup>138</sup> Cs	-82887	9	33.41 m	0.18	3 <sup>-</sup>	03		1943	$\beta^-=100$	*			
<sup>138</sup> Cs <sup>m</sup>	-82807	9	79.9	0.3	2.91 m	0.08		6 <sup>-</sup>	03	1971	IT=81 2; $\beta^-=19$ 2		
<sup>138</sup> Cs <sup>x</sup>	-82847	25	40	23	R=?			fsmix					
<sup>138</sup> Ba	-88261.6	0.3	STABLE		0 <sup>+</sup>	03		1925	IS=71.698 42	*			
<sup>138</sup> Ba <sup>m</sup>	-86171.1	0.3	2090.54	0.06	800 ns	100		6 <sup>+</sup>	03	1971	IT=100		
... A-group is continued on next page ...													

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...												
<sup>138</sup> La	-86519	3			102 Gy	1	5 <sup>+</sup>	03	1947	IS=0.08881 71; $\beta^+$ =65.6 5; $\beta^-$ =34.4 5		
<sup>138</sup> La <sup>m</sup>	-86446	3	72.57	0.03	116 ns	5	(3) <sup>+</sup>	03	1975	IT=100		
<sup>138</sup> La <sup>n</sup>	-85780	3	739.2	0.5	2.0 $\mu$ s	0.3	7 <sup>-</sup>		14As02 ETJ	2014	IT=100	
<sup>138</sup> Ce	-87571	5			STABLE	(>57 Py)	0 <sup>+</sup>	03	11Be02 T	1936	IS=0.251 2; 2 $\beta^+$ ?	
<sup>138</sup> Ce <sup>m</sup>	-85442	5	2129.17	0.12	8.65 ms	0.20	7 <sup>-</sup>	03		1960	IT=100	
<sup>138</sup> Pr	-83134	11			1.45 m	0.05	1 <sup>+</sup>	03		1951	$\beta^+$ =100	
<sup>138</sup> Pr <sup>m</sup>	-82782	17	352	19	BD		2.12 h	0.04	7 <sup>-</sup>	03	1958	$\beta^+$ =100
<sup>138</sup> Nd	-82018	12			5.04 h	0.09	0 <sup>+</sup>	03		1965	$\beta^+$ =100	
<sup>138</sup> Nd <sup>m</sup>	-78843	12	3174.9	0.4	370 ns	5	10 <sup>+</sup>	03	13Va10 T	1975	IT=100	
<sup>138</sup> Pm	-74940	28			10 s	2	1 <sup>+</sup> #	03		1981	$\beta^+$ =100	
<sup>138</sup> Pm <sup>m</sup>	-74911	13	30	30	BD	*	3.24 m	0.05	5 <sup>-</sup> #	03	1973	$\beta^+$ =100
<sup>138</sup> Pm <sup>n</sup>			non existent		EU	*	3.24 m	0.05	(3 <sup>+</sup> )			$\beta^+$ =100
<sup>138</sup> Sm	-71498	12			3.1 m	0.2	0 <sup>+</sup>	03	81De38 I	1982	$\beta^+$ =100	
<sup>138</sup> Eu	-61750	28			12.1 s	0.6	(6 <sup>-</sup> )	03		1982	$\beta^+$ =100	
<sup>138</sup> Gd	-55800#	200#			4.7 s	0.9	0 <sup>+</sup>	03		1985	$\beta^+$ =100	
<sup>138</sup> Gd <sup>m</sup>	-53570#	200#	2233.1	0.5	6.2 $\mu$ s	0.2	(8 <sup>-</sup> )	03	11Pr02 T	1997	IT=100	
<sup>138</sup> Tb	-43670#	300#			800# ms	(>200 ns)		03	00So11 I	1993	$\beta^+$ ?; $\beta^+$ p ?; p=0	
<sup>138</sup> Dy	-34930#	500#			200# ms		0 <sup>+</sup>				$\beta^+$ ?; $\beta^+$ p ?	
* <sup>138</sup> Sn	T : symmetrized from 15Lo04=140(+30-20)											
* <sup>138</sup> Sb	J : expected pg7/2 nf7/2 config and strong repulsive residual interaction											
* <sup>138</sup> I <sup>m</sup>	J : 67.9 E2 $\gamma$ ray (delayed) to (1 <sup>-</sup> )											
* <sup>138</sup> Xe	T : average of 12Wa21=14.18(0.10) 72Mo33=14.08(0.08) 69Ca03=14.17(0.07)											
* <sup>138</sup> Ce	T : also 01Da22>150Ty; both for 2v- $\beta\beta$ and 1 $\sigma$											
* <sup>138</sup> Pm <sup>n</sup>	D : arguments for a second isomer of intermediate spin are not convincing											
* <sup>138</sup> Gd <sup>m</sup>	E : for least-squares fit to $\gamma$ -ray energies in 11Pr02											
* <sup>138</sup> Tb	D : from 93Li40											
<sup>139</sup> Sn	-38440#	500#			130 ms	60	5/2 <sup>-</sup> #	15		2015	$\beta^-$ =100; $\beta^-$ n=80#; $\beta^-$ 2n=20#	
<sup>139</sup> Sb	-49790#	400#			93 ms	13	7/2 <sup>+</sup> #	01	11Ar18 TD	1994	$\beta^-$ =100; $\beta^-$ n=90 10; $\beta^-$ 2n=3#	
<sup>139</sup> Te	-60205	4			500# ms	(>150 ns)	5/2 <sup>-</sup> #	01	94Be24 I	1994	$\beta^-$ ?; $\beta^-$ n=2#	
<sup>139</sup> I	-68471	4			2.282 s	0.010	7/2 <sup>+</sup> #	01	93Ru01 T	1949	$\beta^-$ =100; $\beta^-$ n=10.0 3	
<sup>139</sup> Xe	-75644.6	2.1			39.68 s	0.14	3/2 <sup>-</sup>	01		1951	$\beta^-$ =100	
<sup>139</sup> Cs	-80701	3			9.27 m	0.05	7/2 <sup>+</sup>	01		1939	$\beta^-$ =100	
<sup>139</sup> Ba	-84913.8	0.3			83.13 m	0.06	(7/2 <sup>-</sup> )	01	12Da17 T	1937	$\beta^-$ =100	
<sup>139</sup> La	-87226.2	2.0			STABLE		7/2 <sup>+</sup>	01		1924	IS=99.91119 71	
<sup>139</sup> La <sup>m</sup>	-85426.3	2.1	1799.9	0.5	315 ns	35	(17/2 <sup>+</sup> )		12As06 ETJ	2012	IT=100	
<sup>139</sup> Ce	-86948	7			137.641 d	0.020	3/2 <sup>+</sup>	01		1948	$\epsilon$ =100	
<sup>139</sup> Ce <sup>m</sup>	-86194	7	754.24	0.08	56.54 s	0.13	11/2 <sup>-</sup>	01	94It.A T	1967	IT=100	
<sup>139</sup> Pr	-84819	8			4.41 h	0.04	5/2 <sup>+</sup>	01		1951	$\beta^+$ =100	
<sup>139</sup> Nd	-82014	28			29.7 m	0.5	3/2 <sup>+</sup>	01		1951	$\beta^+$ =100	
<sup>139</sup> Nd <sup>m</sup>	-81783	28	231.15	0.05	5.50 h	0.20	11/2 <sup>-</sup>	01		1951	$\beta^+$ =88.2 4; IT=11.8 4	
<sup>139</sup> Nd <sup>n</sup>	-79398	28	2616	2	276.8 ns	1.8	23/2 <sup>+</sup>	01	13Va10 ETJ	1980	IT ?	
<sup>139</sup> Pm	-77500	14			4.15 m	0.05	(5/2 <sup>+</sup> )	01		1967	$\beta^+$ =100	
<sup>139</sup> Pm <sup>m</sup>	-77311	14	188.7	0.3	180 ms	20	(11/2 <sup>-</sup> )	01		1975	IT $\approx$ 100; $\beta^+$ =0.16#	
<sup>139</sup> Sm	-72380	11			2.57 m	0.10	1/2 <sup>+</sup>	01		1971	$\beta^+$ =100	
<sup>139</sup> Sm <sup>m</sup>	-71923	11	457.40	0.22	10.7 s	0.6	11/2 <sup>-</sup>	01		1973	IT=93.7 5; $\beta^+$ =6.3 5	
<sup>139</sup> Eu	-65398	13			17.9 s	0.6	(11/2 <sup>-</sup> )	01		1975	$\beta^+$ =100	
<sup>139</sup> Eu <sup>m</sup>	-65250	13	148.2	0.2	10 $\mu$ s	2	(7/2 <sup>+</sup> )		11Cu01 ETJ	2011	IT=100	
<sup>139</sup> Gd	-57630#	200#			5.7 s	0.3	9/2 <sup>-</sup> #	01	99Xi04 T	1983	$\beta^+$ =100; $\beta^+$ p=?	
<sup>139</sup> Gd <sup>m</sup>	-57380#	250#	150#		4.8 s	0.9	1/2 <sup>+</sup> #	01		1983	$\beta^+$ =100; $\beta^+$ p=?	
<sup>139</sup> Tb	-48130#	300#			1.6 s	0.2	11/2 <sup>-</sup> #	01		1999	$\beta^+$ =100; $\beta^+$ p ?	
<sup>139</sup> Dy	-37640#	500#			600 ms	200	(7/2 <sup>+</sup> )	01		1999	$\beta^+$ =100; $\beta^+$ p ?	
* <sup>139</sup> I	T : average 93Ru01=2.280(0.011) 80Al15=2.29(0.02)											
* <sup>139</sup> Ba	T : average 12Da17=83.01(0.14) 12Da04=83.25(0.08) 72Em01=82.71(0.18)											
* <sup>139</sup> Nd <sup>n</sup>	T : average 13Va10=278(2) 08Fe02=272(4)											
* <sup>139</sup> Nd <sup>n</sup>	T : 80Mu10 > 141 ns											
* <sup>139</sup> Gd	T : average 99Xi04=5.8(0.9) 88Be.A=5.8(0.4); other 83Ni05=4.9(1.0) not used											
* <sup>139</sup> Gd	T : since it corresponds to a mixture of ground-state and isomer											
* <sup>139</sup> Gd <sup>m</sup>	D : assuming that the delayed protons reported in 83Ni05 are from both states											
<sup>140</sup> Sb	-43940#	600#			100# ms	(>400 ns)	(4 <sup>-</sup> , 3 <sup>-</sup> )	16		2010	$\beta^-$ ?; $\beta^-$ n=40#; $\beta^-$ 2n=20#	
<sup>140</sup> Sb <sup>m</sup>	-43610#	600#	330	10	41 $\mu$ s	8	(6 <sup>-</sup> , 7 <sup>-</sup> )	16		2016	IT=100	
<sup>140</sup> Te	-56580	60			300# ms	(>300 ns)	0 <sup>+</sup>	07		1994	$\beta^-$ ?; $\beta^-$ n=3#	
<sup>140</sup> I	-63606	12			860 ms	40	(4 <sup>-</sup> )	07		1972	$\beta^-$ =100; $\beta^-$ n=9.3 10; $\beta^-$ 2n=0#	
<sup>140</sup> Xe	-72986.5	2.3			13.60 s	0.10	0 <sup>+</sup>	07		1951	$\beta^-$ =100	
<sup>140</sup> Cs	-77050	8			63.7 s	0.3	1 <sup>-</sup>	07		1950	$\beta^-$ =100	
<sup>140</sup> Cs <sup>m</sup>	-77036	8	13.931	0.021	471 ns	51	(2) <sup>-</sup>	07		1974	IT=100	
... A-group is continued on next page ...												

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>140</sup> Ba	-83269	8	12.7527 d	0.0023	0 <sup>+</sup>	07	1939	$\beta^-$ =100
<sup>140</sup> La	-84315.9	2.0	40.285 h	0.003	3 <sup>-</sup>	07	1935	$\beta^-$ =100
<sup>140</sup> Ce	-88076.1	1.6	STABLE		0 <sup>+</sup>	07	1925	IS=88.450 51
<sup>140</sup> Ce <sup>m</sup>	-85968.2	1.6	2107.86	0.03	7.3 $\mu$ s	1.5	1966	IT=100
<sup>140</sup> Pr	-84688	6	3.39 m	0.01	1 <sup>+</sup>	07	1938	$e^+$ =51.3 18; $\epsilon$ =48.7 18
<sup>140</sup> Pr <sup>m</sup>	-84560	6	127.8	0.3	350 ns	20	1964	IT=100
<sup>140</sup> Pr <sup>n</sup>	-83924	6	763.7	0.5	3.05 $\mu$ s	0.20	1964	IT=100
<sup>140</sup> Nd	-84259	3	3.37 d	0.02	0 <sup>+</sup>	07	1949	$\epsilon$ =100
<sup>140</sup> Nd <sup>m</sup>	-82038	3	2221.4	0.1	600 $\mu$ s	50	1962	IT=100
<sup>140</sup> Nd <sup>n</sup>	-76829	3	7429.6	0.7	1.22 $\mu$ s	0.06	2008	IT=100
<sup>140</sup> Pm	-78214	24	9.2 s	0.2	1 <sup>+</sup>	07	1966	$\beta^+$ =100
<sup>140</sup> Pm <sup>m</sup>	-77783	13	431	28	5.95 m	0.05	1966	$\beta^+$ =100
<sup>140</sup> Sm	-75456	12	14.82 m	0.12	0 <sup>+</sup>	07	1967	$\beta^+$ =100
<sup>140</sup> Eu	-66990	50	1.51 s	0.02	1 <sup>+</sup>	07	1982	$\beta^+$ =100
<sup>140</sup> Eu <sup>m</sup>	-66780	50	210	15	125 ms	2	1988	IT $\approx$ 100; $\beta^+$ <1
<sup>140</sup> Eu <sup>n</sup>	-66320	50	669	15	299.8 ns	2.1	2002	IT=100
<sup>140</sup> Gd	-61782	28	15.8 s	0.4	0 <sup>+</sup>	07	1985	$\beta^+$ =100
<sup>140</sup> Tb	-50480	800	2.32 s	0.16	(7 <sup>+</sup> )	07	1986	$\beta^+$ =100; $\beta^+$ p=0.26 13
<sup>140</sup> Dy	-42830#	400#	700# ms		0 <sup>+</sup>	07	2002	$\beta^+$ ?; $\beta^+$ p?
<sup>140</sup> Dy <sup>m</sup>	-40660#	400#	2166.1	0.5	7.0 $\mu$ s	0.5	2002	IT=100
<sup>140</sup> Ho	-29260#	500#	6 ms	3	8 <sup>+</sup> #	07	1999	p=?; $\beta^+$ =1#; $\beta^+$ p?
* <sup>140</sup> Sb <sup>m</sup>	E : 16Lo01=298.2 + x, x estimated by authors x=30#							
* <sup>140</sup> Pr	T : other: 07Li71=7.3(0.4) for q=59 <sup>+</sup> (bare ion) 3.04(0.10) for q=58 <sup>+</sup>							
* <sup>140</sup> Pr	T : (H-like ion) and 3.84(0.15) for q=57 <sup>+</sup> (He-like ion)							
* <sup>140</sup> Pr	D : e <sup>+</sup> =42.4(2.3)%; $\epsilon$ =57.6(2.3)% for q=58 <sup>+</sup> (H-like ion) and							
* <sup>140</sup> Pr	D : e <sup>+</sup> =51.2(3.1)%; $\epsilon$ =48.8(3.1)% for q=57 <sup>+</sup> (He-like ion)							
* <sup>140</sup> Nd <sup>n</sup>	E : uncertainty not given, estimated by evaluator							
* <sup>140</sup> Nd <sup>n</sup>	T : average 13Va10=1.2(0.1) 08Fe02=1.23(0.07)							
* <sup>140</sup> Eu <sup>m</sup>	E : less than 50 keV above 185.3 level, from ENSDF, thus 185.3 + 25(15)							
* <sup>140</sup> Eu <sup>n</sup>	E : 459.5(0.3) keV above <sup>140</sup> Eu <sup>m</sup>							
* <sup>140</sup> Tb	T : average 06Xu03=2.0(0.5) 00Xu08=2.1(0.4) 91Fi03=2.4(0.2) 86Wi15=2.4(0.4)							
* <sup>140</sup> Ho	D : from estimated $\beta^+$ half-life 400# ms; p observed in 99Ry04							
<sup>141</sup> Sb	-39110#	500#	100# ms		7/2 <sup>+</sup> #			$\beta^-$ ?; $\beta^-$ n=90#; $\beta^-$ 2n=3#
<sup>141</sup> Te	-50490#	400#	150# ms	(>150 ns)	5/2 <sup>-</sup> #	14	94Be24 I	1994
<sup>141</sup> I	-59927	16	430 ms	20	7/2 <sup>+</sup> #	14		1974
<sup>141</sup> Xe	-68197.3	2.9	1.73 s	0.01	5/2 <sup>(-#)</sup>	14		1951
<sup>141</sup> Cs	-74478	9	24.84 s	0.16	7/2 <sup>+</sup>	14		1962
<sup>141</sup> Ba	-79733	5	18.27 m	0.07	3/2 <sup>-</sup>	14		1945
<sup>141</sup> La	-82932	4	3.92 h	0.03	(7/2 <sup>+</sup> )	14		1951
<sup>141</sup> Ce	-85432.9	1.6	32.511 d	0.013	7/2 <sup>-</sup>	14		1948
<sup>141</sup> Pr	-86015.6	1.7	STABLE		5/2 <sup>+</sup>	14		1924
<sup>141</sup> Nd	-84193	3	2.49 h	0.03	3/2 <sup>+</sup>	14		1949
<sup>141</sup> Nd <sup>m</sup>	-83436	3	756.51	0.05	62.0 s	0.8	70Ab05 D	1960
<sup>141</sup> Pm	-80523	14	20.90 m	0.05	5/2 <sup>+</sup>	14		1952
<sup>141</sup> Pm <sup>m</sup>	-79894	14	628.62	0.07	630 ns	20		1970
<sup>141</sup> Pm <sup>n</sup>	-77992	14	2530.75	0.17	> 2 $\mu$ s			1985
<sup>141</sup> Sm	-75934	9	10.2 m	0.2	1/2 <sup>+</sup>	14		1967
<sup>141</sup> Sm <sup>m</sup>	-75758	9	175.9	0.3	22.6 m	0.2		1967
<sup>141</sup> Eu	-69926	13	40.7 s	0.7	5/2 <sup>+</sup>	14		1977
<sup>141</sup> Eu <sup>m</sup>	-69830	13	96.45	0.07	2.7 s	0.3		1973
<sup>141</sup> Gd	-63224	20	14 s	4	(1/2 <sup>+</sup> )	14		1986
<sup>141</sup> Gd <sup>m</sup>	-62846	20	377.76	0.09	24.5 s	0.5		1986
<sup>141</sup> Tb	-54540	110	3.5 s	0.2	(5/2 <sup>-</sup> )	14		1986
<sup>141</sup> Tb <sup>m</sup>	-54540#	230#	0#	200#	7.9 s	0.6	88Be.A I	1988
<sup>141</sup> Dy	-45380#	300#	900 ms	140	11/2 <sup>-</sup> #	14		1984
<sup>141</sup> Ho	-34360#	400#	4.1 ms	0.1	(7/2 <sup>-</sup> )	14		1998
<sup>141</sup> Ho <sup>m</sup>	-34290#	400#	66	2	7.3 $\mu$ s	0.3		1998
* <sup>141</sup> I	D : rounded from 21.2(3.0); 80Al15=21.2(3.0) included in 93Ru01=22(3)							
* <sup>141</sup> Eu <sup>m</sup>	D : symmetrized from IT=87(+2-4)% and $\beta^+$ =13(+4-2)%							
* <sup>141</sup> Gd	J : weak arguments in ENSDF'2001 for $J^\pi$ assignment; same for <sup>141</sup> Gd <sup>m</sup>							
* <sup>141</sup> Tb <sup>m</sup>	I : existence discussed in 88Be.A. Provisionally accepted							
* <sup>141</sup> Ho	D : from estimated $\beta^+$ half-life 200# ms							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>142</sup> Te	-46370# 500#		100# ms (>150 ns)	0 <sup>+</sup>	11	94Be24 I	1994	$\beta^-$ ?; $\beta^-n=10\#$ ; $\beta^-2n=0\#$
<sup>142</sup> I	-54770 370		222 ms 12	2 <sup>-</sup> #	11		1975	$\beta^-n=100$ ; $\beta^-n=20\#$ ; $\beta^-2n=1\#$
<sup>142</sup> Xe	-65229.6 2.7		1.23 s 0.02	0 <sup>+</sup>	11		1960	$\beta^-n=100$ ; $\beta^-n=0.21$ 6 *
<sup>142</sup> Cs	-70515 7		1.684 s 0.014	0 <sup>-</sup>	11		1962	$\beta^-n=100$ ; $\beta^-n=0.090$ 4
<sup>142</sup> Ba	-77842 6		10.6 m 0.2	0 <sup>+</sup>	11		1959	$\beta^-n=100$ *
<sup>142</sup> La	-80024 6		91.1 m 0.5	2 <sup>-</sup>	11		1953	$\beta^-n=100$
<sup>142</sup> La <sup>m</sup>	-79878 6	145.82 0.08	870 ns 170	(4) <sup>-</sup>	11		1983	IT=100
<sup>142</sup> Ce	-84533.2 2.5		STABLE (>50 Py)	0 <sup>+</sup>	11		1925	IS=11.114 51; $\alpha$ ?; $2\beta^-$ ? *
<sup>142</sup> Pr	-83787.5 1.7		19.12 h 0.04	2 <sup>-</sup>	11		1935	$\beta^-n\approx 100$ ; $\epsilon=0.0164$ 8
<sup>142</sup> Pr <sup>m</sup>	-83783.8 1.7	3.694 0.003	14.6 m 0.5	5 <sup>-</sup>	11		1967	IT=100
<sup>142</sup> Nd	-85950.0 1.4		STABLE	0 <sup>+</sup>	11		1924	IS=27.152 40
<sup>142</sup> Nd <sup>m</sup>	-83740.7 1.4	2209.303 0.021	16.5 $\mu$ s	6 <sup>+</sup>	14		1964	IT=100
<sup>142</sup> Pm	-81142 24		40.5 s 0.5	1 <sup>+</sup>	11		1959	$e^+=77.1$ 27; $\epsilon=22.9$ 27 *
<sup>142</sup> Pm <sup>m</sup>	-80259 24	883.17 0.16	2.0 ms 0.2	(8) <sup>-</sup>	11		1971	IT=100
<sup>142</sup> Pm <sup>n</sup>	-78313 24	2828.7 0.6	67 $\mu$ s 5	(13) <sup>-</sup>	11		1974	IT=100
<sup>142</sup> Sm	-78986 3		72.49 m 0.05	0 <sup>+</sup>	11		1959	$\beta^+n=100$
<sup>142</sup> Sm <sup>m</sup>	-76614 3	2372.1 0.4	170 ns 2	7 <sup>-</sup>	11		1975	IT=100
<sup>142</sup> Sm <sup>n</sup>	-75324 3	3662.2 0.7	480 ns 60	10 <sup>+</sup>	11		1979	IT=100
<sup>142</sup> Eu	-71310 30		2.36 s 0.10	1 <sup>+</sup>	11	91Fi03 T	1966	$\beta^+n=100$ *
<sup>142</sup> Eu <sup>m</sup>	-70856 12	460 30 BD	1.223 m 0.008	8 <sup>-</sup>	11		1966	$\beta^+n=100$
<sup>142</sup> Gd	-66960 28		70.2 s 0.6	0 <sup>+</sup>	11		1986	$\epsilon=52$ 5; $e^+=48$ 5
<sup>142</sup> Tb	-56560 700		597 ms 17	1 <sup>+</sup>	11		1991	$\beta^+n=100$ ; $\beta^+p=0.0022$ 11
<sup>142</sup> Tb <sup>m</sup>	-56280 700	279.7 0.4	303 ms 17	5 <sup>-</sup>	11		1986	IT=100
<sup>142</sup> Tb <sup>n</sup>	-55910 700	652.1 0.6	26 $\mu$ s 1	8 <sup>+</sup>	11		1989	IT=100
<sup>142</sup> Dy	-50120# 730#		2.3 s 0.3	0 <sup>+</sup>	11		1986	$\beta^+n=100$ ; $\beta^+p=0.06$ 3
<sup>142</sup> Ho	-37250# 400#		400 ms 100	(7 <sup>-</sup> , 8 <sup>+</sup> )	11		2001	$\beta^+n\approx 100$ ; $\beta^+p=?$ ; $p\approx 0$ *
<sup>142</sup> Er	-28030# 500#		10# $\mu$ s	0 <sup>+</sup>				$p?$
* <sup>142</sup> Xe	D: 03Be05=0.21(6) 75As04=0.406(0.034) T: 03Be05=1.250(0.025) **							
* <sup>142</sup> Ba	D: $\beta^-n=0.091(0.003)\%$ in ENSDF'00 contradicts $Q(\beta^-n)=-2979(7)$ keV **							
* <sup>142</sup> Ce	T: lower limit is for $\alpha$ decay; for $\beta\beta$ decay 11Be02>300Py 01Da22>260 Py **							
* <sup>142</sup> Pm	T: other: 09Wi09=56(3) for $q=61^+$ (bare ion) 39.2(0.7) for $q=60^+$ **							
* <sup>142</sup> Pm	T: (H-like ion) and 39.6(1.4) for $q=59^+$ (He-like ion) **							
* <sup>142</sup> Pm	D: $e^+=71.0(1.3)\%$ ; $\epsilon=29.0(1.3)\%$ for $q=60^+$ (H-like ion) and **							
* <sup>142</sup> Pm	D: $e^+=79.8(1.0)\%$ ; $\epsilon=20.2(1.0)\%$ for $q=59^+$ (He-like ion) **							
* <sup>142</sup> Eu	T: average 91Fi03=2.34(0.12) 75Ke08=2.4(0.2) **							
* <sup>142</sup> Ho	D: $p=0$ from 93Li40 **							
<sup>143</sup> Te	-40280# 500#		100# ms (>400 ns)	7/2 <sup>+</sup> #	12		2010	$\beta^-$ ?; $\beta^-n=20\#$ ; $\beta^-2n=2\#$
<sup>143</sup> I	-50630# 200#		130 ms 45	7/2 <sup>+</sup> #	12		1994	$\beta^-$ ?; $\beta^-n=70\#$ ; $\beta^-2n=0.02\#$
<sup>143</sup> Xe	-60203 5		511 ms 6	5/2 <sup>-</sup>	12	03Be05 D	1951	$\beta^-n=100$ ; $\beta^-n=1.00$ 15
<sup>143</sup> Cs	-67676 8		1.791 s 0.007	3/2 <sup>+</sup>	12		1962	$\beta^-n=100$ ; $\beta^-n=1.64$ 7
<sup>143</sup> Ba	-73937 7		14.5 s 0.3	5/2 <sup>-</sup>	12		1962	$\beta^-n=100$
<sup>143</sup> La	-78172 7		14.2 m 0.1	(7/2) <sup>+</sup>	12		1951	$\beta^-n=100$
<sup>143</sup> Ce	-81606.7 2.5		33.039 h 0.006	3/2 <sup>-</sup>	12		1948	$\beta^-n=100$
<sup>143</sup> Pr	-83068.2 1.9		13.57 d 0.02	7/2 <sup>+</sup>	12		1948	$\beta^-n=100$
<sup>143</sup> Nd	-84002.2 1.4		STABLE	7/2 <sup>-</sup>	12		1933	IS=12.174 26
<sup>143</sup> Pm	-82960.7 3.0		265 d 7	5/2 <sup>+</sup>	12		1952	$\epsilon=100$ ; $e^+<5.7e-6$
<sup>143</sup> Sm	-79517.2 2.8		8.75 m 0.06	3/2 <sup>+</sup>	12		1956	$\beta^+n=100$
<sup>143</sup> Sm <sup>m</sup>	-78763.2 2.8	753.99 0.16	66 s 2	11/2 <sup>-</sup>	12		1960	IT $\approx 100$ ; $\beta^+=0.24$ 5
<sup>143</sup> Sm <sup>n</sup>	-76723 3	2793.8 1.3	30 ms 3	23/2 <sup>-</sup>	12	FGK128 J	1969	IT=100 *
<sup>143</sup> Eu	-74241 11		2.59 m 0.02	5/2 <sup>+</sup>	12		1965	$\beta^+n=100$
<sup>143</sup> Eu <sup>m</sup>	-73851 11	389.51 0.04	50.0 $\mu$ s 0.5	11/2 <sup>-</sup>	12		1978	IT=100
<sup>143</sup> Gd	-68230 200		39 s 2	(1/2) <sup>+</sup>	12	78Fi02 D	1975	$\beta^+n=100$ ; $\beta^+p=?$ ; $\beta^+\alpha=?$ *
<sup>143</sup> Gd <sup>m</sup>	-68080 200	152.6 0.5	110.0 s 1.4	11/2 <sup>-</sup>	12	78Fi02 D	1973	$\beta^+n=100$ ; $\beta^+p=?$ ; $\beta^+\alpha=?$ *
<sup>143</sup> Tb	-60420 50		12 s 1	(11/2) <sup>-</sup>	12		1985	$\beta^+n=100$
<sup>143</sup> Tb <sup>m</sup>	-60420# 110#	0# 100#	s <21s	5/2 <sup>+</sup> #	12		1986	$\beta^+?$
<sup>143</sup> Dy	-52169 13		5.6 s 1.0	(1/2) <sup>+</sup>	12	03Xu04 J	1983	$\beta^+n=100$ ; $\beta^+p=?$ *
<sup>143</sup> Dy <sup>m</sup>	-51858 13	310.7 0.6	3.0 s 0.3	(11/2) <sup>-</sup>	12	03Xu04 EJD	2003	$\beta^+n=100$ ; $\beta^+p=?$
<sup>143</sup> Dy <sup>n</sup>	-51763 13	406.3 0.8	1.2 $\mu$ s 0.3		12	05Ri17 E	2005	IT=100 *
<sup>143</sup> Ho	-42050# 300#		300# ms (>200 ns)	11/2 <sup>-</sup> #	12	00So11 I	2000	$\beta^+?$ ; $\beta^+p?$
<sup>143</sup> Er	-31260# 400#		200# ms	9/2 <sup>-</sup> #	12			$\beta^+?$ ; $\beta^+p?$
* <sup>143</sup> Sm <sup>n</sup>	J: E3 to 17/2 <sup>+</sup> **							
* <sup>143</sup> Gd	D: 78Fi02: $\beta^+p$ and/or $\beta^+\alpha$ for <sup>143</sup> Gd+ <sup>143</sup> Gd <sup>m</sup> =0.001%, 39 particles detected **							
* <sup>143</sup> Gd <sup>m</sup>	J: from 05Ba64 **							
* <sup>143</sup> Dy	T: 03Xu04=5.6(1.0); 84Ni03=3.2(0.6) 83Ni05=4.1(0.3) in diff. experiments **							
* <sup>143</sup> Dy <sup>n</sup>	E: 95.6(0.5) above 11/2 <sup>-</sup> isomer **							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>144</sup> I	-45280# 400#		100# ms (>150 ns)	1 <sup>-</sup> #	01	94Be24 I	1994	$\beta^-$ ?; $\beta^-$ -n=40#; $\beta^-$ -2n=1#
<sup>144</sup> Xe	-56872 5		388 ms 7	0 <sup>+</sup>	01	03Be05 TD	2003	$\beta^-$ =100; $\beta^-$ -n=3.0 3
<sup>144</sup> Cs	-63271 20		994 ms 6	1 <sup>(-)</sup>	10		1967	$\beta^-$ =100; $\beta^-$ -n=3.03 13
<sup>144</sup> Cs <sup>m</sup>	-63179 20	92.2 0.5	1.1 $\mu$ s 0.1	(4 <sup>-</sup> )	10		2009	IT=100
<sup>144</sup> Cs <sup>n</sup>	-62970# 200#	300# 200#	< 1 s	(> 3)	10		1978	$\beta^-$ =?; IT ?
<sup>144</sup> Ba	-71767 7		11.5 s 0.2	0 <sup>+</sup>	01		1967	$\beta^-$ =100
<sup>144</sup> La	-74850 13		40.8 s 0.4	(3 <sup>-</sup> )	01		1967	$\beta^-$ =100
<sup>144</sup> Ce	-80431.9 2.9		284.91 d 0.05	0 <sup>+</sup>	01		1945	$\beta^-$ =100
<sup>144</sup> Pr	-80750.5 2.8		17.28 m 0.05	0 <sup>-</sup>	01		1951	$\beta^-$ =100
<sup>144</sup> Pr <sup>m</sup>	-80691.5 2.8	59.03 0.03	7.2 m 0.3	3 <sup>-</sup>	01		1970	IT $\approx$ 100; $\beta^-$ =0.07
<sup>144</sup> Nd	-83748.0 1.4		2.29 Py 0.16	0 <sup>+</sup>	01		1924	IS=23.798 19; $\alpha$ =100
<sup>144</sup> Pm	-81416.1 3.0		363 d 14	5 <sup>-</sup>	01	94Hi05 D	1952	$\epsilon$ =100; $e^+$ <8e-5
<sup>144</sup> Pm <sup>m</sup>	-80575 3	840.90 0.05	780 ns 200	(9 <sup>+</sup> )	01		1993	IT=100
<sup>144</sup> Pm <sup>n</sup>	-72820 4	8595.8 2.2	2.7 $\mu$ s	(27 <sup>+</sup> )	01		1994	IT=100
<sup>144</sup> Sm	-81965.5 1.6		STABLE	0 <sup>+</sup>	01		1933	IS=3.07 7; 2 $\beta^+$ ?
<sup>144</sup> Sm <sup>m</sup>	-79641.9 1.6	2323.60 0.08	880 ns 25	6 <sup>+</sup>	01		1972	IT=100
<sup>144</sup> Eu	-75619 11		10.2 s 0.1	1 <sup>+</sup>	01		1965	$\beta^+$ =100
<sup>144</sup> Eu <sup>m</sup>	-74491 11	1127.6 0.6	1.0 $\mu$ s 0.1	8 <sup>-</sup>	01	FGK127 J	1976	IT=100
<sup>144</sup> Gd	-71760 28		4.47 m 0.06	0 <sup>+</sup>	01		1968	$\beta^+$ =100
<sup>144</sup> Gd <sup>m</sup>	-68327 28	3433.1 0.5	145 ns 30	(10 <sup>+</sup> )	01		1978	IT=100
<sup>144</sup> Tb	-62368 28		1 s	1 <sup>+</sup>	01		1982	$\beta^+$ =100
<sup>144</sup> Tb <sup>m</sup>	-61971 28	396.9 0.5	4.25 s 0.15	(6 <sup>-</sup> )	01		1982	IT=66; $\beta^+$ =34
<sup>144</sup> Tb <sup>n</sup>	-61892 28	476.2 0.5	2.8 $\mu$ s 0.3	(8 <sup>-</sup> )	01		1996	IT=100
<sup>144</sup> Tb <sup>p</sup>	-61851 28	517.1 0.5	670 ns 60	(9 <sup>+</sup> )	01		1996	IT=100
<sup>144</sup> Tb <sup>q</sup>	-61824 28	544.5 0.6	< 300 ns	(10 <sup>+</sup> )	01		1996	IT=100
<sup>144</sup> Dy	-56570 7		9.1 s 0.4	0 <sup>+</sup>	01		1986	$\beta^+$ =100; $\beta^+$ p=?
<sup>144</sup> Ho	-44610 8		700 ms 100	(5 <sup>-</sup> )	08		1986	$\beta^+$ =100; $\beta^+$ p=?
<sup>144</sup> Ho <sup>m</sup>	-44345 8	265.3 0.3	519 ns 5	(8 <sup>+</sup> )	08	10Ma08 T	2001	IT=100
<sup>144</sup> Er	-36610# 200#		400# ms (>200 ns)	0 <sup>+</sup>	06		2003	$\beta^+$ ?
<sup>144</sup> Tm	-22260# 400#		2.3 $\mu$ s 0.9	(10 <sup>+</sup> )	08		2005	p=?; $\beta^+$ =0#
* <sup>144</sup> Ba	D : $\beta^-$ -n=3.6(0.7)% in ENSDF'01 belongs in fact to <sup>144</sup> Cs; $\beta^-$ -n not allowed							
* <sup>144</sup> Eu <sup>m</sup>	J : E2 to 6							
* <sup>144</sup> Tb <sup>m</sup>	T : other 03Li42=12(2)s for q=65 <sup>+</sup> (bare ion)							
* <sup>144</sup> Tm	T : symmetrized from 1.9(+1.2-0.5) $\mu$ s							
<sup>145</sup> I	-40940# 500#		100# ms (>400 ns)	7/2 <sup>+</sup> #	10	100h02 I	2010	$\beta^-$ ?; $\beta^-$ -n=40#; $\beta^-$ -2n=0.3#
<sup>145</sup> Xe	-51493 11		188 ms 4	3/2 <sup>-</sup> #	09		2003	$\beta^-$ =100; $\beta^-$ -n=5.0 6; $\beta^-$ -2n=0#
<sup>145</sup> Cs	-60054 9		582 ms 6	3/2 <sup>+</sup>	09	93Ru01 T	1971	$\beta^-$ =100; $\beta^-$ -n=14.7 9
<sup>145</sup> Cs <sup>m</sup>	-59291 9	762.9 0.4	500 ns 100	19/2 <sup>-</sup> #	09	15YaZW TD	2015	IT=100
<sup>145</sup> Ba	-67516 8		4.31 s 0.16	5/2 <sup>-</sup>	09		1974	$\beta^-$ =100
<sup>145</sup> La	-72835 12		24.8 s 2.0	(5/2 <sup>+</sup> )	09		1974	$\beta^-$ =100
<sup>145</sup> Ce	-77070 30		3.01 m 0.06	5/2 <sup>-</sup> #	09		1954	$\beta^-$ =100
<sup>145</sup> Pr	-79626 7		5.984 h 0.010	7/2 <sup>+</sup>	09		1954	$\beta^-$ =100
<sup>145</sup> Nd	-81432.0 1.4		STABLE	7/2 <sup>-</sup>	09		1933	IS=8.293 12
<sup>145</sup> Pm	-81267.5 2.9		17.7 y 0.4	5/2 <sup>+</sup>	09		1951	$\epsilon$ =100; $\alpha$ =2.8e-7
<sup>145</sup> Sm	-80651.3 1.6		340 d 3	7/2 <sup>-</sup>	09		1947	$\epsilon$ =100
<sup>145</sup> Sm <sup>m</sup>	-71865.1 1.7	8786.2 0.7	990 ns 170	(49/2 <sup>+</sup> )	09		1993	IT=100
<sup>145</sup> Eu	-77992 3		5.93 d 0.04	5/2 <sup>+</sup>	09		1951	$\beta^+$ =100
<sup>145</sup> Eu <sup>m</sup>	-77276 3	716.0 0.3	490 ns 30	11/2 <sup>-</sup>	09		1975	IT=100
<sup>145</sup> Gd	-72926 20		23.0 m 0.4	1/2 <sup>+</sup>	09		1959	$\beta^+$ =100
<sup>145</sup> Gd <sup>m</sup>	-72177 20	749.1 0.2	85 s 3	11/2 <sup>-</sup>	09		1969	IT=94.3 5; $\beta^+$ =5.7 5
<sup>145</sup> Tb	-66390 110		30.9 s 0.6	(11/2 <sup>-</sup> )	09		1981	$\beta^+$ =100
<sup>145</sup> Tb <sup>m</sup>	-65540 200	850 230	BD	(3/2 <sup>+</sup> )	09		1993	$\beta^+$ ?
<sup>145</sup> Dy	-58243 7		9.5 s 1.0	(1/2 <sup>+</sup> )	09	93Al03 T	1982	$\beta^+$ =100; $\beta^+$ p=?
<sup>145</sup> Dy <sup>m</sup>	-58125 7	118.2 0.2	14.1 s 0.7	(11/2 <sup>-</sup> )	09		1982	$\beta^+$ =100; $\beta^+$ p $\approx$ 50
<sup>145</sup> Ho	-49120 7		2.4 s 0.1	11/2 <sup>-</sup> #	09		1987	$\beta^+$ =100
<sup>145</sup> Ho <sup>m</sup>	-49020# 100#	100# 100#	100# ms	5/2 <sup>+</sup> #				$\beta^+$ ?; IT ?
<sup>145</sup> Er	-39240# 200#		900 ms 300	1/2 <sup>+</sup> #	09		1989	$\beta^+$ =100; $\beta^+$ p=?
<sup>145</sup> Er <sup>m</sup>	-39040# 200#	205 4	p	11/2 <sup>-</sup> #	09	10Ma20 T	2010	$\beta^+$ ?
<sup>145</sup> Tm	-27580# 200#		3.17 $\mu$ s 0.20	(11/2 <sup>-</sup> )	09		1998	p=100
* <sup>145</sup> Cs	T : average 93Ru01=579(6) 82Ra13=594(13); other 16Wu.At=613(+32-24)							
* <sup>145</sup> Cs <sup>m</sup>	E : 16Ya.A=762.9(0.4)							
* <sup>145</sup> Sm <sup>m</sup>	T : symmetrized from 960(+190-150)							
* <sup>145</sup> Dy	T : average 93Al03=10.5(1.5) 93To04=6(2) 84Sc.C=10(1)							
* <sup>145</sup> Er	T : 89Vi02=900(300) for mixture gs+isomer; similarly 900(200) from 10Ma20							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>146</sup> Xe	-47955	24			146 ms	6	0 <sup>+</sup>	97 03Be05	TD	1989	$\beta^-$ =100; $\beta^-$ n=6.9 15	
<sup>146</sup> Cs	-55310.4	2.9			323 ms	6	1 <sup>-</sup>	97 93Ru01	T	1971	$\beta^-$ =100; $\beta^-$ n=14.2 5; $\beta^-$ 2n=0#	*
<sup>146</sup> Cs <sup>m</sup>	-55263.7	2.9	46.7	0.1	1.25 $\mu$ s	0.05	4 <sup>-</sup> #	15YaZW	TD	2015	IT=100	*
<sup>146</sup> Ba	-64947	21			2.22 s	0.07	0 <sup>+</sup>	97 93Ru01	D	1970	$\beta^-$ =100	*
<sup>146</sup> La	-69050	30			6.27 s	0.10	2 <sup>-</sup>	97 93Ru01	D	1970	$\beta^-$ =100	*
<sup>146</sup> La <sup>m</sup>	-68920	130	130	130	10.0 s	0.1	(6 <sup>-</sup> )	97 79Ke02	E	1969	$\beta^-$ =100	*
<sup>146</sup> Ce	-75635	16			13.52 m	0.13	0 <sup>+</sup>	97		1953	$\beta^-$ =100	
<sup>146</sup> Pr	-76680	30			24.15 m	0.18	(2) <sup>-</sup>	97		1953	$\beta^-$ =100	
<sup>146</sup> Nd	-80925.9	1.4			STABLE		0 <sup>+</sup>	97		1924	IS=17.189 32; $2\beta^-$ ?; $\alpha$ ?	
<sup>146</sup> Pm	-79454	4			5.53 y	0.05	3 <sup>-</sup>	99		1960	$\epsilon$ =66.0 13; $\beta^-$ =34.0 13	
<sup>146</sup> Sm	-80996	3			68 My	7	0 <sup>+</sup>	97 12Ki16	T	1953	$\alpha$ =100	
<sup>146</sup> Eu	-77118	6			4.61 d	0.03	4 <sup>-</sup>	97		1957	$\beta^+$ =100	
<sup>146</sup> Eu <sup>m</sup>	-76452	6	666.37	0.16	235 $\mu$ s	3	9 <sup>+</sup>	97		1962	IT=100	
<sup>146</sup> Gd	-76086	4			48.27 d	0.10	0 <sup>+</sup>	01		1957	$\epsilon$ =100	
<sup>146</sup> Tb	-67760	40			8 s	4	1 <sup>+</sup>	97		1974	$\beta^+$ =100	
<sup>146</sup> Tb <sup>m</sup>	-67610#	110#	150#	100#	24.1 s	0.5	5 <sup>-</sup>	97 93Al03	T	1974	$\beta^-$ =100	
<sup>146</sup> Tb <sup>n</sup>	-66830#	110#	930#	100#	1.18 ms	0.02	(10 <sup>+</sup> )	97		1989	IT=100	*
<sup>146</sup> Dy	-62555	7			33.2 s	0.7	0 <sup>+</sup>	97 93Al03	T	1981	$\beta^+$ =100	
<sup>146</sup> Dy <sup>m</sup>	-59619	7	2935.7	0.6	150 ms	20	(10 <sup>+</sup> )	97 FGK128	J	1982	IT=100	*
<sup>146</sup> Ho	-51238	7			2.8 s	0.5	(6 <sup>-</sup> )	97 10Ma37	TJ	1982	$\beta^+$ =100; $\beta^+$ p=?	*
<sup>146</sup> Er	-44322	7			1.7 s	0.6	0 <sup>+</sup>	97 93To05	D	1993	$\beta^+$ =100; $\beta^+$ p=?	*
<sup>146</sup> Tm	-31060#	200#			155 ms	20	(1 <sup>+</sup> )	05Ro40	TJD	1993	p $\approx$ 100; $\beta^+$ ?; $\beta^+$ p ?	*
<sup>146</sup> Tm <sup>m</sup>	-30750#	200#	304	6	75 ms	7	(5 <sup>-</sup> )	02 06Ta08	TJ	1993	p $\approx$ 100; $\beta^+$ ?; $\beta^+$ p ?	*
<sup>146</sup> Tm <sup>n</sup>	-30620#	200#	437	7	200 ms	3	(10 <sup>+</sup> )	02 06Ta08	TJ	1993	p=?; $\beta^+$ =16#; $\beta^+$ p ?	*
* <sup>146</sup> Cs	T : average 93Ru01=321(2) 76Lu02=343(7); other 16Wu.A=288(13)										**	
* <sup>146</sup> Cs <sup>m</sup>	E : 16Ya.A=46.7(0.1)										**	
* <sup>146</sup> Ba	D : 93Ru01 $\beta^-$ n<0.02% is not relevant since $Q(\beta^-$ n)=-176(24) is negative										**	
* <sup>146</sup> La	D : 93Ru01 $\beta^-$ n<0.007% is not relevant since $Q(\beta^-$ n)=-50(50) is negative										**	
* <sup>146</sup> La <sup>m</sup>	E : derived from $Q(^{146}\text{La}^m)$ =6660(120) in 79Ke02										**	
* <sup>146</sup> Tb <sup>n</sup>	E : 779.6 keV above <sup>146</sup> Tb <sup>m</sup> , from ENSDF										**	
* <sup>146</sup> Dy <sup>m</sup>	J : E3 to (7 <sup>-</sup> )										**	
* <sup>146</sup> Ho	J : from $\beta^+$ p branching in 10Ma37; supported by $\beta^+$ p spectrum from 85Wi15										**	
* <sup>146</sup> Tm	T : also 05Bb02=190(80) ms										**	
* <sup>146</sup> Tm <sup>m</sup>	T : unweighed average 06Ta08=68(3) 05Ro40=82(4); 05Bb02=75(3)superseded in 06Ta08										**	
* <sup>146</sup> Tm <sup>n</sup>	T : average 07DaZU=213(9) 06Ta08=198(3)										**	
<sup>147</sup> Xe	-42360#	200#			130 ms	80	3/2 <sup>-</sup> #	09		1994	$\beta^-$ =100; $\beta^-$ n=4.0 23; $\beta^-$ 2n=0.01#	*
<sup>147</sup> Cs	-51920	8			230 ms	1	(3/2 <sup>+</sup> )	09		1978	$\beta^-$ =100; $\beta^-$ n=28.5 17	
<sup>147</sup> Cs <sup>m</sup>	-51219	8	701.4	0.4	190 ns	20	19/2 <sup>-</sup> #	15YaZW	TD	2015	IT=100	*
<sup>147</sup> Ba	-60264	20			894 ms	10	5/2 <sup>-</sup>	09 13Rz01	J	1978	$\beta^-$ =100; $\beta^-$ n=0.06 3	
<sup>147</sup> La	-66678	11			4.06 s	0.04	(5/2 <sup>+</sup> )	09 96Ur02	J	1979	$\beta^-$ =100; $\beta^-$ n=0.041 4	
<sup>147</sup> Ce	-72014	9			56.4 s	1.0	(5/2 <sup>-</sup> )	09		1964	$\beta^-$ =100	
<sup>147</sup> Pr	-75444	16			13.4 m	0.3	3/2 <sup>+</sup>	09 15Wa28	J	1964	$\beta^-$ =100	
<sup>147</sup> Nd	-78146.7	1.4			10.98 d	0.01	5/2 <sup>-</sup>	09		1947	$\beta^-$ =100	
<sup>147</sup> Pm	-79042.3	1.4			2.6234 y	0.0002	7/2 <sup>+</sup>	09		1947	$\beta^-$ =100	
<sup>147</sup> Sm	-79266.4	1.4			106.6 Gy	0.7	7/2 <sup>-</sup>	09 09Ko15	T	1933	IS=14.99 18; $\alpha$ =100	
<sup>147</sup> Eu	-77544.8	2.6			24.1 d	0.6	5/2 <sup>+</sup>	09		1951	$\beta^+$ $\approx$ 100; $\alpha$ =0.0022 6	
<sup>147</sup> Eu <sup>m</sup>	-76919.5	2.6	625.27	0.05	765 ns	15	11/2 <sup>-</sup>	09		1970	IT=100	
<sup>147</sup> Gd	-75356.9	2.0			38.06 h	0.12	7/2 <sup>-</sup>	09		1957	$\beta^+$ =100	
<sup>147</sup> Gd <sup>m</sup>	-66769.1	2.1	8587.8	0.5	510 ns	20	(49/2 <sup>+</sup> )	09		1982	IT=100	
<sup>147</sup> Tb	-70743	8			1.64 h	0.03	(1/2 <sup>+</sup> )	09		1969	$\beta^+$ =100	
<sup>147</sup> Tb <sup>m</sup>	-70692	8	50.6	0.9	1.87 m	0.05	11/2 <sup>-</sup> #	09 93Al03	T	1987	$\beta^+$ =100	*
<sup>147</sup> Dy	-64196	9			67 s	7	(1/2 <sup>+</sup> )	09		1975	$\beta^+$ =100; $\beta^+$ p $\approx$ 0.05	
<sup>147</sup> Dy <sup>m</sup>	-63446	9	750.5	0.4	55.2 s	0.5	(11/2 <sup>-</sup> )	09		1976	$\beta^+$ =68.9 23; IT=31.1 23	
<sup>147</sup> Dy <sup>n</sup>	-60789	9	3407.2	0.8	400 ns	10	(27/2 <sup>-</sup> )	09		1985	IT=100	
<sup>147</sup> Ho	-55757	5			5.8 s	0.4	(11/2 <sup>-</sup> )	09		1982	$\beta^+$ =100	
<sup>147</sup> Ho <sup>m</sup>	-53070	5	2687.1	0.4	315 ns	30	(27/2 <sup>-</sup> )	09		1982	IT=100	
<sup>147</sup> Er	-46610	40			3.2 s	1.2	(1/2 <sup>+</sup> )	09 10Ma27	T	1992	$\beta^+$ =100; $\beta^+$ p=?	
<sup>147</sup> Er <sup>m</sup>	-46510#	60#	100#	50#	1.6 s	0.2	(11/2 <sup>-</sup> )	09 10Ma27	T	1982	$\beta^+$ =100; $\beta^+$ p=?	*
<sup>147</sup> Tm	-35974	7			580 ms	30	11/2 <sup>-</sup>	09		1982	$\beta^+$ =85 5; p=15 5	
<sup>147</sup> Tm <sup>m</sup>	-35913	7	62	5	360 $\mu$ s	40	3/2 <sup>+</sup>	09		1984	p=100	
* <sup>147</sup> Xe	T : symmetrized from 100(+100-50) D : from $\beta^-$ n<8%										**	
* <sup>147</sup> Cs <sup>m</sup>	E : 16Ya.A=701.4(0.4)										**	
* <sup>147</sup> Tb <sup>m</sup>	T : average 93Al03=1.92(0.07) 73Bo13=1.83(0.06)										**	
* <sup>147</sup> Er <sup>m</sup>	E : estimated from 11/2 <sup>-</sup> level in isotones <sup>141</sup> Sm=175 <sup>143</sup> Gd=152 <sup>145</sup> Dy=118										**	



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
<sup>148</sup> Xe	-38600#	300#		100# ms (>400 ns)	0 <sup>+</sup>	14	100h02	I	2010	$\beta^-$ ?; $\beta^-$ -n=10#; $\beta^-$ -2n=0#		
<sup>148</sup> Cs	-46911	13		145 ms	4	14	16Wu.A	T	1978	$\beta^-$ =100; $\beta^-$ -n=25.1 25; $\beta^-$ -2n=0#	*	
<sup>148</sup> Cs <sup>m</sup>	-46866	13	45.2 0.1	4.8 $\mu$ s	0.2	4 <sup>-</sup> #	15YaZW	TD	2015	IT=100	*	
<sup>148</sup> Ba	-57590	60		620 ms	5	0 <sup>+</sup>	16Wu.A	T	1979	$\beta^-$ =100; $\beta^-$ -n=0.4 3	*	
<sup>148</sup> La	-62709	19		1.35 s	0.04	(2 <sup>-</sup> )	16Wu.A	T	1969	$\beta^-$ =100; $\beta^-$ -n=0.15 3	*	
<sup>148</sup> Ce	-70398	11		56.8 s	0.3	0 <sup>+</sup>			1964	$\beta^-$ =100		
<sup>148</sup> Pr	-72535	15		2.29 m	0.02	1 <sup>-</sup>			1964	$\beta^-$ =100		
<sup>148</sup> Pt <sup>m</sup>	-72458	15	76.80 0.20	2.01 m	0.07	(4)			1964	$\beta^-$ =64 10; IT=36 10		
<sup>148</sup> Nd	-77408.0	2.1		STABLE	(>3.0 Ey)	0 <sup>+</sup>	82Be20	T	1937	IS=5.756 21; 2 $\beta^-$ ?; $\alpha$ ?	*	
<sup>148</sup> Pm	-76866	6		5.368 d	0.007	1 <sup>-</sup>			1947	$\beta^-$ =100		
<sup>148</sup> Pm <sup>m</sup>	-76728	6	137.9 0.3	41.29 d	0.11	5 <sup>-</sup> , 6 <sup>-</sup>			1951	$\beta^-$ =95.8 6; IT=4.2 6		
<sup>148</sup> Sm	-79336.3	1.4		6.3 Py	1.3	0 <sup>+</sup>	16Ca.1	T	1933	IS=11.24 10; $\alpha$ =100	*	
<sup>148</sup> Eu	-76299	10		54.5 d	0.5	5 <sup>-</sup>			1951	$\beta^+$ =100; $\alpha$ =9.4e-7 28		
<sup>148</sup> Eu <sup>m</sup>	-75579	10	720.4 0.3	162 ns	8	9 <sup>+</sup>			1980	IT=100		
<sup>148</sup> Gd	-76269.3	1.6		70.9 y	1.0	0 <sup>+</sup>	03Fu10	T	1953	$\alpha$ =100; 2 $\beta^+$ ?	*	
<sup>148</sup> Tb	-70537	12		60 m	1	2 <sup>-</sup>			1960	$\beta^+$ =100		
<sup>148</sup> Tb <sup>m</sup>	-70447	12	90.1 0.3	2.20 m	0.05	(9) <sup>+</sup>			1973	$\beta^+$ =100		
<sup>148</sup> Tb <sup>n</sup>	-61918	12	8618.6 1.0	1.310 $\mu$ s	0.007	(27 <sup>+</sup> )			1980	IT=100		
<sup>148</sup> Dy	-67860	9		3.3 m	0.2	0 <sup>+</sup>			1974	$\beta^+$ =100		
<sup>148</sup> Dy <sup>m</sup>	-64941	9	2919.1 1.0	471 ns	20	10 <sup>+</sup>			1978	IT=100		
<sup>148</sup> Ho	-57990	80		2.2 s	1.1	(1 <sup>+</sup> )			1979	$\beta^+$ =100		
<sup>148</sup> Ho <sup>m</sup>	-57740#	130#	250# 100#	9.49 s	0.12	5# <sup>(-)</sup>	93Al03	T	1979	$\beta^+$ =100; $\beta^+$ p=0.08 1	*	
<sup>148</sup> Ho <sup>n</sup>	-57050#	130#	940# 100#	2.36 ms	0.06	(10) <sup>+</sup>			1984	IT=100	*	
<sup>148</sup> Er	-51479	10		4.6 s	0.2	0 <sup>+</sup>			1982	$\beta^+$ =100; $\beta^+$ p $\approx$ 0.15		
<sup>148</sup> Er <sup>m</sup>	-48566	10	2913.2 0.4	13 $\mu$ s	3	(10 <sup>+</sup> )			1982	IT=100		
<sup>148</sup> Tm	-38765	10		700 ms	200	(10 <sup>+</sup> )			1982	$\beta^+$ =100; $\beta^+$ p ?		
<sup>148</sup> Yb	-30330#	400#		250# ms		0 <sup>+</sup>				$\beta^+$ ?; $\beta^+$ p ?		
* <sup>148</sup> Cs	T : average 16Wu.A=144(5) 93Ru01=140(12) 86Hi08=158(7) 86Wa17=130(10) and										**	
* <sup>148</sup> Cs	T : 78Ko29=130(40)										**	
* <sup>148</sup> Cs <sup>m</sup>	E : 16Ya.A=45.2(0.1)										**	
* <sup>148</sup> Ba	T : average 16Wu.A=621(11) 86Wa17=620(5) 84Ch02=607(25) 82Ga24=630(50)										**	
* <sup>148</sup> La	T : unweighed average 16Wu.A=1.27(+0.10-0.09) 86Wa17=1.40(0.02)										**	
* <sup>148</sup> La	T : 93Ru01=1.428(0.012) and 69Wi.A=1.29(0.08)										**	
* <sup>148</sup> Nd	T : lower limit is for 2 $\beta^-$ decay										**	
* <sup>148</sup> Sm	T : symmetrized from 16Ca.1=6.4(+1.2-1.3) Py										**	
* <sup>148</sup> Gd	T : 81Pr06=74.6(3.0) unweighed not used										**	
* <sup>148</sup> Ho <sup>m</sup>	T : average 93Al03=9.30(0.20) 89Ta11=9.59(0.15)										**	
* <sup>148</sup> Ho <sup>n</sup>	E : 694.4 keV above <sup>148</sup> Ho <sup>m</sup> , from ENSDF										**	
<sup>149</sup> Cs	-43250#	400#		113 ms	8	3/2 <sup>+</sup> #	04 16Wu.A	TD	1979	$\beta^-$ =100; $\beta^-$ -n=60#; $\beta^-$ -2n=0#		
<sup>149</sup> Ba	-53120	440		348 ms	4	3/2 <sup>-</sup> #	04 16Wu.A	T	1993	$\beta^-$ =100; $\beta^-$ -n=0.43 12	*	
<sup>149</sup> La	-60220	200		1.07 s	0.02	(3/2 <sup>-</sup> )	07 16Wu.A	T	1979	$\beta^-$ =100; $\beta^-$ -n=1.43 28	*	
<sup>149</sup> Ce	-66670	10		4.94 s	0.04	3/2 <sup>-</sup> #	04 96Ya.A	T	1974	$\beta^-$ =100		
<sup>149</sup> Pr	-71039	10		2.26 m	0.07	(5/2 <sup>+</sup> )	04		1964	$\beta^-$ =100		
<sup>149</sup> Nd	-74375.5	2.1		1.728 h	0.001	5/2 <sup>-</sup>	04		1938	$\beta^-$ =100		
<sup>149</sup> Pm	-76064.3	2.3		53.08 h	0.05	7/2 <sup>+</sup>	04		1947	$\beta^-$ =100		
<sup>149</sup> Pm <sup>m</sup>	-75824.1	2.3	240.214 0.007	35 $\mu$ s	3	11/2 <sup>-</sup>	04		1966	IT=100		
<sup>149</sup> Sm	-77135.7	1.3		STABLE	(>2 Py)	7/2 <sup>-</sup>	04		1933	IS=13.82 7; $\alpha$ ?		
<sup>149</sup> Eu	-76441	4		93.1 d	0.4	5/2 <sup>+</sup>	04		1959	$\epsilon$ =100		
<sup>149</sup> Eu <sup>m</sup>	-75945	4	496.386 0.002	2.45 $\mu$ s	0.05	11/2 <sup>-</sup>	04		1961	IT=100		
<sup>149</sup> Gd	-75127	3		9.28 d	0.10	7/2 <sup>-</sup>	04		1951	$\beta^+$ =100; $\alpha$ =4.3e-4 10		
<sup>149</sup> Tb	-71489	4		4.118 h	0.025	1/2 <sup>+</sup>	04		1950	$\beta^+$ =83.3 17; $\alpha$ =16.7 17		
<sup>149</sup> Tb <sup>m</sup>	-71453	4	35.78 0.13	4.16 m	0.04	11/2 <sup>-</sup>	04		1962	$\beta^+$ $\approx$ 100; $\alpha$ =0.022 3		
<sup>149</sup> Dy	-67696	9		4.20 m	0.14	7/2 <sup>(-)</sup>	04	88Ah02	J	1958	$\beta^+$ =100	
<sup>149</sup> Dy <sup>m</sup>	-65035	9	2661.1 0.4	490 ms	15	(27/2 <sup>-</sup> )	04		1976	IT=99.3 3; $\beta^+$ =0.7 3	*	
<sup>149</sup> Ho	-61647	12		21.1 s	0.2	(11/2 <sup>-</sup> )	04		1979	$\beta^+$ =100		
<sup>149</sup> Ho <sup>m</sup>	-61598	12	48.80 0.20	56 s	3	(1/2 <sup>+</sup> )	04		1988	$\beta^+$ =100		
<sup>149</sup> Er	-53742	28		4 s	2	(1/2 <sup>+</sup> )	04		1984	$\beta^+$ =100; $\beta^+$ p=7 2		
<sup>149</sup> Er <sup>m</sup>	-53000	28	741.8 0.2	8.9 s	0.2	(11/2 <sup>-</sup> )	04		1984	$\beta^+$ =96.5 7; IT=3.5 7; $\beta^+$ p=0.18 7		
<sup>149</sup> Er <sup>n</sup>	-51131	28	2611.1 0.3	610 ns	80	(19/2 <sup>+</sup> )	04		1987	IT=100		
<sup>149</sup> Er <sup>p</sup>	-50470	30	3272 20	4.8 $\mu$ s	0.1	(27/2 <sup>-</sup> )	04		1987	IT=100	*	
<sup>149</sup> Tm	-43880#	200#		900 ms	200	(11/2 <sup>-</sup> )	04		1987	$\beta^+$ =100; $\beta^+$ p=0.26 15	*	
<sup>149</sup> Yb	-33200#	300#		700 ms	200	(1/2 <sup>+</sup> )	04	05Xu04	J	2001	$\beta^+$ =100; $\beta^+$ p $\approx$ 100	*
* <sup>149</sup> Ba	T : average 16Wu.A=352(6) 93Ru01=324(18) 86Wa17=346(6)										**	
* <sup>149</sup> La	T : average 16Wu.A=1.11(0.04) 93Ru01=1.066(0.034) 86Wa17=1.04(0.04)										**	
* <sup>149</sup> Dy <sup>m</sup>	T : other 03Li42=11(1)s for q=66 <sup>+</sup> (bare ion)										**	
* <sup>149</sup> Er <sup>p</sup>	E : 3242.7 + 30(20) keV										**	
* <sup>149</sup> Tm	D : symmetrized from $\beta^+$ p=0.2(+0.2-0.1)%										**	
* <sup>149</sup> Yb	J : (1/2 <sup>+</sup> , 3/2 <sup>+</sup> ) in ENSDF2004 and 1/2 in 05Xu04; 06Xu07=(1/2 <sup>-</sup> ) however,										**	
* <sup>149</sup> Yb	J : no 1/2 <sup>-</sup> ground-state or isomer for e-o in this region										**	

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>150</sup> Cs	-38170#	400#	84.4 ms	8.2		13 16Wu.A TD	1979	$\beta^-$ =100; $\beta^-$ n=80#; $\beta^-$ 2n=2#	
<sup>150</sup> Ba	-49900#	300#	259 ms	5		13 16Wu.A T	1994	$\beta^-$ =100; $\beta^-$ n=1#	
<sup>150</sup> La	-56130	440	504 ms	15	(3 <sup>+</sup> )	13 16Wu.A T	1993	$\beta^-$ =100; $\beta^-$ n=2.7 3 *	
<sup>150</sup> Ce	-64847	12	6.05 s	0.07	0 <sup>+</sup>	13 15Ko23 T	1970	$\beta^-$ =100	
<sup>150</sup> Pr	-68300	9	6.19 s	0.16	1 <sup>-</sup>	13 15Ko23 J	1970	$\beta^-$ =100 *	
<sup>150</sup> Nd	-73679.8	1.3	8.2 Ey	0.9	0 <sup>+</sup>	13 15Ba11 T	1937	IS=5.638 28; $2\beta^-$ =100 *	
<sup>150</sup> Pm	-73597	20	2.698 h	0.015	(1 <sup>-</sup> )	13	1952	$\beta^-$ =100	
<sup>150</sup> Sm	-77051.1	1.3	STABLE		0 <sup>+</sup>	13	1934	IS=7.38 1	
<sup>150</sup> Eu	-74792	6	36.9 y	0.9	5 <sup>(-#)</sup>	13	1950	$\beta^+$ =100	
<sup>150</sup> Eu <sup>m</sup>	-74750	6	41.7	1.0	12.8 h	0.1	1953	$\beta^-$ =89 2; $\beta^+$ =11 2; IT <sub>≤</sub> 5e-8	
<sup>150</sup> Gd	-75764	6	1.79 My	0.08	0 <sup>+</sup>	13	1953	$\alpha$ =100; $2\beta^+$ ?	
<sup>150</sup> Tb	-71106	7	3.48 h	0.16	(2 <sup>-</sup> )	13	1959	$\beta^+$ ≈100; $\alpha$ <0.05	
<sup>150</sup> Tb <sup>m</sup>	-70645	26	461	27	5.8 m	0.2	1993	$\beta^+$ ≈100; IT ?	
<sup>150</sup> Dy	-69310	4	7.17 m	0.05	0 <sup>+</sup>	13	1959	$\beta^+$ =64 5; $\alpha$ =36 5	
<sup>150</sup> Ho	-61946	14	76.8 s	1.8	(2 <sup>-</sup> )	13 93Al03 T	1963	$\beta^+$ =100 *	
<sup>150</sup> Ho <sup>m</sup>	-61950	50	-0	50	23.3 s	0.3	1980	$\beta^+$ =100	
<sup>150</sup> Ho <sup>n</sup>	-54050	50	7900	50	787 ns	36	2006	IT=100 *	
<sup>150</sup> Er	-57831	17	18.5 s	0.7	0 <sup>+</sup>	13	1982	$\beta^+$ =100	
<sup>150</sup> Er <sup>m</sup>	-55035	17	2796.5	0.5	2.55 μs	0.10	1984	IT=100	
<sup>150</sup> Tm	-46490#	200#	* &	3#	s	(1 <sup>+</sup> )	88Ni02 J	1982	$\beta^+$ =100
<sup>150</sup> Tm <sup>m</sup>	-46350#	240#	140#	140#	* &	2.20 s	0.06	1981	$\beta^+$ =100; $\beta^+$ p=1.2 3
<sup>150</sup> Tm <sup>n</sup>	-45680#	240#	810#	140#	5.2 ms	0.3	1984	IT=100 *	
<sup>150</sup> Yb	-38640#	300#	700#	ms	(>200ns)	0 <sup>+</sup>	13	2000	$\beta^+$ ?
<sup>150</sup> Lu	-24640#	300#	45 ms	3	(5 <sup>-</sup> , 6 <sup>-</sup> )	13 00Gi01 J	1993	p=?; $\beta^+$ =29#	
<sup>150</sup> Lu <sup>m</sup>	-24620#	300#	22	5	40 μs	7	1998	p=100	
* <sup>150</sup> La	T : average 16Wu.At=510(+10-22) 95Ok02=510(30) **								
* <sup>150</sup> Pr	T : also 15Ko23=8.2 s (no unc.) is "apparent" value direct+growth from 150Ce **								
* <sup>150</sup> Nd	T : and 15Ba11=120(+30-20) to first exc. 0 <sup>+</sup> state **								
* <sup>150</sup> Ho	T : average 93Al03=78(2) 82No08=72(4) **								
* <sup>150</sup> Ho <sup>n</sup>	E : 7912.1(2.3) keV above the (9) <sup>+</sup> isomer **								
* <sup>150</sup> Tm <sup>n</sup>	E : 671.3(1.0) keV above <sup>150</sup> Tm <sup>m</sup> , from ENSDF **								
* <sup>150</sup> Lu <sup>m</sup>	T : symmetrized from 03Gi10=39(+8-6) **								
<sup>151</sup> Cs	-34230#	500#	69 ms	26	3/2 <sup>+</sup> #	09 16Wu.A TD	1979	$\beta^-$ =100; $\beta^-$ n=90#; $\beta^-$ 2n=0.4#	
<sup>151</sup> Ba	-44940#	400#	167 ms	5	3/2 <sup>-</sup> #	09 16Wu.A TD	1994	$\beta^-$ =100; $\beta^-$ n=7#	
<sup>151</sup> La	-53310	440	465 ms	24	5/2 <sup>+</sup> #	09 16Wu.A TD	1994	$\beta^-$ =100; $\beta^-$ n=6# *	
<sup>151</sup> Ce	-61225	18	1.76 s	0.06	(3/2 <sup>-</sup> )	09 10Si03 J	1997	$\beta^-$ =100	
<sup>151</sup> Pr	-66780	12	18.90 s	0.07	(3/2 <sup>-</sup> )	09	1990	$\beta^-$ =100	
<sup>151</sup> Pr <sup>m</sup>	-66745	12	35.10	0.10	50 μs	8	2006	IT=100	
<sup>151</sup> Nd	-70943.0	1.3	12.44 m	0.07	3/2 <sup>+</sup>	09	1938	$\beta^-$ =100	
<sup>151</sup> Pm	-73386	5	28.40 h	0.04	5/2 <sup>+</sup>	09	1952	$\beta^-$ =100	
<sup>151</sup> Sm	-74576.3	1.3	90 y	8	5/2 <sup>-</sup>	09	1947	$\beta^-$ =100	
<sup>151</sup> Sm <sup>m</sup>	-74315.2	1.3	261.13	0.04	1.4 μs	0.1	1973	IT=100	
<sup>151</sup> Eu	-74652.9	1.3	4.6 Ey	1.2	5/2 <sup>+</sup>	09 14Ca13 T	1933	IS=47.81 6; $\alpha$ =100	
<sup>151</sup> Eu <sup>m</sup>	-74456.7	1.3	196.245	0.010	58.9 μs	0.5	1958	IT=100	
<sup>151</sup> Gd	-74189	3	123.9 d	1.0	7/2 <sup>-</sup>	09	1950	$\epsilon$ =100; $\alpha$ =1.0e-6 6 *	
<sup>151</sup> Tb	-71624	4	17.609 h	0.001	1/2 <sup>(+)</sup>	09	1953	$\beta^+$ ≈100; $\alpha$ =0.0095 15	
<sup>151</sup> Tb <sup>m</sup>	-71524	4	99.53	0.05	25 s	3	1978	IT=93.4 20; $\beta^+$ =6.6 20	
<sup>151</sup> Dy	-68752	3	17.9 m	0.3	7/2 <sup>(-)</sup>	09	1959	$\beta^+$ =?; $\alpha$ =5.6 4	
<sup>151</sup> Ho	-63623	8	35.2 s	0.1	11/2 <sup>(-)</sup>	09 87Ne.A J	1963	$\beta^+$ =?; $\alpha$ =22 3	
<sup>151</sup> Ho <sup>m</sup>	-63582	8	41.0	0.2	47.2 s	1.3	1963	$\alpha$ =77 18; $\beta^+$ ? *	
<sup>151</sup> Er	-58266	16	23.5 s	2.0	(7/2 <sup>-</sup> )	09	1970	$\beta^+$ =100	
<sup>151</sup> Er <sup>m</sup>	-55680	16	2586.0	0.5	580 ms	20	1980	IT=95.3 3; $\beta^+$ =4.7 3 *	
<sup>151</sup> Er <sup>n</sup>	-47979	16	10286.6	1.0	420 ns	50	1990	IT=100	
<sup>151</sup> Tm	-50773	19	4.17 s	0.11	(65/2 <sup>-</sup> , 61/2 <sup>+</sup> )	09 09Fu05 J	1982	$\beta^+$ =100	
<sup>151</sup> Tm <sup>m</sup>	-50679	20	94	6	AD	6.6 s	2.0	1987	$\beta^+$ =100
<sup>151</sup> Tm <sup>n</sup>	-48117	19	2655.67	0.22	451 ns	34	1982	IT=100	
<sup>151</sup> Yb	-41540	300	1.6 s	0.5	(1/2 <sup>+</sup> )	09 86To12 T	1985	$\beta^+$ =100; $\beta^+$ p=? *	
<sup>151</sup> Yb <sup>m</sup>	-40790#	320#	750#	100#	1.6 s	0.5	1986	$\beta^+$ ≈100; $\beta^+$ p=?; IT=0.4# *	
<sup>151</sup> Yb <sup>n</sup>	-39000#	580#	2450#	500#	2.6 μs	0.7	1993	IT=100 *	
<sup>151</sup> Yb <sup>p</sup>	-39090#	580#	2450#	500#	20 μs	1	1987	IT=100 *	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>151</sup> Lu	-30110# 300#		78.4 ms 0.9	(11/2 <sup>-</sup> )	09	15Ta12 TJ	1982	p=?; β <sup>+</sup> =37#
<sup>151</sup> Lu <sup>m</sup>	-30060# 300#	53	16.5 μs 0.7	(3/2 <sup>+</sup> )	09	15Ta12 TJ	1998	p=100
* <sup>151</sup> La	T: symmetrized from 457(+30-18)							**
* <sup>151</sup> Ce	T: average 16Wu.A=1.71(0.09) 06Ko25=1.76 (0.06)							**
* <sup>151</sup> Ce	I: isomer with T=1.02(0.06) suggested in ENSDF2009 not trusted by NUBASE							**
* <sup>151</sup> Gd	D: symmetrized from α=0.8(+0.8-0.4)e-6%							**
* <sup>151</sup> Ho <sup>m</sup>	D: symmetrized from α=80(+15-20)%							**
* <sup>151</sup> Er <sup>m</sup>	T: other 03Li42=19(3)s for q=68 <sup>+</sup> (bare ion)							**
* <sup>151</sup> Yb	T: derived from 1.6(0.1), for mixture of ground-state and isomer with almost same half-life							**
* <sup>151</sup> Yb <sup>m</sup>	E: 740# estimated in 90Ak01 (see ENSDF'09)							**
* <sup>151</sup> Yb <sup>n</sup>	E: above the 1791.2 keV level above <sup>151</sup> Yb <sup>m</sup> (see ENSDF'09)							**
* <sup>151</sup> Yb <sup>p</sup>	E: 2448 keV above <sup>151</sup> Yb <sup>m</sup> (see ENSDF'97)							**
* <sup>151</sup> Lu	D: p=63.4(0.9)% in ENSDF'09, based on predicted β <sup>+</sup> decay half-life≈220 ms							**
* <sup>151</sup> Lu	T: average 15Ta12=78(1) 99Bi14=80(2)							**
* <sup>151</sup> Lu <sup>m</sup>	T: average 15Ta12=17(1) 99Bi14=16(1)							**
<sup>152</sup> Cs	-28930# 500#		30# ms					β <sup>-</sup> ?; β <sup>-</sup> n ?
<sup>152</sup> Ba	-41710# 400#		139 ms 8	0 <sup>+</sup>	13	16Wu.A TD	2010	β <sup>-</sup> =100; β <sup>-</sup> n=5#
<sup>152</sup> La	-49290# 300#		287 ms 16		13	16Wu.A TD	1994	β <sup>-</sup> =100; β <sup>-</sup> n=50#
<sup>152</sup> Ce	-58980# 200#		1.42 s 0.02	0 <sup>+</sup>	13	16Wu.A T	1990	β <sup>-</sup> =100
<sup>152</sup> Pr	-63758 19		3.57 s 0.11	4 <sup>+</sup>	13	99To04 J	1983	β <sup>-</sup> =100
<sup>152</sup> Pr <sup>m</sup>	-63643 19	114.8	4.1 μs 0.1	(3 <sup>+</sup> )	13		1990	IT=100
<sup>152</sup> Nd	-70149 24		11.4 m 0.2	0 <sup>+</sup>	13		1969	β <sup>-</sup> =100
<sup>152</sup> Pm	-71254 26		4.12 m 0.08	1 <sup>+</sup>	13		1958	β <sup>-</sup> =100
<sup>152</sup> Pm <sup>m</sup>	-71110 80	140	7.52 m 0.08	4 <sup>-</sup>	13		1971	β <sup>-</sup> =100
<sup>152</sup> Pm <sup>n</sup>	-71000# 150#	250#	13.8 m 0.2	(8)	13		1971	β <sup>-</sup> ≈100; IT=?
<sup>152</sup> Sm	-74762.6 1.2		STABLE	0 <sup>+</sup>	13		1933	IS=26.75 16
<sup>152</sup> Eu	-72888.3 1.3		13.517 y 0.009	3 <sup>-</sup>	13		1938	β <sup>+</sup> =72.08 13; β <sup>-</sup> =27.92 13
<sup>152</sup> Eu <sup>m</sup>	-72842.7 1.3	45.5998	9.3116 h 0.0013	0 <sup>-</sup>	13		1958	β <sup>-</sup> =73 3; β <sup>+</sup> =27 3
<sup>152</sup> Eu <sup>n</sup>	-72823.0 1.3	65.2969	940 ns 80	1 <sup>-</sup>	13		1978	IT=100
<sup>152</sup> Eu <sup>p</sup>	-72810.1 1.3	78.2331	165 ns 10	1 <sup>+</sup>	13		1978	IT=100
<sup>152</sup> Eu <sup>q</sup>	-72798.5 1.3	89.8496	384 ns 10	4 <sup>+</sup>	13		1970	IT=100
<sup>152</sup> Eu <sup>r</sup>	-72740.4 1.3	147.86	95.8 m 0.4	8 <sup>-</sup>	13	15Hu02 T	1963	IT=100
<sup>152</sup> Gd	-74706.9 1.2		108 Ty 8	0 <sup>+</sup>	13		1938	IS=0.20 1; α=100; 2β <sup>+</sup> ?
<sup>152</sup> Tb	-70720 40		17.5 h 0.1	2 <sup>-</sup>	13		1959	β <sup>+</sup> =100; α<7e-7
<sup>152</sup> Tb <sup>m</sup>	-70380 40	342.15	960 ns 10	5 <sup>-</sup>	13		1972	IT=100
<sup>152</sup> Tb <sup>n</sup>	-70220 40	501.74	4.2 m 0.1	8 <sup>+</sup>	13		1971	IT=78.9 6; β <sup>+</sup> =21.1 6
<sup>152</sup> Dy	-70118 5		2.38 h 0.02	0 <sup>+</sup>	13		1958	ε≈100; α=0.100 7
<sup>152</sup> Ho	-63605 13		161.8 s 0.3	2 <sup>-</sup>	13		1963	β <sup>+</sup> =88 3; α=12 3
<sup>152</sup> Ho <sup>m</sup>	-63445 13	160	49.8 s 0.2	9 <sup>+</sup>	13		1963	β <sup>+</sup> =89.2 17; α=10.8 17
<sup>152</sup> Ho <sup>n</sup>	-60585 13	3019.59	8.4 μs 0.3	19 <sup>-</sup>	13		1997	IT=100
<sup>152</sup> Er	-60500 9		10.3 s 0.1	0 <sup>+</sup>	13		1963	α=90 4; β <sup>+</sup> =10 4
<sup>152</sup> Tm	-51720 50		8.0 s 1.0	(2#) <sup>-</sup>	13		1980	β <sup>+</sup> =100
<sup>152</sup> Tm <sup>m</sup>	-51820 240	-100	5.2 s 0.6	(9) <sup>+</sup>	13		1980	β <sup>+</sup> =100
<sup>152</sup> Tm <sup>n</sup>	-49060# 140#	2665#	294 ns 12	(17 <sup>+</sup> )	13		1986	IT=100
<sup>152</sup> Yb	-46270 150		3.03 s 0.06	0 <sup>+</sup>	13		1982	β <sup>+</sup> =100
<sup>152</sup> Yb <sup>m</sup>	-43530 150	2744.5	30 μs 1	(10 <sup>+</sup> )	13		1995	IT=100
<sup>152</sup> Lu	-33420# 200#		650 ms 70	(4 <sup>-</sup> , 5 <sup>-</sup> , 6 <sup>-</sup> )	13	88Ni02 T	1987	β <sup>+</sup> =100; β <sup>+</sup> p=15 7
* <sup>152</sup> La	T: symmetrized from 298(+6-23)							**
* <sup>152</sup> Pr	T: average 90An31=3.7(0.2) 85Br08=3.8(0.2) 83Hi05=3.24(0.19)							**
* <sup>152</sup> Pm <sup>n</sup>	E: ENSDF: "Probably feeds 7.52 m level" at 140 keV							**
* <sup>152</sup> Tm <sup>n</sup>	E: 2555.05(0.19) above <sup>152</sup> Tm <sup>m</sup>							**
* <sup>152</sup> Lu	T: average 88Ni02=600(100) 87To02=700(100)							**
<sup>153</sup> Ba	-36470# 400#		116 ms 52	5/2 <sup>-</sup> #		16Wu.A TD	2016	β <sup>-</sup> =100; β <sup>-</sup> n=3#; β <sup>-</sup> 2n=0#
<sup>153</sup> La	-46060# 300#		245 ms 18	5/2 <sup>+</sup> #	06	16Wu.A TD	1994	β <sup>-</sup> =100; β <sup>-</sup> n=50#
<sup>153</sup> Ce	-54910# 200#		865 ms 25	3/2 <sup>-</sup> #	06	16Wu.A TD	1994	β <sup>-</sup> =100; β <sup>-</sup> n=0.01#
<sup>153</sup> Pr	-61568 12		4.28 s 0.11	5/2 <sup>-</sup> #	06		1987	β <sup>-</sup> =100; β <sup>-</sup> n=0.02#
<sup>153</sup> Nd	-67330.3 2.7		31.6 s 1.0	(3/2) <sup>-</sup>	06		1987	β <sup>-</sup> =100
<sup>153</sup> Nd <sup>m</sup>	-67138.6 2.9	191.7	1.10 μs 0.04	(5/2 <sup>+</sup> )	06	10Si03 TJ	1996	IT=100
<sup>153</sup> Pm	-70648 9		5.25 m 0.02	5/2 <sup>-</sup>	06		1962	β <sup>-</sup> =100
<sup>153</sup> Sm	-72559.7 1.2		46.284 h 0.004	3/2 <sup>+</sup>	06		1938	β <sup>-</sup> =100
<sup>153</sup> Sm <sup>m</sup>	-72461.3 1.2	98.37	10.6 ms 0.3	11/2 <sup>-</sup>	06		1971	IT=100
<sup>153</sup> Eu	-73367.2 1.3		STABLE (>550 Py)	5/2 <sup>+</sup>	06	12Da16 T	1933	IS=52.19 6
<sup>153</sup> Eu <sup>m</sup>	-71596.2 1.4	1771.0	475 ns 10	19/2 <sup>-</sup>	06		2000	IT=100
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>153</sup> Gd	-72882.6	1.2	240.4 d	1.0	3/2 <sup>-</sup>	06	1947	$\epsilon=100$
<sup>153</sup> Gd <sup>m</sup>	-72787.4	1.2	95.1736	0.0008	3.5 $\mu$ s	0.4	1979	IT=100
<sup>153</sup> Gd <sup>n</sup>	-72711.4	1.2	171.188	0.004	76.0 $\mu$ s	1.4	1967	IT=100
<sup>153</sup> Tb	-71313	4			2.34 d	0.01	1957	$\beta^+=100$
<sup>153</sup> Tb <sup>m</sup>	-71150	4	163.175	0.005	186 $\mu$ s	4	1965	IT=100
<sup>153</sup> Dy	-69143	4			6.4 h	0.1	1958	$\beta^+\approx 100; \alpha=0.0094$ 14
<sup>153</sup> Ho	-65012	5			2.01 m	0.03	1963	$\beta^+\approx 100; \alpha=0.051$ 25
<sup>153</sup> Ho <sup>m</sup>	-64943	5	68.7	0.3	9.3 m	0.5	1963	$\beta^+\approx 100; \alpha=0.18$ 8
<sup>153</sup> Ho <sup>n</sup>	-62240	11	2772	10	229 ns	2	1980	IT=100
<sup>153</sup> Er	-60469	9			37.1 s	0.2	1963	$\alpha=53$ 3; $\beta^+=47$ 3
<sup>153</sup> Er <sup>m</sup>	-57671	9	2798.2	1.0	373 ns	9	1979	IT=100
<sup>153</sup> Er <sup>n</sup>	-55221	9	5248.1	1.0	248 ns	32	1979	IT=100
<sup>153</sup> Tm	-53973	12			1.48 s	0.01	1964	$\alpha=91$ 3; $\beta^+=9$ 3
<sup>153</sup> Tm <sup>m</sup>	-53930	12	43.2	0.2	2.5 s	0.2	1988	$\alpha=92$ 3; $\beta^+=?$
<sup>153</sup> Yb	-47210#	200#			4.2 s	0.2	1977	$\beta^+=?$ ; $\alpha=50$ #; $\beta^+p=0.008$ 2
<sup>153</sup> Yb <sup>n</sup>	-44510#	220#	2700	100	15 $\mu$ s	1	1989	IT=100
<sup>153</sup> Lu	-38370	150			900 ms	200	1989	$\beta^+ ?; \alpha=?; p=0$
<sup>153</sup> Lu <sup>m</sup>	-38290	150	80	5	1#	s	1997	$\beta^+ ?; \alpha=?; IT=?; p=0$
<sup>153</sup> Lu <sup>n</sup>	-35870	150	2502.5	0.4	> 100 ns		1993	IT=100
<sup>153</sup> Lu <sup>p</sup>	-35740	150	2632.9	0.5	15 $\mu$ s	3	1993	IT=100
<sup>153</sup> Hf	-27300#	300#			400# ms	(>200 ns)	2000	$\beta^+ ?$
<sup>153</sup> Hf <sup>m</sup>	-26550#	320#	750#	100#	500# ms			$\beta^+ ?; IT ?$
<sup>153</sup> Nd <sup>m</sup>	T : average 10Si03=1.17(0.07) 96Ya12=1.06(0.05)							
<sup>153</sup> Er	J : and 89Ot.A							
<sup>153</sup> Yb <sup>m</sup>	E : in ENSDF 2578.2 + x							
<sup>153</sup> Lu	D : p=0% decay is from 97Ir01							
<sup>154</sup> Ba	-32820#	500#			53 ms	48	2016	$\beta^-=100$
<sup>154</sup> La	-41530#	300#			161 ms	15	2016	$\beta^-=100; \beta^-n=20\#; \beta^-2n=0.1\#$
<sup>154</sup> Ce	-52220#	200#			722 ms	14	1994	$\beta^-=100; \beta^-n=0.1\#$
<sup>154</sup> Pr	-58100	110			2.3 s	0.1	1988	$\beta^-=100; \beta^-n=0.2\#$
<sup>154</sup> Nd	-65820	50			25.9 s	0.2	1970	$\beta^-=100$
<sup>154</sup> Nd <sup>m</sup>	-64520	50	1297.9	0.4	3.2 $\mu$ s	0.3	1970	IT=100
<sup>154</sup> Pm	-68510	50			2.68 m	0.07	1958	$\beta^-=100$
<sup>154</sup> Pm <sup>m</sup>	-68490	40	20	12	1.73 m	0.10	1958	$\beta^-=100$
<sup>154</sup> Sm	-72455.2	1.5			STABLE	(>2.3 Ey)	1933	IS=22.75 29; $2\beta^- ?$
<sup>154</sup> Eu	-71738.1	1.3			8.601 y	0.010	1947	$\beta^-\approx 100; \epsilon=0.018$ 12
<sup>154</sup> Eu <sup>m</sup>	-71669.9	1.3	68.1702	0.0004	2.2 $\mu$ s	0.1	1964	IT=100
<sup>154</sup> Eu <sup>n</sup>	-71592.8	1.3	145.3	0.3	46.3 m	0.4	1975	IT=100
<sup>154</sup> Gd	-73706.0	1.2			STABLE		1938	IS=2.18 3
<sup>154</sup> Tb	-70160	50			21.5 h	0.4	1950	$\beta^+\approx 100; \beta^- < 0.1$
<sup>154</sup> Tb <sup>m</sup>	-70150	50	12	7	9.994 h	0.039	1972	$\beta^+=78.2$ 7; IT=21.8 7; $\beta^- < 0.1$
<sup>154</sup> Tb <sup>n</sup>	-69960#	160#	200#	150#	22.7 h	0.5	1972	$\beta^+=98.2$ 6; IT=1.8 6
<sup>154</sup> Tb <sup>p</sup>	-62160#	900#	8000#	900#	513 ns	42	1982	IT ?
<sup>154</sup> Dy	-70394	7			3.0 My	1.5	1961	$\alpha=100; 2\beta^+ ?$
<sup>154</sup> Ho	-64639	8			11.76 m	0.19	1966	$\beta^+\approx 100; \alpha=0.019$ 5
<sup>154</sup> Ho <sup>m</sup>	-64397	27	243	28	3.10 m	0.14	1968	$\beta^+=100; \alpha < 0.001; IT\approx 0$
<sup>154</sup> Er	-62605	5			3.73 m	0.09	1963	$\beta^+\approx 100; \alpha=0.47$ 13
<sup>154</sup> Tm	-54427	14			8.1 s	0.3	1964	$\alpha=54$ 5; $\beta^+=46$ 5
<sup>154</sup> Tm <sup>m</sup>	-54350	50	70	50	3.30 s	0.07	1964	$\alpha=58$ 5; $\beta^+=42$ 5; IT ?
<sup>154</sup> Yb	-49932	17			409 ms	2	1964	$\alpha=92.6$ 12; $\beta^+=7.4$ 12
<sup>154</sup> Lu	-39720#	200#			1#	s	1981	$\beta^+ ?$
<sup>154</sup> Lu <sup>m</sup>	-39660#	200#	60	12	1.12 s	0.08	1981	$\beta^+\approx 100; \beta^+p=?; \beta^+\alpha=?; \alpha=0.002$ #
<sup>154</sup> Lu <sup>n</sup>	-37000#	220#	2720#	100#	35 $\mu$ s	3	1990	IT=100
<sup>154</sup> Hf	-32670#	300#			2 s	1	1981	$\beta^+\approx 100; \alpha\approx 0$
<sup>154</sup> Hf <sup>m</sup>	-29960#	300#	2710#	30#	9 $\mu$ s	4	1989	IT=100
<sup>154</sup> Nd <sup>m</sup>	E : from a least-squares fit to $\gamma$ -ray energies in 09Si21							
<sup>154</sup> Tb <sup>m</sup>	E : estimated by NUBASE from 73Ba20<25 keV							
<sup>154</sup> Tm <sup>m</sup>	D : IT decay has not been observed							
<sup>154</sup> Lu <sup>m</sup>	D : $\beta^+p$ and $\beta^+\alpha$ modes observed in 88Vi02; $\beta^+p$ confirmed in 90Sh.A							
<sup>154</sup> Lu <sup>n</sup>	E : 2431.3 + 130.4 + z, above <sup>154</sup> Lu <sup>m</sup> ; z estimated 100#100							
<sup>154</sup> Hf <sup>m</sup>	E : 42#28 above 2671 level, see ENSDF'09							
<sup>155</sup> La	-37930#	400#			101 ms	28	2016	$\beta^-=100; \beta^-n=60\#; \beta^-2n=0\#$
<sup>155</sup> Ce	-47780#	300#			313 ms	7	1994	$\beta^-=100; \beta^-n=0.2\#$
<sup>155</sup> Pr	-55415	17			1.47 s	0.3	1992	$\beta^-=100; \beta^-n=0.3\#$
<sup>155</sup> Nd	-62284	9			8.9 s	0.2	1986	$\beta^-=100$
<sup>155</sup> Pm	-66940	5			41.5 s	0.2	1982	$\beta^-=100$
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...											
<sup>155</sup> Sm	-70190.8	1.5			22.3 m	0.2	3/2 <sup>-</sup>	05		1951	$\beta^-$ =100
<sup>155</sup> Sm <sup>m</sup>	-70174.3	1.6	16.5	0.5	2.8 $\mu$ s	0.5	5/2 <sup>+</sup>		10Si03 ETJ	2010	IT=100
<sup>155</sup> Sm <sup>n</sup>	-69652.2	1.7	538.6	0.7	1.00 $\mu$ s	0.08	11/2 <sup>-</sup>		10Si03 ETJ	2010	IT=100
<sup>155</sup> Eu	-71818.1	1.4			4.741 y	0.009	5/2 <sup>+</sup>	05	14Un01 T	1947	$\beta^-$ =100 *
<sup>155</sup> Gd	-72069.9	1.2			STABLE		3/2 <sup>-</sup>	05		1933	IS=14.80 12
<sup>155</sup> Gd <sup>m</sup>	-71948.8	1.2	121.05	0.19	31.97 ms	0.27	11/2 <sup>-</sup>	05		1967	IT=100
<sup>155</sup> Tb	-71250	10			5.32 d	0.06	3/2 <sup>+</sup>	05		1957	$\epsilon$ =100
<sup>155</sup> Dy	-69156	10			9.9 h	0.2	3/2 <sup>-</sup>	05		1958	$\beta^+$ =100
<sup>155</sup> Dy <sup>m</sup>	-68922	10	234.33	0.03	6 $\mu$ s	1	(11/2 <sup>-</sup> )	05		1970	IT=100
<sup>155</sup> Ho	-66040	17			48 m	1	5/2 <sup>+</sup>	05		1959	$\beta^+$ =100
<sup>155</sup> Ho <sup>m</sup>	-65898	17	141.97	0.11	880 $\mu$ s	80	(11/2 <sup>-</sup> )	05		1984	IT=100
<sup>155</sup> Er	-62209	6			5.3 m	0.3	(7/2 <sup>-</sup> )	05		1969	$\beta^+$ ≈100; $\alpha$ =0.022 7
<sup>155</sup> Tm	-56626	10			21.6 s	0.2	(11/2 <sup>-</sup> )	05		1971	$\beta^+$ =99.11 24; $\alpha$ =0.89 24
<sup>155</sup> Tm <sup>m</sup>	-56585	12	41	6	45 s	3	1/2 <sup>+</sup>	05	FGK12a J	1990	$\beta^+$ >92; $\alpha$ <8 *
<sup>155</sup> Yb	-50503	17			1.793 s	0.019	(7/2 <sup>-</sup> )	05		1964	$\alpha$ =89 4; $\beta^+$ =11 4
<sup>155</sup> Lu	-42545	19			68.6 ms	1.6	(11/2 <sup>-</sup> )	05		1965	$\alpha$ =90 2; $\beta^+$ ?
<sup>155</sup> Lu <sup>m</sup>	-42524	20	21	4	AD		(1/2 <sup>+</sup> )	05		1967	$\alpha$ =76 16; $\beta^+$ ?
<sup>155</sup> Lu <sup>n</sup>	-40764	19	1781.0	2.0	AD		25/2 <sup>-</sup> #	05		1981	$\alpha$ ≈100; IT ?
<sup>155</sup> Hf	-34170#	300#			840 ms	30	7/2 <sup>-</sup> #	05	11Sa59 T	1981	$\beta^+$ ≈100; $\alpha$ ?
<sup>155</sup> Ta	-23930#	300#			3.2 ms	1.3	(11/2 <sup>-</sup> )	07		2007	p=100 *
* <sup>155</sup> Eu	T : average (values in days) 14Un01=1730.1(3.5) 98Si12=1739(8) **										
* <sup>155</sup> Tm <sup>m</sup>	J : favored $\alpha$ decay from <sup>159</sup> Lu 1/2 <sup>+</sup> **										
* <sup>155</sup> Ta	T : symmetrized from 2.9(+1.5-1.1) I : NUBASE expects 1/2 <sup>+</sup> 30#20 below **										
<sup>156</sup> La	-33050#	400#			84 ms	78			16Wu.A TD	2016	$\beta^-$ =100; $\beta^-$ -n ?
<sup>156</sup> Ce	-44820#	300#			233 ms	9	0 <sup>+</sup>		16Wu.A TD	2016	$\beta^-$ =100; $\beta^-$ -n=1#
<sup>156</sup> Pr	-51570#	200#			444 ms	6			16Wu.A TD	1992	$\beta^-$ =100; $\beta^-$ -n=0.7#
<sup>156</sup> Nd	-60470	200			5.06 s	0.13	0 <sup>+</sup>	12	07Sh05 T	1987	$\beta^-$ =100 *
<sup>156</sup> Nd <sup>m</sup>	-59040	200	1431.3	0.4	365 ns	145	(5 <sup>-</sup> )	12	09Si21 ET	1998	IT=100
<sup>156</sup> Pm	-64164	4			27.2 s	0.50	4 <sup>+</sup>	12	16Ko.A TJ	1986	$\beta^-$ =100 *
<sup>156</sup> Pm <sup>m</sup>	-64014	4	150.3	0.1	5.6 s	0.6	1 <sup>+</sup>	12	16Ko.A TJD	2007	IT≈98; $\beta^-$ =2#
<sup>156</sup> Sm	-69360	9			9.4 h	0.2	0 <sup>+</sup>	12		1951	$\beta^-$ =100
<sup>156</sup> Sm <sup>m</sup>	-67962	9	1397.55	0.09	185 ns	7	5 <sup>-</sup>	12		1974	IT=100
<sup>156</sup> Eu	-70083	4			15.19 d	0.08	0 <sup>+</sup>	12		1947	$\beta^-$ =100
<sup>156</sup> Gd	-72534.9	1.2			STABLE		0 <sup>+</sup>	12		1933	IS=20.47 9
<sup>156</sup> Gd <sup>m</sup>	-70397.3	1.2	2137.60	0.05	1.3 $\mu$ s	0.1	7 <sup>-</sup>	12		1969	IT=100
<sup>156</sup> Tb	-70091	4			5.35 d	0.10	3 <sup>-</sup>	12		1950	$\beta^+$ ≈100; $\beta^-$ ?
<sup>156</sup> Tb <sup>m</sup>	-70037	5	54	3	24.4 h	1.0	(7 <sup>-</sup> )	12		1970	IT=100 *
<sup>156</sup> Tb <sup>n</sup>	-70003	4	88.4	0.2	5.3 h	0.2	(0 <sup>+</sup> )	12		1950	IT=?; $\beta^+$ =?
<sup>156</sup> Dy	-70529.0	1.2			STABLE	(>1 Ey)	0 <sup>+</sup>	12	58Ri23 T	1948	IS=0.056 3; $\alpha$ ?; 2 $\beta^+$ ? *
<sup>156</sup> Ho	-65480	60			56 m	1	4 <sup>-</sup>	12		1957	$\beta^+$ =100
<sup>156</sup> Ho <sup>m</sup>	-65430	60	52.37	0.30	9.5 s	1.5	1 <sup>-</sup>	12		1995	IT≈100; $\beta^+$ ? *
<sup>156</sup> Ho <sup>n</sup>	-65304	28	170	70	MD		(9 <sup>+</sup> )	12		1975	$\beta^+$ =75; IT ? *
<sup>156</sup> Er	-64212	25			19.5 m	1.0	0 <sup>+</sup>	12		1967	$\beta^+$ =100; $\alpha$ =17e-6 4
<sup>156</sup> Tm	-56835	14			83.8 s	1.8	2 <sup>-</sup>	12		1971	$\beta^+$ ≈100; $\alpha$ =0.064 10
<sup>156</sup> Tm <sup>m</sup>	-56440#	200#	400#	200#	400 ns		(11 <sup>-</sup> )	12		1985	IT=100 *
<sup>156</sup> Tm <sup>n</sup>			non existent		19 s	3	9 <sup>+</sup>		91To08 I		* *
<sup>156</sup> Yb	-53266	9			26.1 s	0.7	0 <sup>+</sup>	12		1970	$\beta^+$ =90 2; $\alpha$ =10 2
<sup>156</sup> Lu	-43700	50			494 ms	12	(2 <sup>-</sup> )	12		1965	$\alpha$ =?; $\beta^+$ =5#
<sup>156</sup> Lu <sup>m</sup>	-43680	240	20	250	198 ms	2	(9 <sup>+</sup> )	12	96Pa01 D	1979	$\alpha$ =94 6; $\beta^+$ ? *
<sup>156</sup> Hf	-37820	150			23 ms	1	0 <sup>+</sup>	12	96Pa01 D	1979	$\alpha$ =97 3; $\beta^+$ ? *
<sup>156</sup> Hf <sup>m</sup>	-35860	150	1959.0	1.0	AD		8 <sup>+</sup>	12	96Pa01 T	1979	$\alpha$ =100 *
<sup>156</sup> Ta	-25860#	300#			106 ms	4	(2 <sup>-</sup> )	12		1992	p=71 3; $\beta^+$ =29 3
<sup>156</sup> Ta <sup>m</sup>	-25770#	300#	94	8	AD		(9 <sup>+</sup> )	12		1993	$\beta^+$ =95.8 9; p=4.2 9
* <sup>156</sup> Nd	T : others 89Ok.A=5.51(0.10) 87Gr12=5.47(0.11), see discussion in 07Sh05 **										
* <sup>156</sup> Pm	T : unweighed average 16Ko.A=27.78(0.07) 87Gr12=26.70(0.10) **										
* <sup>156</sup> Tb <sup>m</sup>	E : derived from E3 24h to 4 <sup>+</sup> 49.630 level and E(IT)<B(L)=9 keV **										
* <sup>156</sup> Dy	T : lower limit is for $\alpha$ decay **										
* <sup>156</sup> Ho <sup>m</sup>	E : Error not given, estimated by NUBASE **										
* <sup>156</sup> Tm <sup>m</sup>	E : 203.6 keV above unknown level **										
* <sup>156</sup> Tm <sup>n</sup>	I : see also the discussion in ENSDF'03 **										
* <sup>156</sup> Lu <sup>m</sup>	D : derived from original $\alpha$ =98(9)% **										
* <sup>156</sup> Hf	D : derived from original $\alpha$ =100(6)% **										
* <sup>156</sup> Hf <sup>m</sup>	T : average 96Pa01=520(10) 81Ho.A=444(17) **										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>157</sup> Ce	-39930# 400#			175 ms 41	7/2 <sup>+</sup> #		16Wu.A TD	2016	$\beta^- = 100; \beta^- n = 2\#$
<sup>157</sup> Pr	-48540# 300#			307 ms 21	5/2 <sup>-</sup> #		16Wu.A TD	2016	$\beta^- = 100; \beta^- n = 6\#$
<sup>157</sup> Nd	-56462 25			1.15 s 0.03	5/2 <sup>-</sup> #	16	16Wu.A TD	1992	$\beta^- = 100$
<sup>157</sup> Pm	-62297 7			10.56 s 0.10	(5/2 <sup>-</sup> )	16		1987	$\beta^- = 100$
<sup>157</sup> Sm	-66678 4			8.03 m 0.07	3/2 <sup>-</sup> #	16		1973	$\beta^- = 100$
<sup>157</sup> Eu	-69459 4			15.18 h 0.03	5/2 <sup>+</sup>	16		1951	$\beta^- = 100$
<sup>157</sup> Gd	-70823.5 1.2			STABLE	3/2 <sup>-</sup>	16		1933	IS=15.65 2
<sup>157</sup> Gd <sup>m</sup>	-70759.6 1.2	63.916	0.005	460 ns 40	5/2 <sup>+</sup>	16		1964	IT=100
<sup>157</sup> Gd <sup>n</sup>	-70397.0 1.2	426.539	0.023	18.5 $\mu$ s 2.3	11/2 <sup>-</sup>	16		1961	IT=100
<sup>157</sup> Tb	-70763.4 1.2			71 y 7	3/2 <sup>+</sup>	16		1960	$\epsilon = 100$
<sup>157</sup> Dy	-69425 5			8.14 h 0.04	3/2 <sup>-</sup>	16		1953	$\beta^+ = 100$
<sup>157</sup> Dy <sup>m</sup>	-69263 5	161.99	0.03	1.3 $\mu$ s 0.2	9/2 <sup>+</sup>	16		1974	IT=100
<sup>157</sup> Dy <sup>n</sup>	-69226 5	199.38	0.07	21.6 ms 1.6	11/2 <sup>-</sup>	16		1970	IT=100
<sup>157</sup> Ho	-66833 23			12.6 m 0.2	7/2 <sup>-</sup>	16		1966	$\beta^+ = 100$
<sup>157</sup> Er	-63414 27			18.65 m 0.10	3/2 <sup>-</sup>	16		1966	$\beta^+ = 100$
<sup>157</sup> Er <sup>m</sup>	-63259 27	155.4	0.3	76 ms 6	9/2 <sup>+</sup>	16		1971	IT=100
<sup>157</sup> Tm	-58709 28			3.63 m 0.09	1/2 <sup>+</sup>	16		1974	$\beta^+ = 100$
<sup>157</sup> Yb	-53422 11			38.6 s 1.0	7/2 <sup>-</sup>	16		1970	$\beta^+ = 99.5; \alpha = 0.5$
<sup>157</sup> Lu	-46441 12			6.8 s 1.8	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	16		1977	$\beta^+ ?; \alpha = ?$
<sup>157</sup> Lu <sup>m</sup>	-46420 12	20.9	2.0	AD	(11/2 <sup>-</sup> )	16		1972	$\beta^+ = ?; \alpha = 6 2$
<sup>157</sup> Hf	-38900# 200#			115 ms 1	(7/2 <sup>-</sup> )	16		1965	$\alpha = 94 4; \beta^+ = 14 9$
<sup>157</sup> Ta	-29590 150			10.1 ms 0.4	1/2 <sup>+</sup>	16		1979	$\alpha = ?; p = 3.4 12; \beta^+ = 1\#$
<sup>157</sup> Ta <sup>m</sup>	-29570 150	22	5	AD	4.3 ms 0.1	11/2 <sup>-</sup>		1996	$\alpha = ?; \beta^+ = 1\#; p = 0$
<sup>157</sup> Ta <sup>n</sup>	-28000 150	1593	9	AD	1.7 ms 0.1	25/2 <sup>-</sup> #		1996	$\alpha = 100$
<sup>157</sup> W	-19470# 400#			275 ms 40	(7/2 <sup>-</sup> )	16	10Bi03 D	2010	$\beta^+ = 100; \alpha = 0$
<sup>157</sup> W <sup>p</sup>	-19150# 400#	320	30	AD	(9/2 <sup>-</sup> )	16		2010	IT ?
* <sup>157</sup> Pr	T : symmetrized from 295(+29-11)								
* <sup>157</sup> Lu	T : ENSDF'16 average of conflicting 91To09=5.7(0.5) 91Le15=92Po14=9.6(0.8)								
<sup>158</sup> Ce	-36660# 400#			99 ms 93	0 <sup>+</sup>		16Wu.A TD	2016	$\beta^- = 100; \beta^- n ?$
<sup>158</sup> Pr	-44330# 300#			181 ms 14			16Wu.A TD	2016	$\beta^- = 100; \beta^- n = 10\#$
<sup>158</sup> Nd	-54060# 200#			810 ms 30	0 <sup>+</sup>	13	16Wu.A TD	1992	$\beta^- = 100$
<sup>158</sup> Pm	-59089 13			4 s 0.5		04		1987	$\beta^- = 100$
<sup>158</sup> Pm <sup>m</sup>	-58940# 50#	150#	50#	> 16 $\mu$ s			15YoZX TD	2015	IT=100
<sup>158</sup> Sm	-65250 5			5.30 m 0.03	0 <sup>+</sup>	04		1970	$\beta^- = 100$
<sup>158</sup> Sm <sup>m</sup>	-63971 5	1279.1	1.8	115 ns 18	(5 <sup>-</sup> )	04		1973	IT=100
<sup>158</sup> Eu	-67255 10			45.9 m 0.2	(1 <sup>-</sup> )	04		1951	$\beta^- = 100$
<sup>158</sup> Gd	-70689.5 1.2			STABLE	0 <sup>+</sup>	04		1933	IS=24.84 7
<sup>158</sup> Tb	-69470.7 1.4			180 y 11	3 <sup>-</sup>	04		1957	$\beta^+ = 83.4 7; \beta^- = 16.6 7$
<sup>158</sup> Tb <sup>m</sup>	-69360.4 1.8	110.3	1.2	10.70 s 0.17	0 <sup>-</sup>	04		1957	IT $\approx$ 100; $\beta^- < 0.6; \beta^+ < 0.01$
<sup>158</sup> Tb <sup>n</sup>	-69082.3 2.3	388.4	1.8	400 $\mu$ s 40	(7 <sup>-</sup> )	04		1961	IT=100
<sup>158</sup> Dy	-70407.3 2.4			STABLE	0 <sup>+</sup>	04		1938	IS=0.095 3; $\alpha ?; 2\beta^+ ?$
<sup>158</sup> Ho	-66188 27			11.3 m 0.4	5 <sup>+</sup>	04		1961	$\beta^+ \approx 100; \alpha ?$
<sup>158</sup> Ho <sup>m</sup>	-66121 27	67.199	0.010	28 m 2	2 <sup>-</sup>	04		1960	IT>81; $\beta^+ < 19$
<sup>158</sup> Ho <sup>n</sup>	-66010# 80#	180#	70#	21.3 m 2.3	(9 <sup>+</sup> )	04		1970	$\beta^+ > 93; IT < 7\#$
<sup>158</sup> Er	-65304 25			2.29 h 0.06	0 <sup>+</sup>	07		1961	$\epsilon = 100$
<sup>158</sup> Tm	-58703 25			3.98 m 0.06	2 <sup>-</sup>	04		1970	$\beta^+ = 100$
<sup>158</sup> Tm <sup>m</sup>	-58650# 100#	50#	100#	*	20 ns	(5 <sup>+</sup> )	04	81Dr07 T	1981
<sup>158</sup> Yb	-56010 8			1.49 m 0.13	0 <sup>+</sup>	04		1967	$\beta^+ \approx 100; \alpha \approx 0.0021 12$
<sup>158</sup> Lu	-47212 15			10.6 s 0.3	2 <sup>-</sup>	04	95Ga.A J	1979	$\beta^+ = 99.09 20; \alpha = 0.91 20$
<sup>158</sup> Hf	-42102 17			0.99 s 0.03	0 <sup>+</sup>	04	15Li24 T	1965	$\beta^+ = 55.7 19; \alpha = 44.3 19$
<sup>158</sup> Ta	-31170# 200#			49 ms 8	(2 <sup>-</sup> )	04	97Da07 TD	1979	$\alpha = 96 4; \beta^+ ?$
<sup>158</sup> Ta <sup>m</sup>	-31030# 200#	141	11	AD	36.0 ms 0.8	(9 <sup>+</sup> )	04	97Da07 ETJ	1979
<sup>158</sup> Ta <sup>n</sup>	-28360# 200#	2805	16	AD	6.1 $\mu$ s 0.1	(19 <sup>-</sup> )	14Ca03 TJD	2014	IT=98.6 2; $\alpha = 1.4 2$
<sup>158</sup> W	-23630# 300#			1.25 ms 0.21	0 <sup>+</sup>	06		1981	$\alpha = 100$
<sup>158</sup> W <sup>m</sup>	-21740# 300#	1889	8	AD	143 $\mu$ s 19	(8 <sup>+</sup> )	06	1995	$\alpha = 100$
* <sup>158</sup> Nd	T : symmetrized from 820(+15-36)								
* <sup>158</sup> Tm <sup>m</sup>	I : T $\approx$ 20 s in 81Dr07 was a typo. Value in Fig. 2 was correct. See 96Dr.A								
* <sup>158</sup> Ta	T : average 97Da07=72(12) 96Pa01=46(4) with Birge ratio B=2								
* <sup>158</sup> Ta	D : derived from original $\alpha \approx 100(8)\%$								
* <sup>158</sup> Ta <sup>m</sup>	T : average 97Da07=37.7(1.5) 96Pa01=35(1) 79Ho10=36.8(1.6)								
* <sup>158</sup> Ta <sup>n</sup>	E : 14Ca03=2668 above 9 <sup>+</sup> isomer								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>159</sup> Pr	-41090# 400#		134 ms 43	5/2 <sup>-</sup> #		16Wu.A TD	2016	$\beta^-$ =100; $\beta^-$ n=30#
<sup>159</sup> Nd	-49810# 300#		500 ms 30	7/2 <sup>+</sup> #	13	16Wu.A TD	2012	$\beta^-$ =100; $\beta^-$ n=0.02# *
<sup>159</sup> Pm	-56554 10		1.49 s 0.13	5/2 <sup>-</sup> #	12	16Wu.A T	1998	$\beta^-$ =100 *
<sup>159</sup> Pm <sup>m</sup>	-55089 10	1465.0 0.5	4.42 $\mu$ s 0.17			15YoZX ETD	2015	IT=100
<sup>159</sup> Sm	-62208 6		11.37 s 0.15	5/2 <sup>-</sup>	12		1986	$\beta^-$ =100
<sup>159</sup> Sm <sup>m</sup>	-60931 6	1276.8 0.5	116 ns 8	(11/2 <sup>-</sup> )	12	09Ur04 ET	2009	IT=100
<sup>159</sup> Eu	-66043 4		18.1 m 0.1	5/2 <sup>+</sup>	12		1961	$\beta^-$ =100
<sup>159</sup> Gd	-68561.4 1.2		18.479 h 0.004	3/2 <sup>-</sup>	12		1949	$\beta^-$ =100
<sup>159</sup> Tb	-69532.4 1.3		STABLE	3/2 <sup>+</sup>	12	12Vi10 J	1933	IS=100. *
<sup>159</sup> Dy	-69167.1 1.5		144.4 d 0.2	3/2 <sup>-</sup>	12		1951	$\epsilon$ =100
<sup>159</sup> Dy <sup>m</sup>	-68814.3 1.5	352.77 0.14	122 $\mu$ s 3	11/2 <sup>-</sup>	12		1965	IT=100
<sup>159</sup> Ho	-67330 3		33.05 m 0.11	7/2 <sup>-</sup>	12		1958	$\beta^+$ =100
<sup>159</sup> Ho <sup>m</sup>	-67124 3	205.91 0.05	8.30 s 0.08	1/2 <sup>+</sup>	12		1966	IT=100
<sup>159</sup> Er	-64561 4		36 m 1	3/2 <sup>-</sup>	12		1962	$\beta^+$ =100
<sup>159</sup> Er <sup>m</sup>	-64378 4	182.602 0.024	337 ns 14	9/2 <sup>+</sup>	12		1971	IT=100
<sup>159</sup> Er <sup>m</sup>	-64132 4	429.05 0.03	590 ns 60	11/2 <sup>-</sup>	12		1971	IT=100
<sup>159</sup> Tm	-60570 28		9.13 m 0.16	5/2 <sup>+</sup>	12		1971	$\beta^+$ =100
<sup>159</sup> Yb	-55839 18		1.67 m 0.09	5/2 <sup>(-)</sup>	12		1975	$\beta^+$ =100
<sup>159</sup> Lu	-49710 40		12.1 s 1.0	1/2 <sup>+</sup>	12	FGK12a J	1980	$\beta^+$ $\approx$ 100; $\alpha$ =0.1# *
<sup>159</sup> Lu <sup>m</sup>	-49610# 90#	100# 80#	10# s	11/2 <sup>-</sup> #				$\beta^+$ ?; IT ?; $\alpha$ ? *
<sup>159</sup> Hf	-42853 17		5.20 s 0.10	7/2 <sup>-</sup>	12	96Pa01 T	1973	$\beta^+$ =65 7; $\alpha$ =35 7
<sup>159</sup> Ta	-34439 20		1.04 s 0.09	1/2 <sup>+</sup>	12	97Da07 T	1979	$\beta^+$ ?; $\alpha$ =34 5 *
<sup>159</sup> Ta <sup>m</sup>	-34375 19	64 5	560 ms 60	11/2 <sup>-</sup>	12		1994	$\alpha$ =55 1; $\beta^+$ ?
<sup>159</sup> W	-25300# 300#		8.2 ms 0.7	7/2 <sup>-</sup> #	12	96Pa01 TD	1981	$\alpha$ =82 16; $\beta^+$ ? *
<sup>159</sup> Re	-14750# 310#		40# $\mu$ s	1/2 <sup>+</sup> #			2006	p ?; $\alpha$ ?
<sup>159</sup> Re <sup>m</sup>	-14540# 300#	210# 50#	21.6 $\mu$ s 3.3	11/2 <sup>-</sup>	12	07Pa27 T	2006	p=?; $\alpha$ =7.5 35 *
* <sup>159</sup> Nd	T : symmetrized from 485(+39-20)							
* <sup>159</sup> Pm	T : average 16Wu.A=1.48(0.18) 05Ic02=1.5(0.2)							
* <sup>159</sup> Tb	J : 3/2 confirmed by a novel technique in 12Vi10 (see text)							
* <sup>159</sup> Lu	J : favored $\alpha$ decay from <sup>163</sup> Ta 1/2 <sup>+</sup>							
* <sup>159</sup> Ta	T : average 97Da07=0.83(0.18) 96Pa01=1.10(0.10)							
* <sup>159</sup> W	D : derived from original $\alpha$ =92(23)%							
* <sup>159</sup> Re <sup>m</sup>	T : average 07Pa27=23(6) 06Jo10=21(4)							
<sup>160</sup> Pr	-36520# 400#		170 ms 140			16Wu.A TD	2016	$\beta^-$ =100
<sup>160</sup> Nd	-47130# 300#		439 ms 37	0 <sup>+</sup>	13	16Wu.A TD	1985	$\beta^-$ =100; $\beta^-$ n=0.08# *
<sup>160</sup> Pm	-53000# 200#		725 ms 57	0 <sup>-</sup> #	13	16Wu.A TD	2012	$\beta^-$ =100; $\beta^-$ n=0.03# *
<sup>160</sup> Sm	-60235 6		9.6 s 0.3	0 <sup>+</sup>	05		1986	$\beta^-$ =100
<sup>160</sup> Sm <sup>m</sup>	-58874 6	1361.3 0.4	120 ns 46	(5 <sup>-</sup> )		09Si21 ETJ	2009	IT=100
<sup>160</sup> Sm <sup>n</sup>	-57478 6	2757.3 0.4	1.8 $\mu$ s 0.4	(11 <sup>+</sup> )		16Pa01 ETJ	2016	IT=100
<sup>160</sup> Eu	-63480 10		42.4 s 0.2	(5 <sup>-</sup> )	05	16Ha.A TJ	1973	$\beta^-$ =100
<sup>160</sup> Eu <sup>m</sup>	-63400 12	80 7	29.9 s 0.3	(1 <sup>-</sup> )	05	16Ha.A ETJ	2016	$\beta^-$ =100
<sup>160</sup> Gd	-67941.7 1.3		STABLE	0 <sup>+</sup>	05	01Da22 T	1933	IS=21.86 19; 2 $\beta^-$ ?
<sup>160</sup> Tb	-67836.3 1.3		72.3 d 0.2	3 <sup>-</sup>	05		1943	$\beta^-$ =100
<sup>160</sup> Dy	-69672.7 0.8		STABLE	0 <sup>+</sup>	05		1938	IS=2.329 18
<sup>160</sup> Ho	-66383 15		25.6 m 0.3	5 <sup>+</sup>	05		1950	$\beta^+$ =100
<sup>160</sup> Ho <sup>m</sup>	-66323 15	59.98 0.03	5.02 h 0.05	2 <sup>-</sup>	05		1955	IT=73 3; $\beta^+$ =27 3
<sup>160</sup> Ho <sup>n</sup>	-66186 22	197 16	3 s	(9 <sup>+</sup> )	05	GAu E	1988	IT=100 *
<sup>160</sup> Er	-66064 24		28.58 h 0.09	0 <sup>+</sup>	05		1954	$\epsilon$ =100
<sup>160</sup> Tm	-60300 30		9.4 m 0.3	1 <sup>-</sup>	05		1970	$\beta^+$ =100
<sup>160</sup> Tm <sup>m</sup>	-60230 40	70 20	74.5 s 1.5	(5 <sup>+</sup> )	05		1983	IT=85 5; $\beta^+$ =15 5
<sup>160</sup> Tm <sup>n</sup>	-60200# 60#	100# 50#	200 ns	(8)	05		1986	IT=100 *
<sup>160</sup> Yb	-58163 7		4.8 m 0.2	0 <sup>+</sup>	05		1967	$\beta^+$ =100
<sup>160</sup> Lu	-50270 60		36.1 s 0.3	2 <sup>-</sup> #	05		1979	$\beta^+$ =100; $\alpha$ <1e-4
<sup>160</sup> Lu <sup>m</sup>	-50270# 120#	0# 100#	40 s 1		05		1980	$\beta^+$ $\approx$ 100; $\alpha$ ?
<sup>160</sup> Hf	-45939 10		13.6 s 0.2	0 <sup>+</sup>	05		1973	$\beta^+$ =99.3 2; $\alpha$ =0.7 2
<sup>160</sup> Ta	-35820 50		1.70 s 0.20	(2 <sup>-</sup> )	05	96Pa01 JD	1979	$\beta^+$ ?; $\alpha$ =?
<sup>160</sup> Ta <sup>m</sup>	-35710 240	110 250	1.55 s 0.04	(9 <sup>+</sup> )	05	96Pa01 TJ	1979	$\beta^+$ =66#; $\alpha$ =?
<sup>160</sup> W	-29330 150		90 ms 5	0 <sup>+</sup>	05	96Pa01 TD	1979	$\alpha$ =87 8; $\beta^+$ ? *
<sup>160</sup> Re	-16740# 300#		611 $\mu$ s 7	(4 <sup>-</sup> )	05	11Da12 TJD	1992	p=89 1; $\alpha$ =11 1 *
<sup>160</sup> Re <sup>m</sup>	-16560# 300#	182 16	2.8 $\mu$ s 0.1	(9 <sup>+</sup> )		11Da01 JT	2011	IT=100
* <sup>160</sup> Nd	I : first seen in 85Si25 in the thermal fission of <sup>252</sup> Cf; 12Ku26>300ns							
* <sup>160</sup> Ho <sup>n</sup>	E : less than 55 keV above 169.61 level, from ENSDF							
* <sup>160</sup> Tm <sup>n</sup>	E : 98.2 + x, x estimated 0#50							
* <sup>160</sup> Ta	J : from $\alpha$ correlation with <sup>156</sup> Lu line							
* <sup>160</sup> Ta <sup>m</sup>	J : from $\alpha$ correlation with <sup>156</sup> Lu <sup>m</sup> line							
* <sup>160</sup> W	T : average 96Pa01=91(5) 81Ho10=81(15)							
* <sup>160</sup> Re	J : protons from d <sub>3/2</sub> orbital; 92Pa05=(2 <sup>-</sup> )							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>161</sup> Nd	-42590#	400#	215 ms	76		16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ -n=0.6#
<sup>161</sup> Pm	-50240#	300#	1.05 s	0.15		16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ -n=0.4#
<sup>161</sup> Pm <sup>m</sup>	-49270#	300#	966.0	0.5	0.89 $\mu$ s	15YoZX	ETJ	2015	IT=100
<sup>161</sup> Sm	-56672	7	4.8 s	0.4				1998	$\beta^-$ =100
<sup>161</sup> Eu	-61792	10	26.2 s	2.3		16Wu.A	T	1986	$\beta^-$ =100
<sup>161</sup> Gd	-65505.8	1.6	3.646 m	0.003		94It.A	T	1949	$\beta^-$ =100
<sup>161</sup> Tb	-67461.6	1.4	6.89 d	0.02				1949	$\beta^-$ =100
<sup>161</sup> Dy	-68055.8	0.8	STABLE					1934	IS=18.889 42
<sup>161</sup> Dy <sup>m</sup>	-67570.2	0.8	485.56	0.16	760 ns	12Sw01	T	2012	IT=100
<sup>161</sup> Ho	-67197.3	2.2	2.48 h	0.05				1954	$\epsilon$ =100
<sup>161</sup> Ho <sup>m</sup>	-66986.2	2.2	211.15	0.03	6.76 s			1965	IT=100
<sup>161</sup> Er	-65202	9	3.21 h	0.03				1954	$\beta^+$ =100
<sup>161</sup> Er <sup>m</sup>	-64806	9	396.44	0.04	7.5 $\mu$ s			1969	IT=100
<sup>161</sup> Tm	-61899	28	30.2 m	0.8				1959	$\beta^+$ =100
<sup>161</sup> Tm <sup>m</sup>	-61891	28	7.51	0.24	5# m			1981	$\beta^+$ ?; IT ?
<sup>161</sup> Tm <sup>n</sup>	-61821	28	78.20	0.03	110 ns			1981	IT=100
<sup>161</sup> Yb	-57839	15	4.2 m	0.2				1974	$\beta^+$ =100
<sup>161</sup> Lu	-52562	28	77 s	2				1973	$\beta^+$ =100
<sup>161</sup> Lu <sup>m</sup>	-52388	28	174	4	7.3 ms			1973	IT=100
<sup>161</sup> Hf	-46315	23	18.4 s	0.4				1973	$\beta^+$ $\approx$ 100; $\alpha$ <0.13
<sup>161</sup> Hf <sup>m</sup>	-45986	23	329.0	0.5	4.8 $\mu$ s			2014	IT=100
<sup>161</sup> Ta	-38779	24	3# s					1979	$\beta^+$ ?; $\alpha$ ?
<sup>161</sup> Ta <sup>m</sup>	-38718	12	61	23	3.08 s	12Th13	D	1979	$\beta^+$ ?; $\alpha$ =7(3)
<sup>161</sup> W	-30560#	200#	409 ms	16		96Pa01	T	1973	$\alpha$ =73 3; $\beta^+$ =27 3
<sup>161</sup> Re	-20840	150	440 $\mu$ s	1		06La16	T	1979	$p$ $\approx$ 100; $\alpha$ <1.4
<sup>161</sup> Re <sup>m</sup>	-20720	150	123.7	1.3	14.7 ms			1979	$\alpha$ =93.0 3; $p$ =7.0 3
<sup>161</sup> Os	-9980#	400#	640 $\mu$ s	60				2010	$\alpha$ $\approx$ 100
* <sup>161</sup> Eu	T : average 16Wu.A=30.1(9.0) 90An31=24(4) 86Ma12=27(3)								**
* <sup>161</sup> Lu <sup>m</sup>	E : 166.5(0.8) keV above (3/2 <sup>+</sup> ) level at x<15 keV								**
* <sup>161</sup> W	T : average 96Pa01=409(18) 79Ho10=410(40)								**
<sup>162</sup> Nd	-39550#	400#	310 ms	200		16Wu.A	TD	2012	$\beta^-$ =100
<sup>162</sup> Pm	-46370#	300#	630 ms	180		16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ -n=0.8#
<sup>162</sup> Sm	-54530#	200#	2.7 s	0.3		16Wu.A	T	2005	$\beta^-$ =100
<sup>162</sup> Eu	-58700	40	11 s			16Wu.A	T	1987	$\beta^-$ =100
<sup>162</sup> Eu <sup>m</sup>	-58540	40	157	5	7.5 s	16Ko.A	ETJ	2016	$\beta^-$ =100
<sup>162</sup> Gd	-64280	4	8.4 m	0.2				1967	$\beta^-$ =100
<sup>162</sup> Tb	-65680	40	7.60 m	0.15				1965	$\beta^-$ =100
<sup>162</sup> Dy	-68181.5	0.8	STABLE					1934	IS=25.475 36
<sup>162</sup> Dy <sup>m</sup>	-65993.4	0.9	2188.1	0.3	8.3 $\mu$ s	11Sw02	ETD	2011	IT=100
<sup>162</sup> Ho	-66042	3	15.0 m	1.0				1957	$\beta^+$ =100
<sup>162</sup> Ho <sup>m</sup>	-65936	3	105.87	0.06	67.0 m			1961	IT=62; $\beta^+$ =38
<sup>162</sup> Er	-66334.5	0.8	STABLE		(>140 Ty)	56Po16	T	1938	IS=0.139 5; $\alpha$ ?; 2 $\beta^+$ ?
<sup>162</sup> Er <sup>m</sup>	-64308.5	0.8	2026.01	0.13	88 ns	12Sw01	TJ	1974	IT=100
<sup>162</sup> Tm	-61478	26	21.70 m	0.19				1963	$\beta^+$ =100
<sup>162</sup> Tm <sup>m</sup>	-61350	50	130	40	24.3 s	GAu	E	1974	IT ?; $\beta^+$ =19 4
<sup>162</sup> Yb	-59826	15	18.87 m	0.19				1963	$\beta^+$ =100
<sup>162</sup> Lu	-52830	80	*		1.37 m			1978	$\beta^+$ =100
<sup>162</sup> Lu <sup>m</sup>	-52710#	220#	120#	200#	*			1980	$\beta^+$ $\approx$ 100; IT ?
<sup>162</sup> Lu <sup>n</sup>	-52530#	220#	300#	200#	*			1980	$\beta^+$ $\approx$ 100; IT ?
<sup>162</sup> Hf	-49169	9	39.4 s	0.9				1982	$\beta^+$ $\approx$ 100; $\alpha$ =0.008 1
<sup>162</sup> Ta	-39780	50	3.57 s	0.12				1985	$\beta^+$ $\approx$ 100; $\alpha$ =0.074 10; $\beta^+$ p ?
<sup>162</sup> W	-33999	18	1.19 s	0.12				1973	$\beta^+$ ?; $\alpha$ =45.2 16
<sup>162</sup> Re	-22500#	200#	107 ms	13				1979	$\alpha$ =94 6; $\beta^+$ ?
<sup>162</sup> Re <sup>m</sup>	-22330#	200#	175	9	77 ms			1979	$\alpha$ =91 5; $\beta^+$ ?
<sup>162</sup> Os	-14440#	300#	2.1 ms	0.1				1989	$\alpha$ =100
* <sup>162</sup> Eu	T : 16Wu.A=11.8(1.4) 87Gr12=10.6(1.0) but values include both ground-state and isomer								**
* <sup>162</sup> Eu	J : from 16Ko.A, conf p5/2[413]n7/2[633],Kp=1 <sup>+</sup>								**
* <sup>162</sup> Er	T : lower limit is for $\alpha$ decay								**
* <sup>162</sup> Tm <sup>m</sup>	E : above 66.90 level and less than 192 keV, from ENSDF								**



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{163}\text{Pm}$	-43250# 400#		430 ms	350	$5/2^-$ #	13 16Wu.A TD	2012	$\beta^- = 100; \beta^- n = 1\#$
$^{163}\text{Sm}$	-50720# 300#		1.3 s	0.5	$1/2^-$ #	13 16Wu.A TD	2012	$\beta^- = 100$
$^{163}\text{Eu}$	-56480 70		7.7 s	0.4	$5/2^+$ #	10 08Os02 T	2007	$\beta^- = 100$
$^{163}\text{Gd}$	-61314 8		68 s	3	$7/2^+$ #	10	1982	$\beta^- = 100$
$^{163}\text{Gd}^m$	-61176 8	137.8	23.5 s	1.0	$1/2^-$ #	14Ha38 ETD	2014	$\beta^- = 100$
$^{163}\text{Tb}$	-64596 4		19.5 m	0.3	$3/2^+$	10	1966	$\beta^- = 100$
$^{163}\text{Dy}$	-66381.2 0.8		STABLE		$5/2^-$	10	1934	IS=24.896 42
$^{163}\text{Ho}$	-66378.3 0.8		4.570 ky	0.025	$7/2^-$	10	1957	$\epsilon = 100$
$^{163}\text{Ho}^m$	-66080.4 0.8	297.88	1.09 s	0.03	$1/2^+$	10	1957	IT=100
$^{163}\text{Ho}^n$	-64268.9 0.9	2109.4	800 ns	150	$(23/2^+)$	12Sw01 ETJ	2012	IT=100
$^{163}\text{Er}$	-65168 5		75.0 m	0.4	$5/2^-$	10	1953	$\beta^+ = 100$
$^{163}\text{Er}^m$	-64723 5	445.5	580 ns	100	$(11/2^-)$	10	1974	IT=100
$^{163}\text{Tm}$	-62729 6		1.810 h	0.005	$1/2^+$	10	1959	$\beta^+ = 100$
$^{163}\text{Tm}^m$	-62642 6	86.92	380 ns	30	$(7/2^-)$	10	1975	IT=100
$^{163}\text{Yb}$	-59299 15		11.05 m	0.35	$3/2^-$	10	1967	$\beta^+ = 100$
$^{163}\text{Lu}$	-54791 28		3.97 m	0.13	$1/2^{(+)}$	10	1979	$\beta^+ = 100$
$^{163}\text{Hf}$	-49264 25		40.0 s	0.6	$(5/2^-)$	15	1982	$\beta^+ = 100; \alpha < 0.0001$
$^{163}\text{Ta}$	-42530 40		10.6 s	1.8	$1/2^+$	10 FGK12a J	1985	$\beta^+ \approx 100; \alpha \approx 0.2$
$^{163}\text{Ta}^m$	-42390# 40#	140#	10# s		$(9/2^-)$	10 FGK12a J		$\beta^+ ?; \alpha ?; IT ?$
$^{163}\text{W}$	-34910 50		2.63 s	0.09	$7/2^-$	10	1973	$\beta^+ ?; \alpha = 14.2$
$^{163}\text{W}^m$	-34430 50	480.3	154 ns	3	$13/2^+$	10	2010	IT=100
$^{163}\text{Re}$	-26002 19		390 ms	70	$1/2^+$	10	1979	$\beta^+ ?; \alpha = 32.3$
$^{163}\text{Re}^m$	-25882 19	120	214 ms	5	$11/2^-$	10	1979	$\alpha = 66.4; \beta^+ ?$
$^{163}\text{Os}$	-16190# 300#		5.5 ms	0.6	$7/2^-$	10 13Dr06 J	1981	$\alpha \approx 100; \beta^+ ?$
* $^{163}\text{Sm}$	T : symmetrized from 16Wu.A=1.23(+0.51-0.47)							
* $^{163}\text{Ho}$	T : other: 92Ju01=47(+5-4) d for q=66 <sup>+</sup> (bare ion)							
* $^{163}\text{Ta}$	J : favored a-decay from 1/2 <sup>+</sup> isomer in $^{167}\text{Re}$							
* $^{163}\text{Ta}^m$	J : favored a-decay from (9/2 <sup>-</sup> ) ground-state in $^{167}\text{Re}$							
$^{164}\text{Pm}$	-38870# 400#		200# ms					$\beta^- ?; \beta^- n ?$
$^{164}\text{Sm}$	-48100# 300#		1.43 s	0.24	$0^+$	15 16Wu.A TD	2012	$\beta^- = 100; \beta^- n = 0\#$
$^{164}\text{Sm}^m$	-46620# 300#	1485.5	600 ns	140	$(6^-)$	15	2014	IT=100
$^{164}\text{Eu}$	-53380# 110#		4.15 s	0.19	$0^-$ #	08 16Wu.A T	2007	$\beta^- = 100$
$^{164}\text{Gd}$	-59770# 100#		45 s	3	$0^+$	06	1988	$\beta^- = 100$
$^{164}\text{Tb}$	-62080 100		3.0 m	0.1	$(5^+)$	01	1968	$\beta^- = 100$
$^{164}\text{Dy}$	-65968.0 0.8		STABLE		$0^+$	01	1934	IS=28.260 54
$^{164}\text{Ho}$	-64981.5 1.5		29 m	1	$1^+$	01	1938	$\epsilon = 60.5; \beta^- = 40.5$
$^{164}\text{Ho}^m$	-64841.7 1.5	139.77	36.4 m	0.3	$6^-$	01 08Ha21 T	1966	IT=100
$^{164}\text{Er}$	-65942.9 0.8		STABLE		$0^+$	01	1938	IS=1.601 3; $\alpha ?; 2\beta^+ ?$
$^{164}\text{Er}^m$	-62566.8 1.4	3376.1	68 ns	2	$(12^+)$	01 12Sw02 T	1980	IT=100
$^{164}\text{Tm}$	-61904 24		2.0 m	0.1	$1^+$	01	1960	$\epsilon = 61.1; e^+ = 39.1$
$^{164}\text{Tm}^m$	-61894 25	10	5.1 m	0.1	$6^-$	01 GAu E	1971	IT $\approx$ 80; $\beta^+ \approx 20$
$^{164}\text{Yb}$	-61017 15		75.8 m	1.7	$0^+$	01	1960	$\epsilon = 100$
$^{164}\text{Lu}$	-54642 28		3.14 m	0.03	$1^{(-)}$	07	1977	$\beta^+ = 100$
$^{164}\text{Hf}$	-51819 16		111 s	8	$0^+$	01	1981	$\beta^+ = 100$
$^{164}\text{Ta}$	-43283 28		14.2 s	0.3	$(3^+)$	08	1982	$\beta^+ = 100$
$^{164}\text{W}$	-38236 10		6.3 s	0.2	$0^+$	01	1973	$\beta^+ = 96.2.12; \alpha = 3.8.12$
$^{164}\text{Re}$	-27470 50		719 ms	161	$(2^-)$	01 09Ha42 TD	1979	$\alpha = ?; \beta^+ = 42\#$
* $^{164}\text{Re}^m$	-27520 240	-50	890 ms	130	$(9^+)$	01 09Ha42 TD	2009	$\beta^+ ?; \alpha = 3.1$
$^{164}\text{Os}$	-20420 150		21 ms	1	$0^+$	01	1981	$\alpha = ?; \beta^+ = 2\#$
$^{164}\text{Ir}$	-7340# 310#		1# ms		$2^-$ #	06		p ?; $\alpha ?; \beta^+ ?$
$^{164}\text{Ir}^m$	-7080# 300#	260#	70 $\mu\text{s}$	10	$(9^+)$	06 14Dr02 TD	2001	p=?; $\alpha = 4.2; \beta^+ ?$
* $^{164}\text{Eu}$	T : average 16Wu.A=3.80(0.56) 08Os02=4.2(0.2)							
* $^{164}\text{Ho}^m$	T : other 66Jo07=37.5(+1.5-0.5)							
* $^{164}\text{Er}^m$	T : ENSDF'2001 >170 ns							
* $^{164}\text{Tm}^m$	E : less than 20 keV, from ENSDF							
* $^{164}\text{Re}$	T : average 09Ha42=848(+140-105) 96Pa01=380(160) 81Ho10=880(240)							
* $^{164}\text{Re}^m$	T : symmetrized from 864(+150-110)							
$^{165}\text{Sm}$	-43810# 400#		980 ms	210	$5/2^-$ #	13 16Wu.A TD	2012	$\beta^- = 100; \beta^- n = 0.02\#$
$^{165}\text{Eu}$	-50720# 140#		2.53 s	0.25	$5/2^+$ #	08 16Wu.A T	2007	$\beta^- = 100; \beta^- n = 0.2\#$
$^{165}\text{Gd}$	-56450# 120#		11.0 s	0.9	$1/2^-$ #	06 16Wu.A T	1998	$\beta^- = 100$
$^{165}\text{Tb}$	-60570# 100#		2.11 m	0.10	$3/2^+$ #	06	1983	$\beta^- = 100$
$^{165}\text{Dy}$	-63612.6 0.8		2.334 h	0.001	$7/2^+$	06	1935	$\beta^- = 100$
$^{165}\text{Dy}^m$	-63504.4 0.8	108.1552	1.257 m	0.006	$1/2^-$	06	1963	IT=97.76 11; $\beta^- = 2.24.11$
$^{165}\text{Ho}$	-64899.0 1.0		STABLE		$7/2^-$	06	1934	IS=100.
$^{165}\text{Ho}^m$	-64537.3 1.0	361.675	1.512 $\mu\text{s}$	0.004	$3/2^+$	06	1958	IT=100
$^{165}\text{Ho}^n$	-64183.7 1.0	715.33	< 100 ns		$7/2^+$	06	1958	IT=100
$^{165}\text{Er}$	-64521.6 1.0		10.36 h	0.04	$5/2^-$	06	1950	$\epsilon = 100$
$^{165}\text{Er}^m$	-63970.3 1.2	551.3	250 ns	30	$11/2^-$	06	1970	IT=100
$^{165}\text{Er}^n$	-62698.6 1.2	1823.0	370 ns	40	$(19/2)$	12Sw01 EJT	2012	IT=100

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...											
$^{165}\text{Tm}$	-62929.6	1.7			30.06 h	0.03	1/2 <sup>+</sup>	06	1953	$\beta^+=100$	
$^{165}\text{Tm}^m$	-62849.2	1.7	80.37	0.06	80 $\mu\text{s}$	3	7/2 <sup>+</sup>	06	1967	IT=100	
$^{165}\text{Tm}^n$	-62769.1	1.7	160.47	0.06	9.0 $\mu\text{s}$	0.5	7/2 <sup>-</sup>	06	1968	IT=100	
$^{165}\text{Yb}$	-60295	27			9.9 m	0.3	5/2 <sup>-</sup>	06	1964	$\beta^+=100$	
$^{165}\text{Yb}^m$	-60168	27	126.80	0.09	300 ns	30	9/2 <sup>+</sup>	06	1980	IT=100	
$^{165}\text{Lu}$	-56442	27			10.74 m	0.10	1/2 <sup>+</sup>	06	1973	$\beta^+=100$	
$^{165}\text{Hf}$	-51636	28			76 s	4	(5/2 <sup>-</sup> )	06	1981	$\beta^+=100$	
$^{165}\text{Ta}$	-45848	14			31.0 s	1.5	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	06	FGK12a J	$\beta^+=100$	
$^{165}\text{Ta}^m$	-45823	17	24	18	30# s		(9/2 <sup>-</sup> )	06	FGK12a J	$\beta^+ ?; \alpha ?$	
$^{165}\text{W}$	-38861	25			5.1 s	0.5	(5/2 <sup>-</sup> )	06		$\beta^+\approx 100; \alpha < 0.2$	
$^{165}\text{Re}$	-30660	24			2.62 s	0.14	(1/2 <sup>+</sup> )	15	05Sc22 T	$\beta^+ ?; \alpha = 14.8$	
$^{165}\text{Re}^m$	-30632	12	27	22	1.74 s	0.06	(11/2 <sup>-</sup> )	15		$\beta^+ ?; \alpha = 13.1$	
$^{165}\text{Os}$	-21800#	200#			71 ms	3	(7/2 <sup>-</sup> )	14		$\alpha = 90.2; \beta^+ ?$	
$^{165}\text{Ir}$	-11590#	160#			50# ns	<1 $\mu\text{s}$	1/2 <sup>+</sup> #	06	97Da07 I	p ?; $\alpha ?$	
$^{165}\text{Ir}^m$	-11410	150	180#	50#	325 $\mu\text{s}$	33	(11/2 <sup>-</sup> )	06	14Dr02 TD	p=87.4; $\alpha = 12.2$	
* $^{165}\text{Eu}$	T : average 16Wu.A=2.14(0.45) 08Os02=2.7(0.3)										
* $^{165}\text{Gd}$	T : unweighed average 16Wu.A=12.5(1.3) 98Ic02=9.3(2.3) and 11.2(0.3)										
* $^{165}\text{Ta}$	J : favored $\alpha$ decay from $^{169}\text{Re}^m$ ( $J=(1/2^+, 3/2^+)$ )										
* $^{165}\text{Ta}^m$	J : favored $\alpha$ decay from $^{169}\text{Re}$ ( $J=(9/2^-)$ )										
* $^{165}\text{Re}$	T : symmetrized from 05Sc22=2.614(+0.142-0.128); also 12Th13=1.6(0.6)										
* $^{165}\text{Ir}^m$	T : average 14Dr02=340(40) 97Da07=290(60)										
$^{166}\text{Sm}$	-40730#	400#			800 ms	630	0 <sup>+</sup>		16Wu.A TD	2016	$\beta^- = 100$
$^{166}\text{Eu}$	-47210#	360#			1.24 s	0.12	6 <sup>-</sup> #	14	16Wu.A T	2007	$\beta^- = 100; \beta^-_n = 0.6\#$
$^{166}\text{Gd}$	-54530#	200#			5.1 s	0.8	0 <sup>+</sup>	15	16Wu.A T	2005	$\beta^- = 100$
$^{166}\text{Gd}^m$	-52930#	200#	1601.5	1.1	950 ns	60	(6 <sup>-</sup> )	15		2014	IT=100
$^{166}\text{Tb}$	-57880	70			27.1 s	1.5	(2 <sup>-</sup> )	08	16Wu.A T	1996	$\beta^- = 100$
$^{166}\text{Dy}$	-62584.8	0.9			81.6 h	0.1	0 <sup>+</sup>	08		1949	$\beta^- = 100$
$^{166}\text{Ho}$	-63071.3	1.0			26.824 h	0.012	0 <sup>-</sup>	08		1936	$\beta^- = 100$
$^{166}\text{Ho}^m$	-63065.3	1.0	5.969	0.012	1.133 ky	0.05	7 <sup>-</sup>	08	12Ne05 T	1952	$\beta^- = 100$
$^{166}\text{Ho}^n$	-62880.4	1.0	190.9021	0.0020	185 $\mu\text{s}$	15	3 <sup>+</sup>	08		1960	IT=100
$^{166}\text{Er}$	-64926.0	1.2			STABLE		0 <sup>+</sup>	08		1934	IS=33.503 36
$^{166}\text{Tm}$	-61888	12			7.70 h	0.03	2 <sup>+</sup>	08		1948	$\beta^+ = 100$
$^{166}\text{Tm}^m$	-61771	13	117	5	348 ms	21	(6 <sup>-</sup> )	08	96Dr07 T	1996	IT=100
$^{166}\text{Tm}^n$	-61649	13	239	5	2 $\mu\text{s}$	1	(6 <sup>-</sup> )	08	96Dr07 EDT	1995	IT=100
$^{166}\text{Yb}$	-61596	7			56.7 h	0.1	0 <sup>+</sup>	08		1954	$\epsilon = 100$
$^{166}\text{Lu}$	-56021	30			2.65 m	0.10	6 <sup>-</sup>	08		1969	$\beta^+ = 100$
$^{166}\text{Lu}^m$	-55990	30	34.37	0.22	1.41 m	0.10	3 <sup>(-)</sup>	08		1974	$\beta^+ = 58.5; \text{IT} = 42.5$
$^{166}\text{Lu}^n$	-55980	30	43.0	0.4	2.12 m	0.10	0 <sup>-</sup>	08		1974	$\beta^+ > 80; \text{IT} < 20$
$^{166}\text{Hf}$	-53859	28			6.77 m	0.30	0 <sup>+</sup>	08		1965	$\beta^+ = 100$
$^{166}\text{Ta}$	-46098	28			34.4 s	0.5	(2 <sup>+</sup> )	08		1977	$\beta^+ = 100$
$^{166}\text{W}$	-41888	9			19.2 s	0.6	0 <sup>+</sup>	08		1975	$\beta^+ \approx 100; \alpha = 0.035 12$
$^{166}\text{Re}$	-31890	70			2.25 s	0.21	(7 <sup>+</sup> )	08	92Me10 J	1978	$\beta^+ = ?; \alpha = 5.2$
$^{166}\text{Re}^p$	-31740#	90#	150#	50#			3 <sup>-</sup> #	08			
$^{166}\text{Os}$	-25432	18			213 ms	5	0 <sup>+</sup>	16		1977	$\alpha = 72.13; \beta^+ = 28.13$
$^{166}\text{Ir}$	-13350#	200#			10.5 ms	2.2	(2 <sup>-</sup> )	08		1981	$\alpha = 93.3; p = 7.3$
$^{166}\text{Ir}^m$	-13180#	200#	171	6	15.1 ms	0.9	(9 <sup>+</sup> )	08		1996	$\alpha = 98.2 6; p = 1.8 6$
$^{166}\text{Pt}$	-4730#	300#			300 $\mu\text{s}$	100	0 <sup>+</sup>	08		1996	$\alpha = 100$
* $^{166}\text{Eu}$	T : symmetrized from 16Wu.A=1.27(+0.09-0.14)										
* $^{166}\text{Gd}$	T : average 16Wu.A=5.4(1.2) 05Ic02=00As.A=4.8(1.0)										
* $^{166}\text{Tb}$	T : average 16Wu.A=28.3(2.0) 05Ic02=00As.A=25.6(2.2)										
* $^{166}\text{Tm}^m$	E : less than 16 keV above 109.338 level										
* $^{166}\text{Tm}^n$	T : average 340(25) (34.4 keV $\gamma$ -time) 370(40) (74.9 keV $\gamma$ -time)										
* $^{166}\text{Tm}^n$	E : 121.710 keV above the 340 ms isomer										
* $^{166}\text{Tm}^n$	T : other 02Ca46=36(2) ns adopted in ENSDF'08										
* $^{166}\text{Re}$	D : from 2% < $\alpha$ < 8% as discussed in ENSDF J : 92Me10 $\beta^+$ to 6 <sup>+</sup> state										
$^{167}\text{Eu}$	-44010#	400#			1.33 s	0.51	5/2 <sup>+</sup> #	13	16Wu.A TD	2012	$\beta^- = 100; \beta^-_n = 3\#$
$^{167}\text{Gd}$	-50810#	300#			4.2 s	0.3	5/2 <sup>-</sup> #	13	16Wu.A TD	2012	$\beta^- = 100$
$^{167}\text{Tb}$	-55930#	200#			18.9 s	1.6	3/2 <sup>+</sup> #	00	16Wu.A T	1999	$\beta^- = 100$
$^{167}\text{Dy}$	-59930	60			6.20 m	0.08	(1/2 <sup>-</sup> )	00		1960	$\beta^- = 100$
$^{167}\text{Ho}$	-62281	5			3.1 h	0.1	7/2 <sup>-</sup>	00		1955	$\beta^- = 100$
$^{167}\text{Ho}^m$	-62022	5	259.34	0.11	6.0 $\mu\text{s}$	1.0	3/2 <sup>+</sup>	00		1977	IT=100
$^{167}\text{Er}$	-63291.2	1.2			STABLE		7/2 <sup>+</sup>	00		1934	IS=22.869 9
$^{167}\text{Er}^m$	-63083.4	1.2	207.801	0.005	2.269 s	0.006	1/2 <sup>-</sup>	00		1986	IT=100
$^{167}\text{Tm}$	-62543.6	1.3			9.25 d	0.02	1/2 <sup>+</sup>	00		1948	$\epsilon = 100$
$^{167}\text{Tm}^m$	-62364.1	1.3	179.480	0.019	1.16 $\mu\text{s}$	0.06	(7/2 <sup>+</sup> )	00		1964	IT=100
$^{167}\text{Tm}^n$	-62250.8	1.3	292.820	0.020	0.9 $\mu\text{s}$	0.1	7/2 <sup>-</sup>	00		1965	IT=100
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)					
... A-group continued ...															
<sup>167</sup> Yb	-60591	4		17.5	m	0.2	5/2 <sup>-</sup>	00		1954	$\beta^+=100$				
<sup>167</sup> Yb <sup>m</sup>	-60019	4	571.548	0.022			180	ns		1976	IT=100				
<sup>167</sup> Lu	-57500	30		51.5	m	1.0	7/2 <sup>+</sup>	06		1958	$\beta^+=100$				
<sup>167</sup> Lu <sup>m</sup>	-57500#	40#	0#	30#	*		> 1	m		1998	IT ?; $\beta^+ ?$				
<sup>167</sup> Hf	-53468	28		2.05	m	0.05	(5/2) <sup>-</sup>	00		1969	$\beta^+=100$				
<sup>167</sup> Ta	-48351	28		1.33	m	0.07	(3/2 <sup>+</sup> )	00		1982	$\beta^+=100$				
<sup>167</sup> W	-42098	18		19.9	s	0.5	3/2 <sup>-</sup> #	00		1985	$\beta^+=99.96$ 1; $\alpha=0.04$ 1 *				
<sup>167</sup> Re	-34830#	40#		3.4	s	0.4	(9/2 <sup>-</sup> )	00	10An01	J	1992	$\alpha\approx 100$ ; $\beta^+ ?$			
<sup>167</sup> Re <sup>m</sup>	-34700	40	128#	13#	&		5.9	s	0.3	11Ko.B	EJ	1984	$\beta^+\approx 99$ ; $\alpha\approx 1$		
<sup>167</sup> Os	-26500	70		839	ms	5	7/2 <sup>-</sup>	09	10Sc02	TJD	1977	$\alpha=51$ 4; $\beta^+ ?$ *			
<sup>167</sup> Os <sup>m</sup>	-26060	70	435.1	1.0			672	ns	7	10Sc02	E	2009	IT=100 *		
<sup>167</sup> Ir	-17072	18		29.3	ms	0.6	1/2 <sup>+</sup>	02	05Sc22	TD	1981	$\alpha=43$ 2; $p=39.3$ 13; $\beta^+ ?$ *			
<sup>167</sup> Ir <sup>m</sup>	-16897	18	175.5	2.1	p		25.7	ms	0.8	04Ke06	T	1995	$\alpha=90$ 3; $\beta^+ ?$ ; $p=0.42$ 8 *		
<sup>167</sup> Pt	-6610#	300#		800	$\mu$ s	160	7/2 <sup>-</sup> #	00	04Ke06	T	1996	$\alpha=100$ *			
* <sup>167</sup> Gd	T : symmetrized from 4.26(+0.18-0.32)														
* <sup>167</sup> Tb	T : average 16Wu.A=18.6(2.0) 99As03=19.4(2.7)														
* <sup>167</sup> W	J : lowest observed state in 92Th06 is 13/2 <sup>+</sup>														
* <sup>167</sup> Os	D : average 10Sc02=51(5)% 96Pa01=49(7)% 81Ho10=58(12)%														
* <sup>167</sup> Os <sup>m</sup>	E : also 10Sc02=434.3(1.1), unc. estimated by evaluator, based on Table II														
* <sup>167</sup> Ir	T : from p-decay; $\alpha$ -decay 05Sc22=30.9(1.3) 97Da07=35.2(2.0) not used														
* <sup>167</sup> Ir <sup>m</sup>	T : other not used 05Sc22=28.7(3.3) from $\alpha$ -decay and 28.8(1.3) from p-decay														
* <sup>167</sup> Ir <sup>m</sup>	T : 97Da07=30.0(0.6) conflicting, not used D : p from 05Sc22														
* <sup>167</sup> Pt	T : average 04Ke06=900(+300-200) 96Bi07=700(200)														
<sup>168</sup> Eu	-39740#	500#		200	ms	100	2 <sup>+</sup> #	13	16Wu.A	TD	2012	$\beta^-=100$ ; $\beta^-n=10\#$			
<sup>168</sup> Gd	-48360#	400#		3.03	s	0.16	0 <sup>+</sup>	13	16Wu.A	TD	1985	$\beta^-=100$ *			
<sup>168</sup> Tb	-52720#	300#		9.4	s	0.4	(4 <sup>-</sup> )	10	16Wu.A	T	1999	$\beta^-=100$			
<sup>168</sup> Tb <sup>m</sup>	-52510#	300#	211	2			0.71	$\mu$ s	0.03	(6 <sup>+</sup> )	16Gu.A	ETJ	2016	IT=100	
<sup>168</sup> Dy	-58560	140		8.7	m	0.3	0 <sup>+</sup>	10			1982	$\beta^-=100$			
<sup>168</sup> Ho	-60060	30		2.99	m	0.07	3 <sup>+</sup>	10			1960	$\beta^-=100$			
<sup>168</sup> Ho <sup>m</sup>	-60000	30	59	1			132	s	4	(6 <sup>+</sup> )	90Ch37	E	1990	IT $\approx$ 100; $\beta^- < 0.5$	
<sup>168</sup> Ho <sup>n</sup>	-59920	30	143.43	0.17			> 4	$\mu$ s		(1 <sup>-</sup> )	10		1990	IT=100	
<sup>168</sup> Ho <sup>p</sup>	-59870	30	192.57	0.20			108	ns	11	1 <sup>+</sup>	10		1990	IT=100	
<sup>168</sup> Er	-62991.2	1.2		109.0	ns	0.7	0 <sup>+</sup>	10		0 <sup>+</sup>	10		1934	IS=26.978 18	
<sup>168</sup> Er <sup>m</sup>	-61897.2	1.2	1094.0383	0.0016			109.0	ns	0.7	4 <sup>-</sup>	10		1974	IT=100	
<sup>168</sup> Tm	-61312.9	1.7		93.1	d	0.2	3 <sup>+</sup>	10		3 <sup>+</sup>	10		1949	$\beta^+\approx 100$ ; $\beta^-=0.010$ 7	
<sup>168</sup> Yb	-61581.9	1.2		STABLE		(>130 Ty)	0 <sup>+</sup>	10	56Po16	T	1938	IS=0.123 3; $\alpha ?$ ; $2\beta^+ ?$ *			
<sup>168</sup> Lu	-57070	40		5.5	m	0.1	6 <sup>(-)</sup>	10			1960	$\beta^+=100$			
<sup>168</sup> Lu <sup>m</sup>	-56870	40	202.81	0.12			6.7	m	0.4	3 <sup>+</sup>	10		1960	$\beta^+ > 99.6$ 4; IT<0.8	
<sup>168</sup> Hf	-55361	28		25.95	m	0.20	0 <sup>+</sup>	10			1961	$\epsilon\approx 98$ ; $\epsilon^+\approx 2$			
<sup>168</sup> Ta	-48394	28		2.0	m	0.1	(2 <sup>-</sup> , 3 <sup>+</sup> )	10			1969	$\beta^+=100$ *			
<sup>168</sup> W	-44893	13		50.9	s	1.9	0 <sup>+</sup>	10			1971	$\beta^+\approx 100$ ; $\alpha=0.0032$ 10			
<sup>168</sup> Re	-35790	30		4.4	s	0.1	(7 <sup>+</sup> )	10			1992	$\beta^+\approx 100$ ; $\alpha\approx 0.005$			
<sup>168</sup> Os	-29995	10		2.1	s	0.1	0 <sup>+</sup>	10			1977	$\beta^+=57$ 4; $\alpha=43$ 4			
<sup>168</sup> Ir	-18670	60		230	ms	50	(2 <sup>-</sup> )	10			1978	$\alpha\approx 100$ ; $\beta^+ ?$ ; $\beta^+p ?$ *			
<sup>168</sup> Ir <sup>m</sup>	-18620	240	50	250			163	ms	16	(9 <sup>+</sup> )	10	09Ha42	TD	1996	$\alpha=77$ 9; $\beta^+ ?$ ; $\beta^+p ?$ *
<sup>168</sup> Pt	-11010	150		2.02	ms	0.10	0 <sup>+</sup>	10			1981	$\alpha\approx 100$ ; $\beta^+=0.2\#$			
* <sup>168</sup> Gd	I : first seen in 85Si25 via thermal fission of <sup>252</sup> Cf														
* <sup>168</sup> Tb	T : average 16Wu.A=9.49(0.39) 99As03=8.2(1.3) J : 16Gu.A=(4 <sup>-</sup> )														
* <sup>168</sup> Yb	T : lower limit is for $\alpha$ decay														
* <sup>168</sup> Ta	T : other: 02At01=5.2(0.7) for q=73 <sup>+</sup> (bare ion)														
* <sup>168</sup> Ir	T : symmetrized from 09Ha42=222(+60-40)														
* <sup>168</sup> Ir	J : from correlations between $\alpha$ 's depopulating (2 <sup>-</sup> ) isomers down to <sup>152</sup> Tm														
* <sup>168</sup> Ir <sup>m</sup>	T : average 09Ha42=160(+30-20) 09Ha42=153(+40-30)(indep) 96Pa01=161(21)														
* <sup>168</sup> Ir <sup>m</sup>	J : from correlations between $\alpha$ 's depopulating (9 <sup>+</sup> ) isomers down to <sup>152</sup> Tm														
<sup>169</sup> Gd	-44150#	500#		750	ms	210	7/2 <sup>-</sup> #	13	16Wu.A	TD	2012	$\beta^-=100$ ; $\beta^-n=0\#$			
<sup>169</sup> Tb	-50330#	300#		5.13	s	0.32	3/2 <sup>+</sup> #	13	16Wu.A	TD	2012	$\beta^-=100$ ; $\beta^-n=0.01\#$			
<sup>169</sup> Dy	-55600	300		39	s	8	(5/2) <sup>-</sup>	08			1990	$\beta^-=100$			
<sup>169</sup> Ho	-58797	20		4.72	m	0.10	7/2 <sup>-</sup>	08			1963	$\beta^-=100$			
<sup>169</sup> Ho <sup>m</sup>	-57411	20	1386.2	0.4			118	$\mu$ s	6	(19/2 <sup>+</sup> )	10Dr05	ETJ	2010	IT=100	
<sup>169</sup> Er	-60923.1	1.2		9.392	d	0.018	1/2 <sup>-</sup>	08			1956	$\beta^-=100$			
<sup>169</sup> Er <sup>m</sup>	-60831.1	1.2	92.05	0.10			285	ns	20	(5/2) <sup>-</sup>	08		1969	IT=100	
<sup>169</sup> Er <sup>m</sup>	-60679.4	1.2	243.69	0.17			200	ns	10	7/2 <sup>+</sup>	08		1969	IT=100	
<sup>169</sup> Tm	-61275.2	0.8		STABLE			1/2 <sup>+</sup>	08			1934	IS=100.			
<sup>169</sup> Tm <sup>m</sup>	-60959.1	0.8	316.1463	0.0001			659.9	ns	2.3	7/2 <sup>+</sup>	08		1950	IT=100 *	
<sup>169</sup> Yb	-60377.6	1.2		32.018	d	0.005	7/2 <sup>+</sup>	08			1946	$\epsilon=100$			
<sup>169</sup> Yb <sup>m</sup>	-60353.4	1.2	24.1999	0.0016			46	s	2	1/2 <sup>-</sup>	08		1949	IT=100	
<sup>169</sup> Lu	-58085	3		34.06	h	0.05	7/2 <sup>+</sup>	08			1955	$\beta^+=100$			
<sup>169</sup> Lu <sup>m</sup>	-58056	3	29.0	0.5			160	s	10	(1/2 <sup>-</sup> )	08		1965	IT=100	
... A-group is continued on next page ...															

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)				Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...													
<sup>169</sup> Hf	-54717	28				3.24 m	0.04	(5/2 <sup>-</sup> )	08		1969	$\beta^+$ =100	
<sup>169</sup> Ta	-50290	28				4.9 m	0.4	(5/2 <sup>+</sup> )	08	98Zh03	J	1969	$\beta^+$ =100
<sup>169</sup> W	-44918	15				74 s	6	5/2 <sup>-</sup> #	08			1985	$\beta^+$ =100
<sup>169</sup> Re	-38409	11				8.1 s	0.5	(9/2 <sup>-</sup> )	15	92Me10	D	1978	$\beta^+$ =?; $\alpha$ =0.005 3
<sup>169</sup> Re <sup>m</sup>	-38234	14	175	13	AD	15.1 s	1.5	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	15			1984	$\beta^+$ ?; $\alpha$ ≈0.2; IT?
<sup>169</sup> Os	-30723	25				3.46 s	0.11	(5/2 <sup>-</sup> )	08	96Pa01	T	1972	$\beta^+$ ≈86.3 8; $\alpha$ ≈13.7 8
<sup>169</sup> Ir	-22094	23				353 ms	4	(1/2 <sup>+</sup> )	08	12Th13	D	1978	$\alpha$ =53 7; $\beta^+$ ?
<sup>169</sup> Ir <sup>m</sup>	-21941	12	153	22	AD	280 ms	1	(11/2 <sup>-</sup> )	08	12Th13	TD	1984	$\alpha$ =79 5; $\beta^+$ ?; p?
<sup>169</sup> Pt	-12510#	200#				6.99 ms	0.09	(7/2 <sup>-</sup> )	08	09Go16	T	1981	$\alpha$ =?; $\beta^+$ =1#
<sup>169</sup> Au	-1790#	300#				150# $\mu$ s		1/2 <sup>+</sup> #					p?; $\alpha$ ?; $\beta^+$ ?
* <sup>169</sup> Tm <sup>m</sup>	E: ENSDF2008=316.14633 (0.00011)												
* <sup>169</sup> Re	D: $\alpha$ =0.005(3)% derived from original $\alpha$ =0.001% - 0.01%												
* <sup>169</sup> Re	J: favored $\alpha$ decay from (11/2 <sup>-</sup> ) 173Ir to (11/2 <sup>-</sup> ) level at 136.2 keV												
* <sup>169</sup> Os	T: average 96Pa01=3.6(0.2) 95Hi02=3.2(0.3) 84Sc06=3.5(0.2) 82En03=3.4(0.2)												
* <sup>169</sup> Ir	T: also 12Th13=570(30)												
* <sup>169</sup> Ir	D: average of 12Th13=57(9)% 05Sc22=42(15)% 99Po09=50(18)%												
* <sup>169</sup> Ir <sup>m</sup>	D: average 12Th13=78(6)% 99Po09=84(8)% 96Pa01=72(13)%; 05Sc22=59(4)% at												
* <sup>169</sup> Ir <sup>m</sup>	D: variance, not used												
* <sup>169</sup> Pt	T: average 09Go16=6.99(0.10) 04Ke06=7.0(0.2)												
<sup>170</sup> Gd	-41380#	600#				420 ms	130	0 <sup>+</sup>	13	16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ -n=0#
<sup>170</sup> Tb	-46720#	400#				960 ms	68	2 <sup>-</sup> #	13	16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ -n=0.01#
<sup>170</sup> Dy	-53660#	200#				54.9 s	8.0	0 <sup>+</sup>	10	16Wu.A	TD	2010	$\beta^-$ =100
<sup>170</sup> Dy <sup>m</sup>	-52020#	200#	1643.92	0.22		0.94 $\mu$ s	0.16	(6 <sup>+</sup> )		16So.A	ETJ	2016	IT=100
<sup>170</sup> Ho	-56240	50			*	2.76 m	0.05	6 <sup>+</sup> #	02			1960	$\beta^-$ =100
<sup>170</sup> Ho <sup>m</sup>	-56140	60	100	80	BD *	43 s	2	(1 <sup>+</sup> )	02			1960	$\beta^-$ =100
<sup>170</sup> Er	-60108.7	1.5				STABLE	(>320 Py)	0 <sup>+</sup>	02	96De60	T	1934	IS=14.910 36; 2 $\beta^-$ ?; $\alpha$ ?
<sup>170</sup> Tm	-59795.9	0.8				128.6 d	0.3	1 <sup>-</sup>	02			1936	$\beta^-$ ≈100; $\epsilon$ =0.131 10
<sup>170</sup> Tm <sup>m</sup>	-59612.7	0.8	183.197	0.004		4.12 $\mu$ s	0.13	(3) <sup>+</sup>	02			1967	IT=100
<sup>170</sup> Yb	-60763.919	0.010				STABLE		0 <sup>+</sup>	02			1938	IS=2.982 39
<sup>170</sup> Yb <sup>m</sup>	-59505.46	0.14	1258.46	0.14		370 ns	15	4 <sup>-</sup>	02			1981	IT=100
<sup>170</sup> Lu	-57306	17				2.012 d	0.020	0 <sup>+</sup>	02			1951	$\beta^-$ =100
<sup>170</sup> Lu <sup>m</sup>	-57213	17	92.91	0.09		670 ms	100	(4) <sup>-</sup>	02			1965	IT=100
<sup>170</sup> Hf	-56254	28				16.01 h	0.13	0 <sup>+</sup>	06			1961	$\epsilon$ =100
<sup>170</sup> Ta	-50138	28				6.76 m	0.06	(3) <sup>(+)</sup> #	02			1969	$\beta^-$ =100
<sup>170</sup> W	-47291	13				2.42 m	0.04	0 <sup>+</sup>	02			1971	$\beta^-$ ≈100; $\alpha$ <1#
<sup>170</sup> Re	-38913	23				9.2 s	0.2	(5 <sup>+</sup> )	02			1974	$\beta^-$ ≈100; $\alpha$ <0.01#
<sup>170</sup> Os	-33926	10				7.37 s	0.18	0 <sup>+</sup>	08			1972	$\beta^-$ =?; $\alpha$ ≈9.5 10
<sup>170</sup> Ir	-23360#	90#				910 ms	150	(3 <sup>-</sup> )	08			1977	$\beta^-$ ?; $\alpha$ ≈5.2 17
<sup>170</sup> Ir <sup>m</sup>	-23200	70	160#	50#		811 ms	18	(8 <sup>+</sup> )	08			1977	$\alpha$ ≈36 10; $\beta^-$ ?; IT?
<sup>170</sup> Pt	-16299	18				13.93 ms	0.16	0 <sup>+</sup>	02	04Ke06	T	1981	$\alpha$ =?; $\beta^-$ =2#
<sup>170</sup> Au	-3750#	200#				290 $\mu$ s	50	(2 <sup>-</sup> )	02	04Ke06	TD	2002	p=89 10; $\alpha$ =11 10
<sup>170</sup> Au <sup>m</sup>	-3470#	200#	280	13	p	620 $\mu$ s	50	(9 <sup>+</sup> )	02	04Ke06	TD	2002	p=58 5; $\alpha$ =42 5
* <sup>170</sup> Dy <sup>m</sup>	T: symmetrized from 16So.A=0.91(+0.18-0.13)												
* <sup>170</sup> Ir	T: symmetrized from 870(+180-120)												
* <sup>170</sup> Pt	T: average 04Ke06=14.0(0.2) 98Ki20=13.5(0.3) 96Bi07=14.7(0.5)												
* <sup>170</sup> Au	T: symmetrized from 286(+50-40)												
* <sup>170</sup> Au <sup>m</sup>	T: 04Ke06=617(+50-40); other 02Ma61=570(+310-150) D: and 02Ma61=75(15)%												
<sup>171</sup> Tb	-44030#	500#				1.23 s	0.10	3/2 <sup>+</sup> #	13	16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ -n=1#
<sup>171</sup> Dy	-50190#	300#				4.07 s	0.40	7/2 <sup>-</sup> #	13	16Wu.A	TD	2012	$\beta^-$ =100
<sup>171</sup> Ho	-54520	600				53 s	2	7/2 <sup>-</sup> #	02			1989	$\beta^-$ =100
<sup>171</sup> Er	-57719.0	1.6				7.516 h	0.002	5/2 <sup>-</sup>	02			1938	$\beta^-$ =100
<sup>171</sup> Er <sup>m</sup>	-57520.4	1.6	198.6	0.1		210 ns	10	1/2 <sup>-</sup>	02			1969	IT=100
<sup>171</sup> Tm	-59210.3	1.0				1.92 y	0.01	1/2 <sup>+</sup>	02			1948	$\beta^-$ =100
<sup>171</sup> Tm <sup>m</sup>	-58785.3	1.0	424.9560	0.0015		2.60 $\mu$ s	0.02	7/2 <sup>-</sup>	02			1948	IT=100
<sup>171</sup> Tm <sup>n</sup>	-57535.8	1.0	1674.5	0.3		1.7 $\mu$ s	0.2	19/2 <sup>+</sup>		09Wa06	ETJ	2009	IT=100
<sup>171</sup> Yb	-59306.810	0.013				STABLE		1/2 <sup>-</sup>	02			1934	IS=14.09 14
<sup>171</sup> Yb <sup>m</sup>	-59211.528	0.013	95.282	0.002		5.25 ms	0.24	7/2 <sup>+</sup>	02			1968	IT=100
<sup>171</sup> Yb <sup>n</sup>	-59184.394	0.013	122.416	0.002		265 ns	20	5/2 <sup>-</sup>	02			1968	IT=100
<sup>171</sup> Lu	-57828.4	1.9				8.24 d	0.03	7/2 <sup>+</sup>	02			1951	$\beta^-$ =100
<sup>171</sup> Lu <sup>m</sup>	-57757.3	1.9	71.13	0.08		79 s	2	1/2 <sup>-</sup>	02			1965	IT=100
<sup>171</sup> Hf	-55431	29				12.1 h	0.4	7/2 <sup>+</sup>	02	00Ye02	J	1951	$\beta^-$ =100
<sup>171</sup> Hf <sup>m</sup>	-55409	29	21.93	0.09		29.5 s	0.9	1/2 <sup>-</sup>	02	00Ye02	J	1997	IT≈100; $\beta^+$ ?
<sup>171</sup> Ta	-51720	28				23.3 m	0.3	(5/2 <sup>-</sup> )	02			1969	$\beta^-$ =100
<sup>171</sup> W	-47086	28				2.38 m	0.04	(5/2 <sup>-</sup> )	02			1983	$\beta^-$ =100
<sup>171</sup> Re	-41250	28				15.2 s	0.4	(9/2 <sup>-</sup> )	02			1987	$\beta^-$ =100

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...												
<sup>171</sup> Os	-34302	18				8.3 s	0.2	(5/2 <sup>-</sup> )	02	1972	$\beta^+$ ?; $\alpha=1.80$ 21	
<sup>171</sup> Ir	-26410	40				3.1 s	0.3	1/2 <sup>+</sup>	02	11Ko.B TJ	1967	$\beta^+$ ?; $\alpha=15$ 2
<sup>171</sup> Ir <sup>m</sup>	-26250#	40#	167#	12#		1.47 s	0.06	(11/2 <sup>-</sup> )	02	11Ko.B T	1967	$\alpha=54$ 5; $\beta^+$ ?; p ?
<sup>171</sup> Pt	-17470	70				45.5 ms	2.5	7/2 <sup>-</sup>	10	10Sc02 J	1981	$\alpha=90$ 7; $\beta^+$ ?
<sup>171</sup> Pt <sup>m</sup>	-17060	70	412.6	1.0		901 ns	9	13/2 <sup>+</sup>	10	FGK128 J	2010	IT=100
<sup>171</sup> Au	-7562	21				22.3 $\mu$ s	2.4	(1/2 <sup>+</sup> )	02	04Ke06 T	1997	p $\approx$ 100; $\alpha$ ?
<sup>171</sup> Au <sup>m</sup>	-7308	18	255	10	p	1.036 ms	0.016	11/2 <sup>-</sup>	02	04Ke06 TD	1996	$\alpha=60.0$ 28; p=40.0 28
<sup>171</sup> Hg	3480#	300#				70 $\mu$ s	30	3/2 <sup>-</sup> #	04		2004	$\alpha\approx$ 100; $\beta^+=0.01$ #
* <sup>171</sup> Tb	T : symmetrized from 1.24(+0.09-0.10)											
* <sup>171</sup> Ir	T : other 02Ro17=3.2(+1.3-0.7) D : 13An10=15(2)											
* <sup>171</sup> Ir <sup>m</sup>	D : average 10An01=53(5)% 96Pa01=58(11)%											
* <sup>171</sup> Ir <sup>m</sup>	T : average 11Ko.B=1.50(0.07) 10An01=1.40(0.10)											
* <sup>171</sup> Pt <sup>m</sup>	J : M2 to 9/2 <sup>-</sup>											
* <sup>171</sup> Au	T : average 04Ke06=22(+3-2) 99Po09=17(+9-5)											
* <sup>171</sup> Au	T : other 03Ba20=37(+7-5) conflicting, not used											
* <sup>171</sup> Au <sup>m</sup>	T : average 04Ke06=1.09(0.03) 03Ba20=1.014(0.019)											
* <sup>171</sup> Au <sup>m</sup>	D : average 04Ke06=34(4)% 97Da07=46(4)%; Birge ratio B=2.1											
* <sup>171</sup> Hg	T : symmetrized from 59(+36-16)											
<sup>172</sup> Tb	-39850#	500#				760 ms	190	6 <sup>+</sup> #	13	16Wu.A TD	2012	$\beta^-$ =100; $\beta^-$ n=1#
<sup>172</sup> Dy	-48010#	300#				3.4 s	0.2	0 <sup>+</sup>	13	16Wa19 TD	2012	$\beta^-$ =100
<sup>172</sup> Dy <sup>m</sup>	-46730#	300#	1278	1		710 ms	50	(8 <sup>-</sup> )		16Wa19 ETJ	2016	$\beta^-$ =19 3; IT=81 3
<sup>172</sup> Ho	-51480#	200#				25 s	3	0 <sup>+</sup> #	15		1991	$\beta^-$ =100
<sup>172</sup> Er	-56484	4				49.3 h	0.5	0 <sup>+</sup>	15		1956	$\beta^-$ =100
<sup>172</sup> Er <sup>m</sup>	-54983	4	1500.9	0.3		579 ns	62	(6 <sup>+</sup> )	15	10Dr02 ETJ	2006	IT=100
<sup>172</sup> Tm	-57374	6				63.6 h	0.3	2 <sup>-</sup>	15		1956	$\beta^-$ =100
<sup>172</sup> Tm <sup>m</sup>	-56898	6	476.2	0.2		132 $\mu$ s	7	(6 <sup>+</sup> )	15		2008	IT=100
<sup>172</sup> Yb	-59255.446	0.014				STABLE		0 <sup>+</sup>	95		1934	IS=21.68 13
<sup>172</sup> Yb <sup>m</sup>	-57705.02	0.06	1550.43	0.06		3.6 $\mu$ s	0.1	6 <sup>-</sup>	95		1969	IT=100
<sup>172</sup> Lu	-56736.0	2.3				6.70 d	0.03	4 <sup>-</sup>	95		1951	$\beta^+$ =100
<sup>172</sup> Lu <sup>m</sup>	-56694.1	2.3	41.86	0.04		3.7 m	0.5	1 <sup>-</sup>	95		1962	IT=100; $\beta^+ < 0.18$
<sup>172</sup> Lu <sup>n</sup>	-56670.2	2.3	65.79	0.04		332 ns	20	(1) <sup>+</sup>	95		1965	IT=100
<sup>172</sup> Lu <sup>p</sup>	-56626.6	2.3	109.41	0.10		440 $\mu$ s	12	(1) <sup>+</sup>	95		1965	IT=100
<sup>172</sup> Lu <sup>q</sup>	-56522.4	2.3	213.57	0.17		150 ns		(6 <sup>-</sup> )	95		1974	IT=100
<sup>172</sup> Hf	-56402	24				1.87 y	0.03	0 <sup>+</sup>	95		1951	$\epsilon=100$
<sup>172</sup> Hf <sup>m</sup>	-54396	24	2005.84	0.11		163 ns	3	(8 <sup>-</sup> )	95		1976	IT=100
<sup>172</sup> Ta	-51330	28				36.8 m	0.3	(3 <sup>+</sup> )	15		1964	$\beta^+$ =100
<sup>172</sup> W	-49097	28				6.6 m	0.9	0 <sup>+</sup>	95		1964	$\beta^+$ =100
<sup>172</sup> Re	-41540	40			*	15 s	3	(5 <sup>+</sup> )	16		1972	$\beta^+$ =100
<sup>172</sup> Re <sup>m</sup>	-41540#	110#	0#	100#	*	55 s	5	(2)	16		1977	$\beta^+$ =100
<sup>172</sup> Os	-37244	13				19.2 s	0.9	0 <sup>+</sup>	95	95Hi02 D	1971	$\beta^+=?$ ; $\alpha=1.1$ 2
<sup>172</sup> Ir	-27380	30				4.4 s	0.3	(3 <sup>-</sup> , 4 <sup>-</sup> )	16		1967	$\beta^+$ ?; $\alpha=2$
<sup>172</sup> Ir <sup>m</sup>	-27240	30	139	10	AD	2.19 s	0.07	(7 <sup>+</sup> )	16		1967	$\beta^+$ ?; $\alpha=9.5$ 11
<sup>172</sup> Pt	-21107	10				97.6 ms	1.3	0 <sup>+</sup>	10	10An02 D	1981	$\alpha=97$ 3; $\beta^+$ ?
<sup>172</sup> Au	-9320	60				28 ms	4	(2 <sup>-</sup> )	10		1993	$\alpha=?$ ; p<2; $\beta^+$ ?
<sup>172</sup> Au <sup>m</sup>	-9160	240	160	250		11.0 ms	1.0	(9 <sup>+</sup> )	10	09Ha42 T	1993	$\alpha=?$ ; p<2
<sup>172</sup> Hg	-1060	150				231 $\mu$ s	9	0 <sup>+</sup>	10		1999	$\alpha\approx$ 100; $\beta^+=0.1$ #
* <sup>172</sup> Au	T : symmetrized from 09Ha42=22(+6-4)											
* <sup>172</sup> Au	J : from correlations between $\alpha$ 's depopulating (2 <sup>-</sup> ) isomers down to <sup>152</sup> Tm											
* <sup>172</sup> Au <sup>m</sup>	T : average 09Ha42=9(+2-1) 09Ha42=8(+5-2) (independent measurements)											
* <sup>172</sup> Au <sup>m</sup>	T : others 96Pa01=6.3(1.5) 93Se09=4(1)											
<sup>173</sup> Dy	-43940#	400#				1.43 s	0.20	9/2 <sup>+</sup> #	13	16Wu.A TD	2012	$\beta^-$ =100
<sup>173</sup> Ho	-49350#	300#				6.90 s	0.48	7/2 <sup>-</sup> #	13	16Wu.A TD	2012	$\beta^-$ =100
<sup>173</sup> Er	-53650#	200#				1.434 m	0.017	(7/2 <sup>-</sup> )	95	94It.A T	1972	$\beta^-$ =100
<sup>173</sup> Tm	-56256	4				8.24 h	0.08	(1/2 <sup>+</sup> )	95		1961	$\beta^-$ =100
<sup>173</sup> Tm <sup>m</sup>	-55938	4	317.73	0.20		10.7 $\mu$ s	1.7	7/2 <sup>-</sup>	95	12Hu10 TJ	1972	IT=100
<sup>173</sup> Tm <sup>n</sup>	-54350	4	1905.7	0.4		250 ns	69	19/2 <sup>-</sup>	95	12Hu10 ETJ	2012	IT=100
<sup>173</sup> Tm <sup>p</sup>	-52208	4	4047.9	0.5		121 ns	28	35/2 <sup>-</sup>	95	12Hu10 ETJ	2012	IT=100
<sup>173</sup> Yb	-57551.225	0.011				STABLE		5/2 <sup>-</sup>	95		1934	IS=16.103 63
<sup>173</sup> Yb <sup>m</sup>	-57152.3	0.5	398.9	0.5		2.9 $\mu$ s	0.1	1/2 <sup>-</sup>	95		1963	IT=100
<sup>173</sup> Lu	-56880.9	1.6				1.37 y	0.01	7/2 <sup>+</sup>	95		1951	$\epsilon=100$
<sup>173</sup> Lu <sup>m</sup>	-56757.2	1.6	123.672	0.013		74.2 $\mu$ s	1.0	5/2 <sup>-</sup>	95		1962	IT=100
<sup>173</sup> Hf	-55412	28				23.6 h	0.1	1/2 <sup>-</sup>	06		1951	$\beta^+$ =100
<sup>173</sup> Hf <sup>m</sup>	-55305	28	107.16	0.05		180 ns	8	5/2 <sup>-</sup>	06		1973	IT=100
<sup>173</sup> Hf <sup>n</sup>	-55215	28	197.47	0.10		160 ns	40	7/2 <sup>+</sup>	06		1973	IT=100
... A-group is continued on next page ...												

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>173</sup> Ta	-52397	28	3.14 h	0.13	5/2 <sup>-</sup>	95	1960	$\beta^+=100$			
<sup>173</sup> Ta <sup>m</sup>	-52224	28	225 ns	15	9/2 <sup>-</sup>	95	95Ca27 E	1977 IT=100			
<sup>173</sup> Ta <sup>n</sup>	-50678	28	1719.4	1.0	21/2 <sup>-</sup>	06Th07	ETJ	2006 IT=100			
<sup>173</sup> W	-48727	28	7.6 m	0.2	5/2 <sup>-</sup>	95	1963	$\beta^+=100$			
<sup>173</sup> Re	-43554	28	2.0 m	0.3	(5/2 <sup>-</sup> )	95	1986	$\beta^+=100$			
<sup>173</sup> Os	-37438	15	22.4 s	0.9	5/2 <sup>-</sup>	15	1971	$\beta^+\approx 100; \alpha=0.4\ 2$			
<sup>173</sup> Ir	-30268	11	9.0 s	0.8	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	15	01Ko44 J	1967 $\beta^+>93; \alpha<7$			
<sup>173</sup> Ir <sup>m</sup>	-30042	11	226	9	AD	2.20 s	0.05	(11/2 <sup>-</sup> )	15	01Ko44 J	1967 $\beta^+=88\ 1; \alpha=12\ 1$
<sup>173</sup> Pt	-21940	60	382 ms	2	(5/2 <sup>-</sup> )	15	1966	$\alpha=86\ 4; \beta^+?$			
<sup>173</sup> Au	-12832	23	25.5 ms	0.8	(1/2 <sup>+</sup> )	15	12Th13 T	1983 $\alpha=86\ 13; \beta^+=6\#$			
<sup>173</sup> Au <sup>m</sup>	-12619	12	214	21	AD	12.2 ms	0.1	(11/2 <sup>-</sup> )	15	99Po09 D	1984 $\alpha=89\ 11; \beta^+=4\#$
<sup>173</sup> Hg	-2710#	200#	800 $\mu$ s	80	3/2 <sup>-</sup> #	15	1999	$\alpha=100$			
* <sup>173</sup> Tm <sup>m</sup>	T : average 12Hu10=11.1(2.8) 72Pu02=10.4(2.1)										
* <sup>173</sup> Ta <sup>m</sup>	T : other recent 06Th07=163(2), conflicting, not used										
* <sup>173</sup> Ir	J : favored $\alpha$ decay from (1/2 <sup>+</sup> , 3/2 <sup>+</sup> ) <sup>173</sup> Au ground-state										
* <sup>173</sup> Ir <sup>m</sup>	J : favored $\alpha$ decay from (11/2 <sup>-</sup> ) <sup>173</sup> Au isomer										
* <sup>173</sup> Au	T : average 12Th13=26.3(1.2) 99Po09=25(1)										
* <sup>173</sup> Au	D : from 99Po09=94(+6-19)%; and for isomer <sup>173</sup> Au <sup>m</sup> 92(+8-13)%										
* <sup>173</sup> Hg	J : 12Od01=(7/2 <sup>-</sup> ) based on $\alpha$ chain, not trusted										
<sup>174</sup> Dy	-41370#	500#	1#	s	(>300 ns)	0 <sup>+</sup>	13	12Ku26 I	2012 $\beta^-?$ ; $\beta^-n=0\#$		
<sup>174</sup> Ho	-45690#	300#	3.2 s	1.1	8 <sup>-</sup> #	13	16Wu.A	TD	2012 $\beta^-=100$		
<sup>174</sup> Er	-51950#	300#	3.2 m	0.2	0 <sup>+</sup>	99	1989	$\beta^-=100$			
<sup>174</sup> Er <sup>m</sup>	-50840#	300#	1111.6	1.1	3.9 s	0.3	8 <sup>-</sup>	16Wu.A T	2006 IT=100		
<sup>174</sup> Tm	-53860	40	5.4 m	0.1	(4) <sup>-</sup>	99	1960	$\beta^-=100$			
<sup>174</sup> Tm <sup>m</sup>	-53610	40	252.4	0.5	2.29 s	0.01	(0 <sup>+</sup> )	06Ch10 TJD	2006 IT>98.5; $\beta^-<1.5$		
<sup>174</sup> Yb	-56944.512	0.011	STABLE		142 d	2	0 <sup>+</sup>	99	1934 IS=32.026 80		
<sup>174</sup> Yb <sup>m</sup>	-55426.364	0.017	1518.148	0.013	830 $\mu$ s	40	6 <sup>+</sup>	99	1964 IT=100		
<sup>174</sup> Yb <sup>n</sup>	-55179.3	0.5	1765.2	0.5	256 ns	11	7 <sup>-</sup>	05Dr05 EJT	2005 IT=100		
<sup>174</sup> Lu	-55570.2	1.6	170.83	0.05	3.31 y	0.05	1 <sup>-</sup>	99	98Ge13 J	1951 $\beta^+=100$	
<sup>174</sup> Lu <sup>m</sup>	-55399.4	1.6	240.818	0.004	142 d	2	6 <sup>-</sup>	99	98Ge13 J	1960 IT=99.38 2; $\epsilon=0.62\ 2$	
<sup>174</sup> Lu <sup>n</sup>	-55329.4	1.6	365.183	0.006	395 ns	15	(3 <sup>+</sup> )	99	1980 IT=100		
<sup>174</sup> Lu <sup>p</sup>	-55205.0	1.6	1855.7	0.5	145 ns	3	(4) <sup>-</sup>	99	1980 IT=100		
<sup>174</sup> Lu <sup>q</sup>	-53714.5	1.7	5849.6	0.9	194 ns	24	13 <sup>+</sup>	09Ko19 ETJ	2009 IT=100		
<sup>174</sup> Lu <sup>r</sup>	-49720.6	1.8	242 ns	19	(26 <sup>-</sup> )	09Ko19 ETJ	2009	IT=100			
<sup>174</sup> Hf	-55844.5	2.3	2.0 Py	0.4	0 <sup>+</sup>	04	1939	IS=0.16 1; $\alpha=100; 2\beta^+?$			
<sup>174</sup> Hf <sup>m</sup>	-54295.2	2.9	1549.3	1.8	138 ns	4	6 <sup>+</sup>	04	FGK129 J	1976 IT=100	
<sup>174</sup> Hf <sup>n</sup>	-54047.0	2.9	1797.5	1.8	2.39 $\mu$ s	0.04	8 <sup>-</sup>	04	FGK129 J	1974 IT=100	
<sup>174</sup> Hf <sup>p</sup>	-52532.8	2.9	3311.7	1.8	3.7 $\mu$ s	0.2	14 <sup>+</sup>	04	FGK129 J	1974 IT=100	
<sup>174</sup> Ta	-51741	28	1.14 h	0.08	3 <sup>+</sup>	99	1960	$\beta^+=100$			
<sup>174</sup> W	-50227	28	33.2 m	2.1	0 <sup>+</sup>	99	1964	$\beta^+=100$			
<sup>174</sup> W <sup>m</sup>	-48555	28	1672.0	0.5	> 187 ns	99	1976	IT=100			
<sup>174</sup> W <sup>n</sup>	-48307	28	1919.7	0.5	187 ns	25	99	1976 IT=100			
<sup>174</sup> W <sup>p</sup>	-47959	28	2267.8	0.4	158 ns	3	8 <sup>-</sup>	06Ta13 ETJ	2006 IT=100		
<sup>174</sup> W <sup>q</sup>	-46711	28	3515.6	0.4	128 ns	8	12 <sup>+</sup>	06Ta13 ETJ	2006 IT=100		
<sup>174</sup> Re	-43673	28	2.40 m	0.04	3 <sup>+</sup> #	99	1972	$\beta^+=100$			
<sup>174</sup> Re <sup>m</sup>	-43570#	60#	1#	m	(>1 $\mu$ s)	7 <sup>+</sup> #	12Gu14 T	2012 IT?; $\beta^+?$			
<sup>174</sup> Os	-39995	10	44 s	4	0 <sup>+</sup>	99	1971	$\beta^+\approx 100; \alpha=0.024\ 7$			
<sup>174</sup> Ir	-30863	24	7.9 s	0.6	(3 <sup>+</sup> )	99	1967	$\beta^+=99.5\ 3; \alpha=0.5\ 3$			
<sup>174</sup> Ir <sup>m</sup>	-30671	23	192	11	AD	4.9 s	0.3	(7 <sup>+</sup> )	99	1992 $\beta^+=97.5\ 3; \alpha=2.5\ 3$	
<sup>174</sup> Pt	-25318	10	889 ms	17	0 <sup>+</sup>	99	1966	$\alpha=76\ 8; \beta^+?$			
<sup>174</sup> Au	-14240#	90#	139 ms	3	low	99	02Ro17 TD	1983 $\alpha=90\ 6; \beta^+?$			
<sup>174</sup> Au <sup>m</sup>	-13990	70	171 ms	29	high	96Pa01 TJ	1995	$\alpha=?; \beta^+?$			
<sup>174</sup> Hg	-6641	19	2.0 ms	0.4	0 <sup>+</sup>	99	99Se14 T	1997 $\alpha\approx 100; \beta^+=0.4\#$			
* <sup>174</sup> Er <sup>m</sup>	T : average 16Wu.A=3.37(0.73) 09Dr06=4.02(0.35)										
* <sup>174</sup> Er <sup>m</sup>	E : uncertainty estimated by NUBASE										
* <sup>174</sup> Tm <sup>m</sup>	E : uncertainty estimated by NUBASE										
* <sup>174</sup> Hf <sup>m</sup>	J : multiple decay branches, transition mult., magnetic moment; also n and p										
* <sup>174</sup> W <sup>p</sup>	E : derived from least-squares fit to $\gamma$ -ray energies										
* <sup>174</sup> W <sup>q</sup>	E : derived from least-squares fit to $\gamma$ -ray energies										
* <sup>174</sup> Os	D : symmetrized from 71Bo06 $\alpha=0.020(+10-4)\%$										
* <sup>174</sup> Au	T : others 96Pa01=171(29) 83Sc24=120(20)										
* <sup>174</sup> Hg	T : symmetrized from 1.9(+0.4-0.3)										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)				
<sup>175</sup> Ho	-43200#	400#	1.88 s	0.55	7/2 <sup>-</sup> #	13 16Wu.A	TD	2012	$\beta^-$ =100; $\beta^-$ n=0#			
<sup>175</sup> Er	-48650#	400#	1.2 m	0.3	9/2 <sup>+</sup> #	04		1996	$\beta^-$ =100			
<sup>175</sup> Tm	-52310	50	15.2 m	0.5	1/2 <sup>+</sup> #	04		1961	$\beta^-$ =100			
<sup>175</sup> Tm <sup>m</sup>	-51870	50	440.0	1.1	7/2 <sup>-</sup>	04 12Hu10	ETJ	2012	IT=100			
<sup>175</sup> Tm <sup>n</sup>	-50790	50	1517.7	1.2	21	$\mu$ s 14	12Hu10	ETJ	2012	IT=100		
<sup>175</sup> Yb	-54695.55	0.07	4.185	d	0.001	7/2 <sup>-</sup>	04 12FI05	J	1945	$\beta^-$ =100		
<sup>175</sup> Yb <sup>m</sup>	-54180.68	0.07	514.866	0.004	68.2	ms 0.3	1/2 <sup>-</sup>	04	1972	IT=100		
<sup>175</sup> Lu	-55165.6	1.2	STABLE			7/2 <sup>+</sup>	04	1934	IS=97.401 13			
<sup>175</sup> Lu <sup>m</sup>	-54812.1	1.2	353.48	0.13	1.49	$\mu$ s 0.07	5/2 <sup>-</sup>	04	1965	IT=100		
<sup>175</sup> Lu <sup>n</sup>	-53773.4	1.3	1392.2	0.6	984	$\mu$ s 30	19/2 <sup>+</sup>	04 98Wh02	J	1998	IT=100	
<sup>175</sup> Hf	-54481.7	2.3	70.65	d	0.19	5/2 <sup>(-)</sup>	04 12Fa07	T	1949	$\epsilon$ =100		
<sup>175</sup> Hf <sup>m</sup>	-54355.8	2.3	125.89	0.12	53.7	$\mu$ s 1.5	1/2 <sup>-</sup>	04	1964	IT=100		
<sup>175</sup> Hf <sup>n</sup>	-53048.3	2.3	1433.41	0.12	1.10	$\mu$ s 0.08	19/2 <sup>+</sup>	04 95Gj01	J	1990	IT=100	
<sup>175</sup> Hf <sup>p</sup>	-51466.1	2.3	3015.6	0.4	1.21	$\mu$ s 0.15	35/2 <sup>-</sup>	04 95Gj01	J	1980	IT=100	
<sup>175</sup> Hf <sup>q</sup>	-49845.5	2.6	4636.2	1.2	1.9	$\mu$ s 0.1	45/2 <sup>+</sup>	04 04Ko.A	JT	1990	IT=100	
<sup>175</sup> Ta	-52409	28	10.5	h	0.2	7/2 <sup>+</sup>	04	1960	$\beta^+$ =100			
<sup>175</sup> Ta <sup>m</sup>	-52278	28	131.41	0.17	222	ns 8	9/2 <sup>-</sup>	04 96Ko17	JT	1972	IT=100	
<sup>175</sup> Ta <sup>n</sup>	-52070	28	339.2	1.3	170	ns 20	(1/2 <sup>+</sup> )	04	1969	IT=100		
<sup>175</sup> Ta <sup>p</sup>	-50841	28	1567.6	0.3	1.95	$\mu$ s 0.15	21/2 <sup>-</sup>	04 96Ko17	JT	1996	IT=100	
<sup>175</sup> W	-49633	28	35.2	m	0.6	(1/2 <sup>-</sup> )	04	1963	$\beta^+$ =100			
<sup>175</sup> W <sup>m</sup>	-49398	28	234.96	0.15	216	ns 6	(7/2 <sup>+</sup> )	04	1978	IT=100		
<sup>175</sup> Re	-45288	28	5.89	m	0.05	5/2 <sup>-</sup> #	04	1967	$\beta^+$ =100			
<sup>175</sup> Os	-40105	12	1.4	m	0.1	(5/2 <sup>-</sup> )	04	1972	$\beta^+$ =100			
<sup>175</sup> Ir	-33395	12	9	s	2	5/2 <sup>-</sup> #	04	1967	$\beta^+$ =99.15 28; $\alpha$ =0.85 28			
<sup>175</sup> Pt	-25713	18	2.43	s	0.04	(7/2 <sup>-</sup> )	04 14Pe02	TJ	1966	$\alpha$ =64 5; $\beta^+$ ?		
<sup>175</sup> Au	-17400	40	202	ms	6	1/2 <sup>+</sup>	04 13An10	TJD	1975	$\alpha$ =88 3; $\beta^+$ ?		
<sup>175</sup> Au <sup>m</sup>	-17240#	40#	167#	11#	AD	134	ms 4	(11/2 <sup>-</sup> )	04 11Ko.B	TD	1975	$\alpha$ =75 4; $\beta^+$ ?
<sup>175</sup> Hg	-7970	70	10.6	ms	0.4	(7/2 <sup>-</sup> )	09	1983	$\alpha$ ?; $\beta^+$ =1#			
<sup>175</sup> Hg <sup>m</sup>	-7480	70	494	2	340	ns 30	(13/2 <sup>+</sup> )	09	2009	IT=100		
* <sup>175</sup> Au	T : average 13An10=207(7) 11Ko.B=188(12)								**			
* <sup>175</sup> Au	J : favored $\alpha$ decay to 1/2 <sup>+</sup> states in <sup>171</sup> Ir and <sup>167</sup> Re and from 1/2 <sup>+</sup> in <sup>179</sup> Tl								**			
* <sup>175</sup> Au	D : average 13An10=90(7) 11Ko.B=87(4)								**			
* <sup>175</sup> Au	D : $\alpha$ =87(4) from 11Ko.B, after correction for $\alpha$ =64(5) of <sup>175</sup> Pt daughter								**			
* <sup>175</sup> Au <sup>m</sup>	T : average 11Ko.B=124(8) 10An01=138(5); the former supersedes 01Ko44=143(8)								**			
* <sup>175</sup> Au <sup>n</sup>	T : others 02Ro17=158(3) 96Pa01=185(30) 83Sc24=200(22) for mixture ground-state and m								**			
* <sup>175</sup> Au <sup>m</sup>	J : favored $\alpha$ decay to (11/2 <sup>-</sup> ) excited isomer <sup>171</sup> Ir <sup>m</sup>								**			
* <sup>175</sup> Au <sup>m</sup>	D : $\alpha$ =75(4)% from 11Ko.B, after correction for $\alpha$ =64(5)% of <sup>175</sup> Pt daughter								**			
<sup>176</sup> Ho	-39290#	500#	2#	s	(>300 ns)	13 12Ku26	I	2012	$\beta^-$ ?; $\beta^-$ n=0.1#			
<sup>176</sup> Er	-46630#	400#	20#	s	(>300 ns)	0 <sup>+</sup>	13 12Ku26	I	2012	$\beta^-$ ?		
<sup>176</sup> Tm	-49370	100	1.85	m	0.03	(4 <sup>+</sup> )	06 94It.A	T	1961	$\beta^-$ =100		
<sup>176</sup> Yb	-53491.314	0.015	STABLE			(>160 Py)	0 <sup>+</sup>	06 96De60	T	1934	IS=12.996 83; 2 $\beta^-$ ?; $\alpha$ ?	
<sup>176</sup> Yb <sup>m</sup>	-52441.5	0.6	11.4	s	0.3	8 <sup>-</sup>	06	1967	IT=?; $\beta^-$ <10#			
<sup>176</sup> Lu	-53382.2	1.2	36.84	Gy	0.18	7 <sup>-</sup>	06 14Hu07	T	1935	IS=2.599 13; $\beta^-$ =100		
<sup>176</sup> Lu <sup>m</sup>	-53259.4	1.2	3.664	h	0.019	1 <sup>-</sup>	06	1935	$\beta^-$ ≈100; $\epsilon$ =0.095 16			
<sup>176</sup> Lu <sup>n</sup>	-51867.7	1.3	1514.5	0.5	312	ns 69	12 <sup>+</sup>	06	2000	IT=100		
<sup>176</sup> Lu <sup>p</sup>	-51794.7	1.6	1587.5	1.1	40	$\mu$ s 3	14 <sup>+</sup>	06 FGK128	J	2000	IT=100	
<sup>176</sup> Hf	-54576.3	1.5	STABLE			0 <sup>+</sup>	06	1934	IS=5.26 7			
<sup>176</sup> Hf <sup>m</sup>	-53243.2	1.5	1333.07	0.07	9.6	$\mu$ s 0.3	6 <sup>+</sup>	06	1964	IT=100		
<sup>176</sup> Hf <sup>n</sup>	-53017.0	1.5	1559.31	0.09	9.9	$\mu$ s 0.2	8 <sup>-</sup>	06	1967	IT=100		
<sup>176</sup> Hf <sup>p</sup>	-51710.5	1.7	2865.8	0.7	401	$\mu$ s 6	14 <sup>-</sup>	06	1975	IT=100		
<sup>176</sup> Hf <sup>q</sup>	-49712.8	2.2	4863.5	1.6	43	$\mu$ s 4	22 <sup>-</sup>	06 10Mu13	JT	1976	IT=100	
<sup>176</sup> Ta	-51370	30	8.09	h	0.05	(1 <sup>-</sup> )	06	1948	$\beta^+$ =100			
<sup>176</sup> Ta <sup>m</sup>	-51270	30	103.0	1.0	1.08	ms 0.07	(7 <sup>+</sup> )	06 78Du06	ET	1971	IT=100	
<sup>176</sup> Ta <sup>n</sup>	-49900	30	1474.0	1.4	3.8	$\mu$ s 0.4	14 <sup>-</sup>	06	1978	IT=100		
<sup>176</sup> Ta <sup>p</sup>	-48500	30	2874.0	1.4	970	$\mu$ s 70	20 <sup>-</sup>	06	1994	IT=100		
<sup>176</sup> W	-50642	28	2.5	h	0.1	0 <sup>+</sup>	06	1950	$\epsilon$ =100			
<sup>176</sup> Re	-45063	28	5.3	m	0.3	(3 <sup>+</sup> )	06	1967	$\beta^+$ =100			
<sup>176</sup> Os	-42098	28	3.6	m	0.5	0 <sup>+</sup>	06	1970	$\beta^+$ =100			
<sup>176</sup> Ir	-33878	17	8.7	s	0.5	06		1967	$\beta^+$ =96.9 6; $\alpha$ =3.1 6			
<sup>176</sup> Pt	-28934	13	6.33	s	0.15	0 <sup>+</sup>	06	1966	$\beta^+$ ?; $\alpha$ =40 2			
<sup>176</sup> Au	-18520	30	1.05	s	0.01	(3 <sup>-</sup> , 4 <sup>-</sup> )	06 14An10	J	1975	$\alpha$ =75 8; $\beta^+$ ?		
<sup>176</sup> Au <sup>m</sup>	-18380	30	139	13	AD *	860	ms 160	(7 <sup>+</sup> )	06 02Ro17	T	2002	$\alpha$ ?; $\beta^+$ ?

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)						
... A-group continued ...														
<sup>176</sup> Hg	-11785	11	20.3 ms	1.4	0 <sup>+</sup>	06	1983	$\alpha=90.9$ ; $\beta^+$ ?	*					
<sup>176</sup> Tl	580	80	6.2 ms	2.3	(3 <sup>-</sup> , 4 <sup>-</sup> , 5 <sup>-</sup> )	09	2004	$p\approx 100$ ; $\alpha$ ?; $\beta^+$ ?	*					
* <sup>176</sup> Lu	T : average 14Hu07=37.22(0.29) 13Ko20=36.40(0.35) 06Lu03=35.6(0.7)													
* <sup>176</sup> Lu	T : 03Ni11=36.77(0.75) 92Da03=37.3(0.5) 65Br25=36.8(6)													
* <sup>176</sup> Lu <sup>p</sup>	J : 73.0 $\gamma$ (E2) to 12 <sup>+</sup> state													
* <sup>176</sup> Ta <sup>m</sup>	T : average 78Du06=1.05(0.10) 71Go21=1.1(0.1) J : from 98Ko09													
* <sup>176</sup> Ta <sup>n</sup>	E : 1371(1) keV above <sup>176</sup> Ta <sup>m</sup>													
* <sup>176</sup> Ta <sup>p</sup>	E : 2771(1) keV above <sup>176</sup> Ta <sup>m</sup>													
* <sup>176</sup> Au	D : $\alpha=75.8$ as quoted in 14An10													
* <sup>176</sup> Au <sup>m</sup>	T : symmetrized from 840(+170-140) J : from $\alpha$ decay to <sup>172</sup> Ir <sup>m</sup>													
* <sup>176</sup> Hg	D : $\alpha$ symmetrized from 99Po09=94(+6-12)%													
* <sup>176</sup> Tl	T : symmetrized from 5.2(+3.0-1.4)													
<sup>177</sup> Er	-42860#	500#	3#	s (>300 ns)	1/2 <sup>-</sup> #	13	12Ku26	I	2012	$\beta^-$ ?				
<sup>177</sup> Tm	-47470#	300#	90	s	6 (7/2 <sup>-</sup> )	03			1989	$\beta^-$ =100				
<sup>177</sup> Yb	-50986.40	0.22	1.911	h	0.003	9/2 <sup>+</sup>	03	12F105	J	1945	$\beta^-$ =100			
<sup>177</sup> Yb <sup>m</sup>	-50654.9	0.4	331.5	0.3	6.41	s	0.02	1/2 <sup>-</sup>	03	12F105	J	1962	IT=100	
<sup>177</sup> Lu	-52383.8	1.2	6.6457	d	0.0026	7/2 <sup>+</sup>	03	12Ko24	T	1945	$\beta^-$ =100	*		
<sup>177</sup> Lu <sup>m</sup>	-52233.4	1.2	150.3967	0.0010	130	ns	3	9/2 <sup>-</sup>	03		1949	IT=100		
<sup>177</sup> Lu <sup>n</sup>	-51814.1	1.2	569.7068	0.0016	155	$\mu$ s	7	1/2 <sup>+</sup>	03		1965	IT=100		
<sup>177</sup> Lu <sup>p</sup>	-51413.6	1.2	970.1750	0.0024	160.44	d	0.06	23/2 <sup>-</sup>	03		1962	$\beta^-$ =78.6 8; IT=21.4 8		
<sup>177</sup> Lu <sup>q</sup>	-49612.2	1.4	2771.6	0.7	625	ns	62	33/2 <sup>+</sup>		04Dr06	ETJ	2004	IT=100	
<sup>177</sup> Lu <sup>r</sup>	-48853.5	1.4	3530.3	0.7	6	$\mu$ s	2	39/2 <sup>-</sup>	03	11Ko.A	T	2003	IT=100	
<sup>177</sup> Hf	-52880.6	1.4	STABLE		7/2 <sup>-</sup>	03					1934	IS=18.60 9		
<sup>177</sup> Hf <sup>m</sup>	-51565.1	1.4	1315.4504	0.0008	1.09	s	0.05	23/2 <sup>+</sup>	03		1966	IT=100		
<sup>177</sup> Hf <sup>n</sup>	-51538.2	1.4	1342.38	0.20	55.9	$\mu$ s	1.2	(19/2 <sup>-</sup> )	03		1976	IT=100		
<sup>177</sup> Hf <sup>p</sup>	-50140.6	1.4	2740.02	0.15	51.4	m	0.5	37/2 <sup>-</sup>	03		1971	IT=100		
<sup>177</sup> Ta	-51715	3	56.56	h	0.06	7/2 <sup>+</sup>	03				1948	$\beta^+$ =100		
<sup>177</sup> Ta <sup>m</sup>	-51642	3	73.36	0.15	410	ns	7	9/2 <sup>-</sup>	03		1973	IT=100		
<sup>177</sup> Ta <sup>n</sup>	-51529	3	186.15	0.06	3.62	$\mu$ s	0.10	5/2 <sup>-</sup>	03		1971	IT=100		
<sup>177</sup> Ta <sup>p</sup>	-50360	3	1355.01	0.19	5.31	$\mu$ s	0.25	21/2 <sup>-</sup>	03		1971	IT=100		
<sup>177</sup> Ta <sup>q</sup>	-47059	3	4656.3	0.5	133	$\mu$ s	4	49/2 <sup>-</sup>	03		1994	IT=100		
<sup>177</sup> W	-49702	28	132	m	2	1/2 <sup>-</sup>	03				1950	$\beta^+$ =100		
<sup>177</sup> Re	-46269	28	14	m	1	5/2 <sup>-</sup>	03				1957	$\beta^+$ =100		
<sup>177</sup> Re <sup>m</sup>	-46184	28	84.71	0.10	50	$\mu$ s	10	5/2 <sup>+</sup>	03		1972	IT=100		
<sup>177</sup> Os	-41956	15	3.0	m	0.2	1/2 <sup>-</sup>	03				1970	$\beta^+$ =100		
<sup>177</sup> Ir	-36047	20	30	s	2	5/2 <sup>-</sup>	03				1967	$\beta^+\approx 100$ ; $\alpha=0.06$ 1		
<sup>177</sup> Pt	-29370	15	10.6	s	0.4	5/2 <sup>-</sup>	03				1966	$\beta^+=94.3$ 5; $\alpha=5.7$ 5		
<sup>177</sup> Pt <sup>m</sup>	-29223	15	147.4	0.4	2.2	$\mu$ s	0.3	1/2 <sup>-</sup>	03		1979	IT=100		
<sup>177</sup> Au	-21545	10	1.46	s	0.03	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	03	01Ko44	TJ	1968	$\alpha=40$ 6; $\beta^+$ ?			
<sup>177</sup> Au <sup>m</sup>	-21356	10	189	8	AD	1.180	s	0.012	11/2 <sup>-</sup>	03	01Ko44	ETJ	1975	$\alpha=66$ 10; $\beta^+$ ?
<sup>177</sup> Hg	-12780	80	127.3	ms	1.8	(7/2 <sup>-</sup> )	03	05Ca43	J	1975	$\alpha=85$ ; $\beta^+=15$			
<sup>177</sup> Hg <sup>m</sup>	-12460	80	323	1	1.50	$\mu$ s	0.15	(13/2 <sup>+</sup> )	03	03Me20	ETJ	2003	IT=100	
<sup>177</sup> Tl	-3341	22	18	ms	5	(1/2 <sup>+</sup> )	03				1999	$\alpha=73$ 13; $p=27$ 13		
<sup>177</sup> Tl <sup>m</sup>	-2534	12	807	18	p	180	$\mu$ s	60	(11/2 <sup>-</sup> )	03	04Ke06	TD	1997	$p=51$ 8; $\alpha=49$ 8
* <sup>177</sup> Lu	T : average 12Ko24=6.639(0.009) 11Po07=6.6465(0.0032) 01Sc23=6.646(0.005)													
* <sup>177</sup> Lu <sup>r</sup>	E : derived by NUBASE from least-squares fit to $\gamma$ -ray energies													
* <sup>177</sup> Lu <sup>r</sup>	T : 04Al04=7(2) m, not trusted													
* <sup>177</sup> Hf <sup>p</sup>	T : other 04Al04=76(+16-9) from decay growth													
* <sup>177</sup> Au	T : average 09An14=1.53(0.07) 01Ko44=1.46(0.03) D : from 09An14													
* <sup>177</sup> Au <sup>m</sup>	D : from 09An14													
* <sup>177</sup> Hg	J : also 09An20													
* <sup>177</sup> Tl <sup>m</sup>	T : 04Ke06=160(+70-40) D : also 04Ke06=55(20)%													
<sup>178</sup> Er	-40260#	600#	1#	s (>300 ns)	0 <sup>+</sup>	13	12Ku26	I	2012	$\beta^-$ ?; $\beta^-$ n=0#				
<sup>178</sup> Tm	-44120#	400#	30#	s (>300 ns)		11	09St16	I	2008	$\beta^-$ ?				
<sup>178</sup> Yb	-49695	10	74	m	3	0 <sup>+</sup>	09				1973	$\beta^-$ =100		
<sup>178</sup> Lu	-50337.8	2.3	28.4	m	0.2	1(+)	09				1957	$\beta^-$ =100		
<sup>178</sup> Lu <sup>m</sup>	-50214	3	123.8	2.6	RQ	23.1	m	0.3	9(-)	09	98Ge13	J	1951	$\beta^-$ =100
<sup>178</sup> Hf	-52435.2	1.4	STABLE		0 <sup>+</sup>	09					1934	IS=27.28 7		
<sup>178</sup> Hf <sup>m</sup>	-51287.8	1.4	1147.416	0.006	4.0	s	0.2	8 <sup>-</sup>	09		1960	IT=100		
<sup>178</sup> Hf <sup>n</sup>	-49989.1	1.4	2446.09	0.08	31	y	1	16 <sup>+</sup>	09		1968	IT=100		
<sup>178</sup> Hf <sup>p</sup>	-49862.8	1.4	2572.4	0.3	68	$\mu$ s	2	14 <sup>-</sup>	09		1977	IT=100		
<sup>178</sup> Ta	-50600#	50#	2.36	h	0.08	7 <sup>-</sup> #	09				1950	$\beta^+$ =100		
<sup>178</sup> Ta <sup>m</sup>	-50498	15	100#	50#	*	9.31	m	0.03	1 <sup>+</sup> #	09	96Ko13	E	1950	$\beta^+$ =100
<sup>178</sup> Ta <sup>n</sup>	-49130#	50#	1467.82	0.16	59	ms	3	15 <sup>-</sup>	09	96Ko13	ETJ	1979	IT=100	
<sup>178</sup> Ta <sup>p</sup>	-47700#	50#	2901.9	0.7	290	ms	12	21 <sup>-</sup>	09	96Ko13	ETJ	1996	IT=100	
... A-group is continued on next page ...														



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>178</sup> W	-50407	15	21.6 d 0.3	0 <sup>+</sup>	09		1950	$\epsilon=100$	
<sup>178</sup> W <sup>m</sup>	-43834	15	6572.7 0.3	220 ns 10	25 <sup>+</sup>	09	1998	IT=100	
<sup>178</sup> Re	-45653	28	13.2 m 0.2	(3 <sup>+</sup> )	09		1957	$\beta^+=100$	
<sup>178</sup> Os	-43544	14	5.0 m 0.4	0 <sup>+</sup>	09		1967	$\beta^+=100$	
<sup>178</sup> Ir	-36252	20	12 s 2		09		1972	$\beta^+=100$	
<sup>178</sup> Pt	-31998	10	20.7 s 0.7	0 <sup>+</sup>	09		1966	$\beta^+=92.3\ 3; \alpha=7.7\ 3$	
<sup>178</sup> Au	-22304	10	2.6 s 0.5		09		1968	$\beta^+ < 60; \alpha > 40$	
<sup>178</sup> Au <sup>m</sup>	-22115	10	189 14	> 1 s		15Ma.A ET	2015	$\beta^+ ?; \alpha ?$	
<sup>178</sup> Au <sup>p</sup>	-21939	24	365 21	AD					
<sup>178</sup> Hg	-16316	11	266.5 ms 2.4	0 <sup>+</sup>	09	12Ve04 D	1971	$\alpha=89\ 4; \beta^+ ?$	
<sup>178</sup> Tl	-4790#	90#	255 ms 9	(4 <sup>-</sup> , 5 <sup>-</sup> )	09	13Li49 TJD	1997	$\alpha=62\ 2; \beta^+=38\ 2; \beta^+SF=0.15\ 6$	
<sup>178</sup> Pb	3574	24	230 $\mu$ s 150	0 <sup>+</sup>	09	01Ro.B T	2001	$\alpha \approx 100; \beta^+ ?$	
* <sup>178</sup> Ta <sup>m</sup>	E: 1 <sup>+</sup> state (p9/2 <sup>-</sup> [514]+n7/2 <sup>-</sup> [514]) is expected 104 keV above the 7 <sup>-</sup> ground-state, **								
* <sup>178</sup> Ta <sup>m</sup>	E: based on E=220 keV for 8 <sup>+</sup> (p9/2 <sup>-</sup> [514]+n7/2 <sup>-</sup> [514]) and residual energy **								
* <sup>178</sup> Ta <sup>m</sup>	E: shift of 50 keV from known Gallagher-Moszkowski splitting energy **								
* <sup>178</sup> Ta <sup>n</sup>	E: from least-squares fit to $\gamma$ -rays in 96Ko13 **								
* <sup>178</sup> Ta <sup>n</sup>	T: average 96Ko13=58(4) 79Du02=60(5) **								
* <sup>178</sup> Ta <sup>p</sup>	E: from least-squares fit to $\gamma$ -rays in 96Ko13 **								
* <sup>178</sup> Tl	T: average 13Li49=252(20) 02Ro17=254(+11-9) **								
* <sup>178</sup> Pb	T: two events at 202 and 147 $\mu$ s, see 84Sc13 **								
<sup>179</sup> Tm	-41600#	500#	20# s (>300 ns)	1/2 <sup>+</sup> #	13	12Ku26 I	2012	$\beta^- ?; \beta^-n=0\#$	
<sup>179</sup> Yb	-46540#	200#	8.0 m 0.4	(1/2 <sup>-</sup> )	09		1982	$\beta^- =100$	
<sup>179</sup> Lu	-49059	5	4.59 h 0.06	7/2 <sup>+</sup>	09		1961	$\beta^- =100$	
<sup>179</sup> Lu <sup>m</sup>	-48467	5	592.4 0.4	1/2 <sup>+</sup>	09		1982	IT=100	
<sup>179</sup> Hf	-50462.9	1.4	STABLE	9/2 <sup>+</sup>	09		1934	IS=13.62 2	
<sup>179</sup> Hf <sup>m</sup>	-50087.9	1.4	375.0352 0.0025	1/2 <sup>-</sup>	09		1962	IT=100	
<sup>179</sup> Hf <sup>n</sup>	-49357.2	1.4	1105.74 0.16	25.05 d 0.25	25/2 <sup>-</sup>	09	1970	IT=100	
<sup>179</sup> Hf <sup>p</sup>	-46687.7	2.5	3775.2 2.1	15 $\mu$ s 5	(43/2 <sup>+</sup> )	09	2000	IT=100	
<sup>179</sup> Ta	-50357.3	1.5	1.82 y 0.03	7/2 <sup>+</sup>	09		1950	$\epsilon=100$	
<sup>179</sup> Ta <sup>m</sup>	-50326.6	1.5	30.7 0.1	1.42 $\mu$ s 0.08	9/2 <sup>-</sup>	09	1964	IT=100	
<sup>179</sup> Ta <sup>n</sup>	-49837.1	1.5	520.23 0.18	280 ns 80	1/2 <sup>+</sup>	09	FGK128 J	1974	IT=100
<sup>179</sup> Ta <sup>p</sup>	-49104.7	1.5	1252.60 0.23	322 ns 16	21/2 <sup>-</sup>	09	97Ko13 J	1982	IT=100
<sup>179</sup> Ta <sup>q</sup>	-49040.1	1.6	1317.2 0.4	9.0 ms 0.2	25/2 <sup>+</sup>	09	97Ko13 J	1982	IT=100
<sup>179</sup> Ta <sup>r</sup>	-49029.3	1.6	1328.0 0.4	1.6 $\mu$ s 0.4	23/2 <sup>-</sup>	09	97Ko13 J	1982	IT=100
<sup>179</sup> Ta <sup>x</sup>	-47718.0	1.6	2639.3 0.5	54.1 ms 1.7	37/2 <sup>+</sup>	09	97Ko13 J	1982	IT=100
<sup>179</sup> W	-49295	15	37.05 m 0.16	7/2 <sup>-</sup>	09		1950	$\beta^+=100$	
<sup>179</sup> W <sup>m</sup>	-49073	15	221.91 0.03	6.40 m 0.07	1/2 <sup>-</sup>	09	1950	IT $\approx$ 100; $\beta^+=0.29\ 4$	
<sup>179</sup> W <sup>n</sup>	-47663	15	1631.90 0.08	390 ns 30	21/2 <sup>+</sup>	09	94Wa05 J	1978	IT=100
<sup>179</sup> W <sup>p</sup>	-45947	15	3348.41 0.14	750 ns 80	35/2 <sup>-</sup>	09	94Wa05 J	1978	IT=100
<sup>179</sup> Re	-46584	25	19.5 m 0.1	5/2 <sup>+</sup>	09		1960	$\beta^+=100$	
<sup>179</sup> Re <sup>m</sup>	-46519	25	65.35 0.09	95 $\mu$ s 25	(5/2 <sup>-</sup> )	09	1972	IT=100	
<sup>179</sup> Re <sup>n</sup>	-44760	60	1822 50	408 ns 12	(23/2 <sup>+</sup> )	09	1972	IT=100	
<sup>179</sup> Re <sup>p</sup>	-41176	25	5408.0 0.5	466 $\mu$ s 15	(47/2 <sup>+</sup> , 49/2 <sup>+</sup> )	09	1989	IT=100	
<sup>179</sup> Os	-43019	17	6.5 m 0.3	1/2 <sup>-</sup>	09		1968	$\beta^+=100$	
<sup>179</sup> Os <sup>m</sup>	-42874	17	145.41 0.12	500 ns	(7/2 <sup>-</sup> )	09	1983	IT=100	
<sup>179</sup> Os <sup>n</sup>	-42776	17	243.0 0.8	783 ns 14	(9/2 <sup>+</sup> )	09	1983	IT=100	
<sup>179</sup> Ir	-38082	10	79 s 1	(5/2 <sup>-</sup> )	09		1992	$\beta^+=100$	
<sup>179</sup> Pt	-32268	8	21.2 s 0.4	1/2 <sup>-</sup>	09		1966	$\beta^+ \approx 100; \alpha = 0.24\ 3$	
<sup>179</sup> Au	-24989	12	7.1 s 0.3	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	09		1968	$\beta^+ = 78.0\ 9; \alpha = 22.0\ 9$	
<sup>179</sup> Au <sup>m</sup>	-24900	12	89.5 0.5	328 ns 2	(3/2 <sup>-</sup> )	09	11Ve01 ETD	2011	IT=100
<sup>179</sup> Hg	-16928	27	1.05 s 0.03	7/2 <sup>-</sup>	09	12Ve04 D	1970	$\alpha=75\ 4; \beta^+ ?; \beta^+p \approx 0.15$	
<sup>179</sup> Hg <sup>m</sup>	-16757	27	171.4 0.4	6.4 $\mu$ s 0.9	13/2 <sup>+</sup>	09	02Je09 J	2002	IT=100
<sup>179</sup> Tl	-8270	40	265 ms 10	1/2 <sup>+</sup>	09	13An10 TJD	1983	$\alpha=60\ 2; \beta^+ ?$	
<sup>179</sup> Tl <sup>m</sup>	-7440#	40#	825# 10#	1.41 ms 0.03	(11/2 <sup>-</sup> )	09	11Ko.B TJ	1983	$\alpha \approx 100; IT ?; \beta^+ ?$
<sup>179</sup> Pb	2050	80	3.9 ms 1.1	(9/2 <sup>-</sup> )	10	10An01 TDJ	2010	$\alpha=100$	
* <sup>179</sup> Re <sup>n</sup>	E: x keV above 1772.20(0.22) level; x estimated 50(50) by NUBASE **								
* <sup>179</sup> Au <sup>m</sup>	E: uncertainty estimated by NUBASE **								
* <sup>179</sup> Au <sup>p</sup>	E: 44(15) above 89.5 keV level **								
* <sup>179</sup> Tl	T: others 11Ko.B=489(21) 02Ro17=415(55) **								
* <sup>179</sup> Tl	J: $\alpha$ decay to 1/2 <sup>+</sup> in <sup>175</sup> Au **								
* <sup>179</sup> Tl <sup>m</sup>	J: from $\alpha$ decay to <sup>175</sup> Au <sup>m</sup> E: estimated from TNN in <sup>177,181,183</sup> Tl **								
* <sup>179</sup> Tl <sup>m</sup>	T: average 11Ko.B=1.36(0.04) 10An01=1.46(0.04) **								
* <sup>179</sup> Pb	T: symmetrized from 3.5(+1.4-0.8) **								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>180</sup> Tm	-37920#	500#	5# s (>300 ns)		15		2012	$\beta^-$ ?; $\beta^-$ -n=0#		
<sup>180</sup> Yb	-44600#	300#	2.4 m 0.5	0 <sup>+</sup>	15		1987	$\beta^-$ =100		
<sup>180</sup> Lu	-46680	70	5.7 m 0.1	5 <sup>+</sup>	15		1971	$\beta^-$ =100		
<sup>180</sup> Lu <sup>m</sup>	-46670	70	1 s	3 <sup>-</sup>	15	95Me03 JT	1995	$\beta^-$ ?; IT ?		
<sup>180</sup> Lu <sup>n</sup>	-46060	70	624.0	0.5			2001	IT=100		
<sup>180</sup> Hf	-49779.3	1.4	STABLE		0 <sup>+</sup>		1934	IS=35.08 16		
<sup>180</sup> Hf <sup>m</sup>	-48637.7	1.4	1141.552	0.015	5.53 h 0.02		1951	IT≈100; $\beta^-$ =0.31 8		
<sup>180</sup> Hf <sup>n</sup>	-48404.9	1.4	1374.36	0.04	570 $\mu$ s 20		1990	IT=100		
<sup>180</sup> Hf <sup>p</sup>	-47293.8	1.5	2485.5	0.5	940 ns 110		2000	IT=100		
<sup>180</sup> Hf <sup>q</sup>	-46181.8	1.7	3597.5	1.0	90 $\mu$ s 10		1999	IT=100		
<sup>180</sup> Ta	-48932.9	1.9	8.154 h 0.006		1 <sup>+</sup>		1938	$\epsilon$ =85 3; $\beta^-$ =15 3		
<sup>180</sup> Ta <sup>m</sup>	-48857.5	1.4	75.3	1.4	RQ STABLE (>7.1 Py)		1940	IS=0.01201 32; $\beta^-$ ?		
<sup>180</sup> Ta <sup>n</sup>	-47480.5	1.9	1452.39	0.22	31.2 $\mu$ s 1.4		1996	IT=100		
<sup>180</sup> Ta <sup>p</sup>	-45254.0	2.1	3678.9	1.0	2.0 $\mu$ s 0.5		2000	IT=100		
<sup>180</sup> Ta <sup>q</sup>	-44760.7	2.5	4172.2	1.6	17 $\mu$ s 5		2000	IT=100		
<sup>180</sup> W	-49636.1	1.4	1.8 Ey 0.2		0 <sup>+</sup>		1937	IS=0.12 1; $\alpha$ ≈100; 2 $\beta^+$ ?		
<sup>180</sup> W <sup>m</sup>	-48107.0	1.4	1529.05	0.04	5.47 ms 0.09		1978	IT=100		
<sup>180</sup> W <sup>n</sup>	-46371.2	1.4	3264.9	0.3	2.3 $\mu$ s 0.2		1966	IT=100		
<sup>180</sup> Re	-45837	21	2.46 m 0.03		(1) <sup>-</sup>		1955	$\beta^+$ =100		
<sup>180</sup> Re <sup>m</sup>	-45750#	40#	90#	30#	> 1 $\mu$ s	(4 <sup>+</sup> , 5 <sup>+</sup> )	05E110 J	2005	IT≈100; $\beta^+$ ?	
<sup>180</sup> Re <sup>n</sup>	-42280#	40#	3561#	30#	9.0 $\mu$ s 0.7	(21) <sup>-</sup>	15 05E110 TJD	2005	IT=100	
<sup>180</sup> Os	-44358	16	21.5 m 0.4		0 <sup>+</sup>		1967	$\beta^+$ =100		
<sup>180</sup> Ir	-37978	22	1.5 m 0.1		(5 <sup>+</sup> )		1972	$\beta^+$ =100		
<sup>180</sup> Pt	-34436	11	56 s 3		0 <sup>+</sup>		1966	$\beta^+$ ≈100; $\alpha$ ≈0.3		
<sup>180</sup> Au	-25626	5	8.4 s 0.6		15		1977	$\beta^+$ <98.2; $\alpha$ >1.8		
<sup>180</sup> Hg	-20250	13	2.59 s 0.01		0 <sup>+</sup>		1970	$\beta^+$ =52 2; $\alpha$ =48 2		
<sup>180</sup> Tl	-9390	60	1.09 s 0.01		4 <sup>(-)</sup>	15 12Bi.A J	1987	$\beta^+$ =94 4; $\alpha$ =6 4; $\beta^+$ SF=0.0032 2		
<sup>180</sup> Pb	-1941	12	4.1 ms 0.3		0 <sup>+</sup>		1996	$\alpha$ =100		
* <sup>180</sup> Hf <sup>n</sup>	I : isomer at 2425.8(1.0) 15(5) $\mu$ s (10 <sup>+</sup> ) reported then retracted by authors									
* <sup>180</sup> W	T : 03Da09>80 Py for 2 $\beta^-$ decay									
* <sup>180</sup> Re <sup>n</sup>	E : 3471.8(0.6) above (5 <sup>+</sup> ) level, most likely isomer, estimated to be 90#30 keV									
<sup>181</sup> Tm	-35170#	600#	5# s (>300 ns)	1/2 <sup>+</sup> #	13	12Ku26 I	2012	$\beta^-$ ?; $\beta^-$ -n=0.4#		
<sup>181</sup> Yb	-41090#	300#	1# m (>300 ns)	3/2 <sup>+</sup> #	13	09St16 I	2000	$\beta^-$ ?		
<sup>181</sup> Lu	-44800	130	3.5 m 0.3	7/2 <sup>+</sup> #	06		1982	$\beta^-$ =100		
<sup>181</sup> Hf	-47402.8	1.4	42.39 d 0.06	1/2 <sup>-</sup>	06		1935	$\beta^-$ =100		
<sup>181</sup> Hf <sup>m</sup>	-46807.5	1.4	595.27	0.04	80 $\mu$ s 5	01Sh36 T	2001	IT=100		
<sup>181</sup> Hf <sup>n</sup>	-46359.3	1.6	1043.5	0.8	100 $\mu$ s	(17/2 <sup>+</sup> )	06	2001	IT=100	
<sup>181</sup> Hf <sup>p</sup>	-45660.9	1.9	1741.9	1.3	1.5 ms 0.5	(25/2 <sup>-</sup> )	06	2001	IT=100	
<sup>181</sup> Ta	-48438.3	1.4	STABLE		7/2 <sup>+</sup>		1932	IS=99.98799 32		
<sup>181</sup> Ta <sup>m</sup>	-48432.1	1.4	6.237	0.020	6.05 $\mu$ s 0.12	9/2 <sup>-</sup>	06	1979	IT=100	
<sup>181</sup> Ta <sup>n</sup>	-47823.1	1.4	615.19	0.03	18 $\mu$ s 1	1/2 <sup>+</sup>	06	1948	IT=100	
<sup>181</sup> Ta <sup>p</sup>	-47010	14	1428	14	140 ns 36	(19/2 <sup>+</sup> )	06	1998	IT=100	
<sup>181</sup> Ta <sup>q</sup>	-46954.9	1.4	1483.43	0.21	25.2 $\mu$ s 1.8	21/2 <sup>-</sup>	06	98Wh02 T	1998	IT=100
<sup>181</sup> Ta <sup>r</sup>	-46210.4	1.7	2227.9	0.9	210 $\mu$ s 20	29/2 <sup>-</sup>	06	98Wh02 J	1998	IT=100
<sup>181</sup> W	-48233.8	1.4	121.2 d 0.2		9/2 <sup>+</sup>		06	1947	$\epsilon$ =100	
<sup>181</sup> W <sup>m</sup>	-47868.3	1.4	365.55	0.13	14.59 $\mu$ s 0.15	5/2 <sup>-</sup>	06	1968	IT=100	
<sup>181</sup> W <sup>n</sup>	-46580.7	1.5	1653.1	0.6	140 ns 20	21/2 <sup>+</sup>	06	1973	IT=100	
<sup>181</sup> Re	-46517	13	19.9 h 0.7		5/2 <sup>+</sup>		06	1957	$\beta^+$ =100	
<sup>181</sup> Re <sup>m</sup>	-46254	13	262.91	0.11	156.7 ns 1.9	9/2 <sup>-</sup>	06	1967	IT=100	
<sup>181</sup> Re <sup>n</sup>	-44861	13	1656.37	0.14	250 ns 10	21/2 <sup>-</sup>	06	1974	IT=100	
<sup>181</sup> Re <sup>p</sup>	-44636	13	1880.57	0.16	11.5 $\mu$ s 0.9	25/2 <sup>+</sup>	06	2000	IT=100	
<sup>181</sup> Re <sup>q</sup>	-42648	13	3869.40	0.18	1.2 $\mu$ s 0.2	(35/2 <sup>-</sup> )	06	2000	IT=100	
<sup>181</sup> Os	-43550	25	105 m 3		1/2 <sup>-</sup>		06	1966	$\beta^+$ =100	
<sup>181</sup> Os <sup>m</sup>	-43501	25	49.20	0.14	2.7 m 0.1	7/2 <sup>-</sup>	06	1966	$\beta^+$ =100	
<sup>181</sup> Os <sup>n</sup>	-43393	25	156.91	0.15	262 ns 6	9/2 <sup>+</sup>	06	1974	IT=100	
<sup>181</sup> Ir	-39463	5	4.90 m 0.15		5/2 <sup>-</sup>		06	1972	$\beta^+$ =100	
<sup>181</sup> Ir <sup>m</sup>	-39174	5	289.33	0.13	298 ns	5/2 <sup>+</sup>	06	1992	IT=100	
<sup>181</sup> Ir <sup>n</sup>	-39097	5	366.30	0.22	126 ns 6	9/2 <sup>-</sup>	06	1992	IT=100	
<sup>181</sup> Pt	-34382	14	52.0 s 2.2		1/2 <sup>-</sup>	06	95Bi01 D	1966	$\beta^+$ ≈100; $\alpha$ =0.074 10	
<sup>181</sup> Pt <sup>m</sup>	-34265	14	116.65	0.08	> 300 ns	(7/2) <sup>-</sup>	06	1992	IT=100	
<sup>181</sup> Au	-27871	20	13.7 s 1.4		(3/2 <sup>-</sup> )		06	1968	$\beta^+$ ?; $\alpha$ =2.7 5	
<sup>181</sup> Hg	-20661	15	3.6 s 0.1		1/2 <sup>(-#)</sup>		06	1969	$\beta^+$ =73 2; $\alpha$ =27 2; $\beta^+$ p=0.013 3; $\beta^+$ $\alpha$ =9e-6 6	
<sup>181</sup> Hg <sup>m</sup>	-20450	50	210	50	480 $\mu$ s 20	13/2 <sup>+</sup>	06	09An17 T	2009	IT ?

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>181</sup> Tl	-12799	9	3.2 s	0.3	1/2 <sup>+</sup>	09 09An14	J	1996	$\beta^+ ?; \alpha < 10$	*	
<sup>181</sup> Tl <sup>m</sup>	-11963	9	835.9	0.4	1.40 ms	0.03	(9/2 <sup>-</sup> )	09 09An14	J	1984	$\Gamma T=99.60$ 4; $\alpha=0.40$ 6; $\beta^+ ?$
<sup>181</sup> Pb	-3120	80	39.0 ms	0.8	(9/2 <sup>-</sup> )	06 09An20	TJ	1989	$\alpha=?; \beta^+=2\#$	*	
<sup>181</sup> Pb <sup>m</sup>	non existent		RN		13/2 <sup>+</sup> #	96To01	I				
* <sup>181</sup> Ta <sup>p</sup>	E: x keV above 1403.2(0.6) level; x<50									**	
* <sup>181</sup> Ta <sup>q</sup>	T: average 98Wh02=25(2) 98Dr09=23(+6-2)									**	
* <sup>181</sup> Tl	T: average 98To14=3.2(0.3) 92Bo.D=3.4(0.6)									**	
* <sup>181</sup> Pb	T: average 09An20=36(2) 05Ca.A=39.6(0.9)									**	
<sup>182</sup> Yb	-38820#	400#	10#	s	(>300 ns)	0 <sup>+</sup>	15 12Ku26	I	2012	$\beta^- ?$	
<sup>182</sup> Lu	-41880#	200#	2.0	m	0.2	1 <sup>-</sup> #	15		1982	$\beta^- =100$	
<sup>182</sup> Hf	-46050	6	8.90	My	0.09	0 <sup>+</sup>	15		1961	$\beta^- =100$	
<sup>182</sup> Hf <sup>m</sup>	-44877	6	1172.87	0.18	61.5 m	1.5	(8) <sup>-</sup>	15 FGK128	J	1971	$\beta^- =54$ 2; $\Gamma T=46$ 2
<sup>182</sup> Hf <sup>m</sup>	-43479	6	2571.3	1.2	40 $\mu$ s	10	(13 <sup>+</sup> )	15		1999	$\Gamma T=100$
<sup>182</sup> Ta	-46429.9	1.4	114.74	d	0.12	3 <sup>-</sup>	15		1938	$\beta^- =100$	
<sup>182</sup> Ta <sup>m</sup>	-46413.6	1.4	16.273	0.004	283 ms	3	5 <sup>+</sup>	15		1968	$\Gamma T=100$
<sup>182</sup> Ta <sup>n</sup>	-45910.3	1.4	519.577	0.016	15.84 m	0.10	10 <sup>-</sup>	15		1947	$\Gamma T=100$
<sup>182</sup> W	-48246.1	0.7	STABLE		(>7.7 Zy)	0 <sup>+</sup>	15		1930	$IS=26.50$ 16; $\alpha ?$	
<sup>182</sup> W <sup>m</sup>	-46015.5	0.7	2230.65	0.14	1.3 $\mu$ s	0.1	(10 <sup>+</sup> )	15		1969	$\Gamma T=100$
<sup>182</sup> Re	-45450	100	64.2	h	0.5	7 <sup>+</sup>	15		1950	$\beta^+ =100$	
<sup>182</sup> Re <sup>m</sup>	-45386	20	60	100	BD *	14.14 h	0.45	2 <sup>+</sup>	15	1950	$\beta^+ =100$
<sup>182</sup> Re <sup>n</sup>	-45150	140	300	100		585 ns	30	(2) <sup>-</sup>	15	1969	$\Gamma T=100$
<sup>182</sup> Re <sup>p</sup>	-44930	140	520	100		780 ns	90	(4) <sup>-</sup>	15	1984	$\Gamma T=100$
<sup>182</sup> Os	-44609	22	21.84	h	0.20	0 <sup>+</sup>	15		1950	$\epsilon=100$	
<sup>182</sup> Os <sup>m</sup>	-42778	22	1831.4	0.3	780 $\mu$ s	70	(8) <sup>-</sup>	15		1966	$\Gamma T=100$
<sup>182</sup> Os <sup>n</sup>	-37560	22	7049.5	0.4	150 ns	10	(25 <sup>+</sup> )	15		1988	$\Gamma T=100$
<sup>182</sup> Ir	-39052	21	15.0	m	1.0	3 <sup>+</sup>	15		1961	$\beta^+ =100$	
<sup>182</sup> Ir <sup>m</sup>	-38981	21	71.02	0.17	170 ns	40	(5 <sup>+</sup> )	15		1990	$\Gamma T=100$
<sup>182</sup> Ir <sup>n</sup>	-38876	21	176.4	0.3	130 ns	50	(6 <sup>-</sup> )	15		1990	$\Gamma T=100$
<sup>182</sup> Pt	-36168	13	2.67	m	0.12	0 <sup>+</sup>	15		1963	$\beta^+ \approx 100; \alpha=0.038$ 2	
<sup>182</sup> Au	-28301	20	15.5	s	0.4	(2 <sup>+</sup> )	15		1970	$\beta^+ \approx 100; \alpha=0.13$ 5	
<sup>182</sup> Au <sup>p</sup>	-28180	30	120	40		high					
<sup>182</sup> Hg	-23577	10	10.83	s	0.06	0 <sup>+</sup>	15	71Ho07	D	1968	$\beta^+ =86.2$ 9; $\alpha=13.8$ 9; $\beta^+ p < 1e-5$
<sup>182</sup> Tl	-13328	12	1.9	s	0.1	(2 <sup>-</sup> )	10	16Va01	TJD	1991	$\beta^+ \approx 100; \alpha < 0.49; \beta^+ SF < 3.4e-6$
<sup>182</sup> Tl <sup>m</sup>	-13280#	50#	50#	50#	*	3#	s	(7 <sup>+</sup> )	91Bo22	J	
<sup>182</sup> Tl <sup>p</sup>	-12830#	100#	500#	100#	*			(10 <sup>-</sup> )			
<sup>182</sup> Pb	-6825	12	55	ms	5	0 <sup>+</sup>	15		1986	$\alpha=?; \beta^+=2\#$	
* <sup>182</sup> Hf <sup>m</sup>	J: E1 to 8 <sup>+</sup>									**	
* <sup>182</sup> Re <sup>m</sup>	E: 235.732(0.022) above <sup>182</sup> Re <sup>m</sup>									**	
* <sup>182</sup> Re <sup>p</sup>	E: 461.3(0.1) above <sup>182</sup> Re <sup>m</sup>									**	
<sup>183</sup> Yb	-35100#	400#	3#	s	(>300 ns)	3/2 <sup>-</sup> #	16 12Ku26	I	2012	$\beta^- ?$	
<sup>183</sup> Lu	-39720	80	58	s	4	(7/2 <sup>+</sup> )	16		1983	$\beta^- =100$	
<sup>183</sup> Hf	-43280	30	1.018	h	0.002	(3/2 <sup>-</sup> )	16		1956	$\beta^- =100$	
<sup>183</sup> Hf <sup>m</sup>	-41820	70	1464	64	40 s	30	27/2 <sup>-</sup> #	16 10Re07	ETJ	2010	$\Gamma T < 100; \beta^- ?$
<sup>183</sup> Ta	-45292.8	1.4	5.1	d	0.1	7/2 <sup>+</sup>	16		1950	$\beta^- =100$	
<sup>183</sup> Ta <sup>m</sup>	-45219.6	1.4	73.164	0.014	106 ns	10	(9/2) <sup>-</sup>	16		1967	$\Gamma T=100$
<sup>183</sup> Ta <sup>n</sup>	-43957	15	1336	15	900 ns	300	(19/2 <sup>+</sup> )	16 09Sh17	ETJ	2009	$\Gamma T=100$
<sup>183</sup> W	-46365.6	0.7	STABLE		(>670 Ey)	1/2 <sup>-</sup>	16		1930	$IS=14.31$ 4; $\alpha ?$	
<sup>183</sup> W <sup>m</sup>	-46056.1	0.7	309.492	0.004	5.30 s	0.08	11/2 <sup>+</sup>	16		1961	$\Gamma T=100$
<sup>183</sup> Re	-45810	8	70.0	d	1.4	5/2 <sup>+</sup>	16		1950	$\epsilon=100$	
<sup>183</sup> Re <sup>m</sup>	-43903	8	1907.21	0.15	1.04 ms	0.04	(25/2 <sup>+</sup> )	16		1966	$\Gamma T=100$
<sup>183</sup> Os	-43660	50	13.0	h	0.5	9/2 <sup>+</sup>	16		1950	$\beta^+ =100$	
<sup>183</sup> Os <sup>m</sup>	-43490	50	170.73	0.07	9.9 h	0.3	1/2 <sup>-</sup>	16			$\beta^+ =85$ 2; $\Gamma T=15$ 2
<sup>183</sup> Ir	-40203	24	58	m	5	5/2 <sup>-</sup>	16	61Di04	T	1961	$\beta^+ \approx 100; \alpha=0.05\#$
<sup>183</sup> Pt	-35772	16	6.5	m	1.0	1/2 <sup>-</sup>	16		1963	$\beta^+ \approx 100; \alpha=0.0096$ 5	
<sup>183</sup> Pt <sup>m</sup>	-35737	16	34.74	0.07	43 s	5	7/2 <sup>-</sup>	16		1979	$\beta^+ \approx 100; \Gamma T=3.1$ 8; $\alpha < 3e-4$
<sup>183</sup> Pt <sup>n</sup>	-35576	16	195.90	0.10	> 150 ns		(9/2) <sup>+</sup>	16		1990	$\Gamma T=100$
<sup>183</sup> Au	-30191	9	42.8	s	1.0	5/2 <sup>-</sup>	16	94Pa37	J	1968	$\beta^+ \approx 100; \alpha=0.55$ 25
<sup>183</sup> Au <sup>m</sup>	-30118	9	> 1 $\mu$ s			(1/2) <sup>+</sup>	16		1984	$\Gamma T=100$	
<sup>183</sup> Au <sup>p</sup>	-29960	9	< 1 $\mu$ s			(11/2) <sup>-</sup>	16		1984	$\Gamma T=100$	
<sup>183</sup> Hg	-23805	7	9.4	s	0.7	1/2 <sup>-</sup>	16		1969	$\beta^+ =88.3$ 20; $\alpha=11.7$ 20; $\beta^+ p=2.6e-4$ 8	
<sup>183</sup> Hg <sup>m</sup>	-23601	13	204	14	AD	> 8# $\mu$ s	13/2 <sup>+</sup> #	81Mi12	I		$\beta^+ ?$
<sup>183</sup> Tl	-16587	9	6.9	s	0.7	1/2 <sup>(+)</sup>	16	13Ba41	J	1980	$\beta^+ =?; \alpha=2\#$
<sup>183</sup> Tl <sup>m</sup>	-15959	9	628.7	0.5	53.3 ms	0.3	(9/2) <sup>-</sup>	16		1980	$\Gamma T=?; \alpha=1.5$ 3; $\beta^+ ?$
<sup>183</sup> Tl <sup>n</sup>	-15612	9	975.01	0.23	1.48 $\mu$ s	0.10	(13/2 <sup>+</sup> )	16		2001	$\Gamma T=100$
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>183</sup> Pb	-7575	28	535 ms	30	3/2 <sup>-</sup>	16 09Se13	J 1980	$\alpha=?; \beta^+=10\%$			
<sup>183</sup> Pb <sup>m</sup>	-7481	28	94	8	AD	415 ms	20	16 09Se13 J 1980	$\alpha\approx 100; \beta^+ ?; IT ?$		
* <sup>183</sup> Hf <sup>m</sup>	T : for q=71 <sup>+</sup> (H+ like ion); symmetrized from 10(+48-5)										
* <sup>183</sup> Ta <sup>n</sup>	E : less than 50 keV above 1310.16 level										
* <sup>183</sup> Ir	T : average 61Di04=55(7) 61La05=60(6)										
* <sup>183</sup> Hg <sup>m</sup>	I : lack of E(a)=6073- $\gamma$ coinc. in <sup>187</sup> Pb <sup>m</sup> decay; no isomer seen in 01Sc41										
* <sup>183</sup> Tl <sup>m</sup>	E : uncertainty estimated by NUBASE D : IT from 11Ve.A										
<sup>184</sup> Yb	-32540#	500#	1#	s	(>300 ns)	0 <sup>+</sup>	13 12Ku26 I 2012	$\beta^- ?$			
<sup>184</sup> Lu	-36410#	300#	20	s	3	(3 <sup>+</sup> )	10 95Kr04 TJ 1989	$\beta^- =100$			
<sup>184</sup> Hf	-41500	40	4.12	h	0.05	0 <sup>+</sup>	10 1973	$\beta^- =100$			
<sup>184</sup> Hf <sup>m</sup>	-40230	40	1272.2	0.4	48	s	10	12Re.A D 1995	IT=?; $\beta^- =?$		
<sup>184</sup> Hf <sup>n</sup>	-39020	40	2477	10	16	m	7	15 <sup>+</sup> #	10 10Re07 ET 2010	$\beta^- ?; IT ?$	
<sup>184</sup> Ta	-42839	26	8.7	h	0.1	(5 <sup>-</sup> )	10	1955	$\beta^- =100$		
<sup>184</sup> W	-45705.4	0.7	STABLE	(>8.9 Zy)		0 <sup>+</sup>	10	04Co26 T 1930	IS=30.64 2; $\alpha ?$		
<sup>184</sup> W <sup>m</sup>	-44420.4	0.7	1284.997	0.008	8.33	$\mu$ s	0.18	5 <sup>-</sup>	10 1969	IT=100	
<sup>184</sup> W <sup>n</sup>	-41842.2	2.6	3863.2	2.5	188	ns	38	(14 <sup>-</sup> , 15, 17 <sup>-</sup> )	10 2004	IT=100	
<sup>184</sup> Re	-44220	4	35.4	d	0.7	3 <sup>(-)</sup>	10	1940	$\beta^+ =100$		
<sup>184</sup> Re <sup>m</sup>	-44032	4	188.0463	0.0017	169	d	8	8 <sup>(+)</sup>	10 1964	IT=74.5 8; $\epsilon=25.5$ 8	
<sup>184</sup> Os	-44252.5	0.8	STABLE	(>56 Ty)		0 <sup>+</sup>	10	1937	IS=0.02 1; $\alpha ?; 2\beta^+ ?$		
<sup>184</sup> Ir	-39611	28	3.09	h	0.03	5 <sup>-</sup>	10	1960	$\beta^+ =100$		
<sup>184</sup> Ir <sup>m</sup>	-39385	28	225.65	0.11	470	$\mu$ s	30	3 <sup>+</sup>	10 1988	IT=100	
<sup>184</sup> Ir <sup>n</sup>	-39283	28	328.40	0.24	350	ns	90	(7 <sup>+</sup> )	10 1988	IT=100	
<sup>184</sup> Pt	-37334	16	17.3	m	0.2	0 <sup>+</sup>	10	95Bi01 D 1963	$\beta^+ \approx 100; \alpha=0.0017$ 7		
<sup>184</sup> Pt <sup>m</sup>	-35494	16	1840.3	0.8	1.01	ms	0.05	8 <sup>-</sup>	10 1966	IT=100	
<sup>184</sup> Au	-30319	22	20.6	s	0.9	5 <sup>+</sup>	10	1969	$\beta^+ \approx 100; \alpha < 0.016$		
<sup>184</sup> Au <sup>m</sup>	-30251	22	68.46	0.04	47.6	s	1.4	2 <sup>+</sup>	10 1969	$\beta^+ =?; IT=30$ 10; $\alpha < 0.016$	
<sup>184</sup> Hg	-26349	10	30.87	s	0.26	0 <sup>+</sup>	10	1969	$\beta^+ =98.89$ 6; $\alpha=1.11$ 6		
<sup>184</sup> Tl	-16883	10	9.5	s	0.2	(2 <sup>-</sup> )	10	16Va01 TJD 1976	$\beta^+ =98.78$ 30; $\alpha=1.22$ 30		
<sup>184</sup> Tl <sup>m</sup>	-16930	30	-50	30	AD	10#	s	(7 <sup>+</sup> )	16Va01 JD 2016	$\beta^+ ?; IT ?; \alpha=0.047$ 6	
<sup>184</sup> Tl <sup>n</sup>	-16430	30	450	30	AD	47.1	ms	0.7	(10 <sup>-</sup> )	10 15Va10 TD 1984	IT $\approx$ 100; $\alpha=0.089$ 19
<sup>184</sup> Pb	-11052	13	490	ms	25	0 <sup>+</sup>	10	04An07 D 1980	$\alpha=80$ 11; $\beta^+ ?$		
<sup>184</sup> Bi	1060	80	6.6	ms	1.5	3 <sup>+</sup> #	10	2003	$\alpha=?$		
<sup>184</sup> Bi <sup>m</sup>	1210#	130#	150#	100#	*	13	ms	2	10 <sup>-</sup> #	10 2002	$\alpha=?$
* <sup>184</sup> Hf <sup>m</sup>	E : 10Re07=1264(10) T : 12Re19=113(+60-47) for q=72 <sup>+</sup> (bare ion)										
* <sup>184</sup> Hf <sup>n</sup>	T : symmetrized from 12Re19=12(+8-6) for q=72 <sup>+</sup> ; supersedes 10Re07=12(+10-4)										
* <sup>184</sup> Os	T : lower limit is for $\alpha$ decay; 13Be07: $2\beta^+ > 25$ Py										
* <sup>184</sup> Tl <sup>n</sup>	E : 506.1(0.1) keV above <sup>184</sup> Tl <sup>m</sup>										
* <sup>184</sup> Pb	D : average 04An07=80(15)% 03Va16=80(15)%										
<sup>185</sup> Yb	-28500#	500#	300#	ms	(>300 ns)	3/2 <sup>-</sup> #	13 12Ku26 I 2012	$\beta^- ?$			
<sup>185</sup> Lu	-33890#	300#	6#	s	(>300 ns)	7/2 <sup>+</sup> #	13 09St16 I 2009	$\beta^- ?$			
<sup>185</sup> Hf	-38320	60	3.5	m	0.6	3/2 <sup>-</sup> #	06 1993	$\beta^- =100$			
<sup>185</sup> Ta	-41394	14	49.4	m	1.5	7/2 <sup>+</sup> #	06 1950	$\beta^- =100$			
<sup>185</sup> Ta <sup>m</sup>	-40988	14	900	ns	300	(3/2 <sup>+</sup> )	06 07Sh42 ETJ 2007	IT=100			
<sup>185</sup> Ta <sup>n</sup>	-40121	14	1273.4	0.4	11.8	ms	1.4	21/2 <sup>-</sup>	06 09La17 EJT 1999	IT=100	
<sup>185</sup> W	-43387.8	0.7	75.1	d	0.3	3/2 <sup>-</sup>	06 1940	$\beta^- =100$			
<sup>185</sup> W <sup>m</sup>	-43190.4	0.7	197.383	0.023	1.597	m	0.004	11/2 <sup>+</sup>	06 94It.A T 1950	IT=100	
<sup>185</sup> Re	-43819.0	0.8	STABLE			5/2 <sup>+</sup>	06 1931	IS=37.40 2			
<sup>185</sup> Re <sup>m</sup>	-41695.2	1.4	2123.8	1.1	121	ns	13	(21/2)	06 1997	IT=100	
<sup>185</sup> Os	-42805.9	0.8	92.95	d	0.09	1/2 <sup>-</sup>	06 1947	$\epsilon=100$			
<sup>185</sup> Os <sup>m</sup>	-42703.5	0.8	3.0	$\mu$ s	0.4	7/2 <sup>-</sup>	06 FGK128 J 1970	IT=100			
<sup>185</sup> Os <sup>n</sup>	-42530.4	0.8	275.53	0.12	780	ns	50	11/2 <sup>+</sup>	06 1970	IT=100	
<sup>185</sup> Ir	-40336	28	14.4	h	0.1	5/2 <sup>-</sup>	06 1958	$\beta^+ =100$			
<sup>185</sup> Ir <sup>m</sup>	-38140	40	2197	23	120	ms	20	06 1979	IT=100		
<sup>185</sup> Pt	-36688	26	70.9	m	2.4	(9/2 <sup>+</sup> )	06 1960	$\beta^+ \approx 100; \alpha=0.0050$ 20			
<sup>185</sup> Pt <sup>m</sup>	-36585	26	103.41	0.05	33.0	m	0.8	(1/2 <sup>-</sup> )	06 1970	$\beta^+ =?; IT < 2$	
<sup>185</sup> Pt <sup>n</sup>	-36487	26	200.89	0.04	728	ns	20	5/2 <sup>-</sup>	06 1996	IT=100	
<sup>185</sup> Au	-31858.1	2.6	4.25	m	0.06	5/2 <sup>-</sup>	06 1960	$\beta^+ \approx 100; \alpha=0.26$ 6			
<sup>185</sup> Au <sup>m</sup>	-31760#	100#	100#	100#	*	6.8	m	0.3	1/2 <sup>+</sup> #	06 1960	$\beta^+ < 100; IT ?$
<sup>185</sup> Hg	-26184	14	49.1	s	1.0	1/2 <sup>-</sup>	06 1960	$\beta^+ =94$ 1; $\alpha=6$ 1			
<sup>185</sup> Hg <sup>m</sup>	-26080	14	21.6	s	1.5	13/2 <sup>+</sup>	06 13Sa43 E 1970	IT=54 10; $\beta^+ =46$ 10; $\alpha \approx 0.03$			
<sup>185</sup> Tl	-19758	21	19.5	s	0.5	1/2 <sup>(+)</sup> #	06 13Ba41 J 1976	$\beta^+ =?; \alpha ?$			
<sup>185</sup> Tl <sup>m</sup>	-19303	21	454.8	1.5	1.93	s	0.08	9/2 <sup>(-)</sup> #	06 13Ba41 J 1976	IT $\approx$ 100; $\alpha=?; \beta^+ ?$	
<sup>185</sup> Pb	-11541	16	6.3	s	0.4	3/2 <sup>-</sup>	06 1975	$\alpha=34$ 25; $\beta^+ ?$			
<sup>185</sup> Pb <sup>m</sup>	-11470	50	70	50	AD	4.07	s	0.15	13/2 <sup>+</sup>	06 02An15 T 1975	$\alpha=50$ 25; $\beta^+ ?$
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>185</sup> Bi	-2240# 80#				*	&	2# ms	9/2 <sup>-</sup> # 96Da06 J 1996 p ?; α ? *
<sup>185</sup> Bi <sup>m</sup>	-2156 13	80#	80#		*	&	58 μs 4	1/2 <sup>+</sup> 06 1996 p=90 2; α=10 2
<sup>185</sup> Bi <sup>n</sup>	-2060# 100#	180#	60#		EU		50 μs 10	13/2 <sup>+</sup> # 04An07 ITD 2004 p=?; α=? *
* <sup>185</sup> Os <sup>m</sup>	J: E1 from 9/2 <sup>+</sup> **							
* <sup>185</sup> Ir <sup>m</sup>	E: x<80 keV above 2157.3(0.5) level **							
* <sup>185</sup> Pt	D: if the 4444(10) keV α line is from ground-state; otherwise α=0.0010(4)% from isomer **							
* <sup>185</sup> Pb <sup>m</sup>	T: average 02An15=4.3(0.2) 80Sc09=3.73(0.24) (excluding the 6.1 s activity) **							
* <sup>185</sup> Bi	T: estimated from 9/2 <sup>-</sup> isomers in odd Bi and Tl isotopes **							
* <sup>185</sup> Bi <sup>n</sup>	E: 100 keV above <sup>185</sup> Bi <sup>m</sup> T: similar to <sup>185</sup> Bi <sup>m</sup> **							
<sup>186</sup> Lu	-30210# 400#						2# s (>300 ns)	13 12Ku26 I 2012 β <sup>-</sup> ?
<sup>186</sup> Hf	-36420 50						2.6 m 1.2	0 <sup>+</sup> 03 1998 β <sup>-</sup> =100
<sup>186</sup> Hf <sup>m</sup>	-33450 70	2968	43				> 20 s	17 <sup>+</sup> # 10Re07 ET 2010 β <sup>-</sup> ?; IT ? *
<sup>186</sup> Ta	-38610 60						10.5 m 0.3	(2 <sup>-</sup> , 3 <sup>-</sup> ) 03 1955 β <sup>-</sup> =100
<sup>186</sup> Ta <sup>m</sup>	-38270 60	336	20				1.54 m 0.05	9 <sup>+</sup> # 04Xu08 T 2010 β <sup>-</sup> ?; IT ? *
<sup>186</sup> W	-42508.5 1.2						STABLE (>4.1 Ey)	0 <sup>+</sup> 03 03Da09 T 1930 IS=28.43 19; 2β <sup>-</sup> ?; α ? *
<sup>186</sup> W <sup>m</sup>	-40991.3 1.3	1517.2	0.6				18 μs 1	7 <sup>-</sup> 03 12La.A J 1998 IT=100
<sup>186</sup> W <sup>n</sup>	-38965.7 2.4	3542.8	2.1				2.0 s 0.2	16 <sup>+</sup> 03 12La.A TJ 1998 IT=100
<sup>186</sup> Re	-41927.1 0.8						3.7183 d 0.0011	1 <sup>-</sup> 03 1939 β <sup>-</sup> =92.53 10; ε=7.47 10
<sup>186</sup> Re <sup>m</sup>	-41778.9 0.9	148.2	0.5				200 ky	(8 <sup>+</sup> ) 03 15Ma60 E 1972 IT=?; β <sup>-</sup> <10
<sup>186</sup> Os	-42999.9 0.8						2.0 Py 1.1	0 <sup>+</sup> 03 1931 IS=1.59 3; α=100
<sup>186</sup> Ir	-39172 17						16.64 h 0.03	5 <sup>+</sup> 03 1958 β <sup>+</sup> =100
<sup>186</sup> Ir <sup>m</sup>	-39171 17	0.8	0.4				1.92 h 0.05	2 <sup>-</sup> 03 91Be25 ET 1962 β <sup>+</sup> ≈75; IT≈25 *
<sup>186</sup> Pt	-37864 22						2.08 h 0.05	0 <sup>+</sup> 03 1961 β <sup>+</sup> =100; α≈1.4e-4
<sup>186</sup> Au	-31715 21						10.7 m 0.5	3 <sup>-</sup> 03 1960 β <sup>+</sup> =100; α=0.0008 2
<sup>186</sup> Au <sup>m</sup>	-31487 21	227.77	0.07				110 ns 10	2 <sup>+</sup> 03 1983 IT=100
<sup>186</sup> Hg	-28539 12						1.38 m 0.06	0 <sup>+</sup> 03 1960 β <sup>+</sup> ≈100; α=0.016 5
<sup>186</sup> Hg <sup>m</sup>	-26322 12	2217.3	0.4				82 μs 5	(8 <sup>-</sup> ) 03 1984 IT=100
<sup>186</sup> Tl	-19887 22						40# s	(2 <sup>-</sup> ) 03 91Va04 I 1975 β <sup>+</sup> ? *
<sup>186</sup> Tl <sup>m</sup>	-19860 30	20	40				27.5 s 1.0	7 <sup>(+)</sup> 03 13Ba41 J 1975 β <sup>+</sup> ≈100; α≈0.006
<sup>186</sup> Tl <sup>n</sup>	-19490 30	400	40		MD		2.9 s 0.2	10 <sup>(-)</sup> 03 13Ba41 J 1977 IT=100 *
<sup>186</sup> Pb	-14682 11						4.82 s 0.03	0 <sup>+</sup> 03 1972 β <sup>+</sup> ?; α=40 8
<sup>186</sup> Bi	-3146 17						14.8 ms 0.7	(3 <sup>+</sup> ) 03 13La02 D 1997 α≈100; β <sup>+</sup> =0.6#; β <sup>+</sup> SF=0.011 *
<sup>186</sup> Bi <sup>m</sup>	-2980# 100#	170#	100#				9.8 ms 0.4	(10 <sup>-</sup> ) 03 13La02 D 1984 α≈100; β <sup>+</sup> =0.6#; β <sup>+</sup> SF=0.011 *
<sup>186</sup> Po	4101 18						34 μs 12	0 <sup>+</sup> 13 13An13 T 2005 α≈100; p ? *
* <sup>186</sup> Hf <sup>m</sup>	T: for q=72 <sup>+</sup> (bare ion) in 10Re07 **							
* <sup>186</sup> Ta <sup>m</sup>	T: 12Re19=3.0(+1.5-0.8) q=72 <sup>+</sup> (H+ like ion); supersedes 10Re07=3.4(+2.4-1.4) **							
* <sup>186</sup> W	T: given limit is for 2β <sup>-</sup> decay **							
* <sup>186</sup> W	T: for α decay 04Co26>8.2 Zy, 03Da05>170 Ey, 03Ce01>27 Ey, 97Ge15>6.5 Ey **							
* <sup>186</sup> Ir <sup>m</sup>	T: average 91Be25=1.90(0.05) 70Fi.A=2.0(0.1) **							
* <sup>186</sup> Ir <sup>m</sup>	E: E is positive and below 1.5 keV **							
* <sup>186</sup> Tl	I: identified as decay level from <sup>190</sup> Bi in 91Va04 **							
* <sup>186</sup> Tl <sup>n</sup>	E: 374.0(0.2) keV above <sup>186</sup> Tl <sup>m</sup> J: also 12Bi.A **							
* <sup>186</sup> Bi	T: average 03An27=14.8(0.8) 97Ba21=15.0(1.7) **							
* <sup>186</sup> Bi	D: 13La02=0.022 13 for both isomers **							
* <sup>186</sup> Bi <sup>m</sup>	T: from 03An27 **							
* <sup>186</sup> Po	T: symmetrized from 13An13=28(+16-6) **							
<sup>187</sup> Lu	-27580# 400#						1# s (>300 ns)	7/2 <sup>+</sup> # 13 12Ku26 I 2012 β <sup>-</sup> ?
<sup>187</sup> Hf	-32820# 300#						30# s (>300 ns)	3/2 <sup>-</sup> # 09 99Be63 I 1999 β <sup>-</sup> ?
<sup>187</sup> Hf <sup>m</sup>	-32320# 420#	500#	300#		*		270 ns 80	9/2 <sup>-</sup> # 09Al30 TD 2009 IT=100
<sup>187</sup> Ta	-36900 60						2.3 m 6	7/2 <sup>+</sup> # 09 10Re07 T 1999 β <sup>-</sup> ?
<sup>187</sup> Ta <sup>m</sup>	-35110 60	1789	13				22 s 9	27/2 <sup>-</sup> # 10Re07 ET 2010 β <sup>-</sup> ?; IT ? *
<sup>187</sup> Ta <sup>n</sup>	-33970 60	2935	14				> 5 m	41/2 <sup>+</sup> # 10Re07 ET 2010 β <sup>-</sup> ?; IT ? *
<sup>187</sup> W	-39904.0 1.2						24.000 h 0.004	3/2 <sup>-</sup> 09 1940 β <sup>-</sup> =100
<sup>187</sup> W <sup>m</sup>	-39493.9 1.2	410.06	0.04				1.38 μs 0.07	(11/2 <sup>+</sup> ) 09 2008 IT=100
<sup>187</sup> Re	-41216.5 0.7						43.3 Gy 0.07	5/2 <sup>+</sup> 09 1931 IS=62.60 2; β <sup>-</sup> =100; α<0.0001 *
<sup>187</sup> Re <sup>m</sup>	-41010.3 0.7	206.2473	0.0010				555.3 ns 1.7	9/2 <sup>-</sup> 09 1949 IT=100
<sup>187</sup> Re <sup>n</sup>	-39534.5 0.9	1682.0	0.6				354 ns 62	21/2 <sup>+</sup> 09 16Re02 ETJ 2003 IT=100
<sup>187</sup> Os	-41218.9 0.7						STABLE	1/2 <sup>-</sup> 09 1931 IS=1.96 2
<sup>187</sup> Os <sup>m</sup>	-41118.5 0.7	100.45	0.04				112 ns 6	7/2 <sup>-</sup> 09 1964 IT=100
<sup>187</sup> Os <sup>n</sup>	-40961.8 0.7	257.10	0.07				231 μs 2	11/2 <sup>+</sup> 09 1964 IT=100
<sup>187</sup> Ir	-39549 28						10.5 h 0.3	3/2 <sup>+</sup> 09 1958 β <sup>+</sup> =100
<sup>187</sup> Ir <sup>m</sup>	-39363 28	186.16	0.04				30.3 ms 0.6	9/2 <sup>-</sup> 09 1963 IT=100
<sup>187</sup> Ir <sup>n</sup>	-39115 28	433.75	0.06				152 ns 12	11/2 <sup>-</sup> 09 1969 IT=100
<sup>187</sup> Ir <sup>p</sup>	-37061 28	2487.7	0.4				1.8 μs 0.5	29/2 <sup>-</sup> 10Mo09 ETJ 2010 IT=100
<sup>187</sup> Pt	-36685 24						2.35 h 0.03	3/2 <sup>-</sup> 09 1961 β <sup>+</sup> =100
<sup>187</sup> Pt <sup>m</sup>	-36511 24	174.38	0.22				311 μs 15	(11/2 <sup>+</sup> ) 09 1976 IT=100
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>187</sup> Au	-33028	22			8.3 m	0.2		1/2 <sup>(+)</sup>	09 1955	$\beta^+ \approx 100; \alpha = 0.003\#$	
<sup>187</sup> Au <sup>m</sup>	-32908	22	120.33	0.14	2.3 s	0.1		9/2 <sup>(-)</sup>	09 1983	IT=100	
<sup>187</sup> Hg	-28118	14			1.9 m	0.3		3/2 <sup>(-)</sup>	09 70Ha18 TD	$\beta^+ = 100; \alpha > 1.2e-4$	
<sup>187</sup> Hg <sup>m</sup>	-28059	19	59	16	2.4 m	0.3		13/2 <sup>(+)</sup>	09 70Ha18 D	$\beta^+ = 100; \alpha > 2.5e-4$	
<sup>187</sup> Tl	-22445	8			51 s			(1/2 <sup>(+)</sup> )	09 1976	$\beta^+ < 100; \alpha = 0.03\#$	
<sup>187</sup> Tl <sup>m</sup>	-22111	8	334	3	15.60 s	0.12		9/2 <sup>(-)</sup>	09 13Ba41 J	IT=?; $\beta^+ ?; \alpha = 0.15$	
<sup>187</sup> Tl <sup>n</sup>	-20970	50	1480	50	1.11 $\mu$ s				09 2000	IT=100	
<sup>187</sup> Tl <sup>p</sup>	-19863	8	2582.5	0.3	690 ns	40		(25/2 <sup>-</sup> , 27/2 <sup>-</sup> ,...)	09 2000	IT=100	
<sup>187</sup> Pb	-14987	5			15.2 s	0.3		3/2 <sup>-</sup>	09 09Se13 J	$\beta^+ ?; \alpha = 9.5$	
<sup>187</sup> Pb <sup>m</sup>	-14968	11	19	10	18.3 s	0.3		13/2 <sup>(+)</sup>	09 09Se13 J	$\beta^+ ?; \alpha = 12$	
<sup>187</sup> Bi	-6383	10			37 ms	2		9/2 <sup>-</sup> #	09 1999	$\alpha = 100$	
<sup>187</sup> Bi <sup>m</sup>	-6275	12	108	8	370 $\mu$ s	20		1/2 <sup>(+)</sup> #	09 1984	$\alpha = 100$	
<sup>187</sup> Bi <sup>n</sup>	-6131	21	252	18	7 $\mu$ s	5		(13/2 <sup>(+)</sup> )	09 02Hu14 ETJ	IT=100	
<sup>187</sup> Po	2830	30			1.40 ms	0.25		(1/2 <sup>-</sup> , 5/2 <sup>-</sup> )	09 2005	$\alpha \approx 100; \beta^+ ?$	
<sup>187</sup> Po <sup>m</sup>	2830	30	4	27	0.5 ms			13/2 <sup>(+)</sup> #	06An11 ETD	$\alpha = ?; \beta^+ ?$	
* <sup>187</sup> Ta <sup>m</sup>	T : for q=73 <sup>+</sup> (bare ion) in 10Re07										
* <sup>187</sup> Ta <sup>n</sup>	T : for q=73 <sup>+</sup> (bare ion) in 10Re07										
* <sup>187</sup> Re	T : other: 96Bo37=32.9(2.0)y for q=75 <sup>+</sup> (bare ion)										
* <sup>187</sup> Hg	T : from 70Ha18; 98Ru04=2.4 m, not documented, no uncertainty given										
* <sup>187</sup> Hg <sup>m</sup>	T : from 70Ha18; 98Ru04=2.2 m, not documented, no uncertainty given										
* <sup>187</sup> Tl <sup>n</sup>	E : x above 1433.23(0.19) level; x=50(50) keV estimated by NUBASE										
* <sup>187</sup> Bi <sup>n</sup>	T : symmetrized from 3.2(+7.6-2.0)										
<sup>188</sup> Lu	-23790#	500#			300# ms	(>300 ns)			13 12Ku26 I	2012	$\beta^- ?$
<sup>188</sup> Hf	-30880#	300#			20# s	(>300 ns)		0 <sup>+</sup>	02 99Be63 I	1999	$\beta^- ?$
<sup>188</sup> Ta	-33610	50			19.6 s	2.0			02 09A130 TD	1999	$\beta^- = 100$
<sup>188</sup> Ta <sup>m</sup>	-33320	50	292.4	0.2	3.6 $\mu$ s	0.4			05Ca02 ET	2005	IT=100
<sup>188</sup> W	-38668	3			69.78 d	0.05		0 <sup>+</sup>	02 1951	$\beta^- = 100$	
<sup>188</sup> W <sup>m</sup>	-36739	3	1929.3	1.6	109.5 ns	3.5		8 <sup>-</sup>	10La16 ETJ	2010	IT=100
<sup>188</sup> Re	-39016.8	0.7			17.0040 h	0.0022		1 <sup>-</sup>	02 1939	$\beta^- = 100$	
<sup>188</sup> Re <sup>m</sup>	-38844.7	0.7	172.069	0.009	18.59 m	0.04		(6) <sup>-</sup>	02 1953	IT=100	
<sup>188</sup> Os	-41137.2	0.7			STABLE			0 <sup>+</sup>	02 1931	IS=13.24	
<sup>188</sup> Ir	-38345	9			41.5 h	0.5		1 <sup>-</sup>	02 1950	$\beta^- = 100$	
<sup>188</sup> Ir <sup>m</sup>	-37380	30	970	30	4.2 ms	0.2		11 <sup>-</sup> #	02 GAu E	1971	IT $\approx$ 100; $\beta^+ ?$
<sup>188</sup> Pt	-37821	5			10.2 d	0.3		0 <sup>+</sup>	02 1954	$\epsilon = 100; \alpha = 2.6e-5$	
<sup>188</sup> Au	-32371.3	2.7			8.84 m	0.06		1 <sup>(-)</sup>	02 1955	$\beta^- = 100$	
<sup>188</sup> Hg	-30202	12			3.25 m	0.15		0 <sup>+</sup>	02 1960	$\beta^+ = 100; \alpha = 3.7e-5$	
<sup>188</sup> Hg <sup>m</sup>	-27478	12	2724.3	0.4	134 ns	15		(12 <sup>(+)</sup> )	02 1983	IT=100	
<sup>188</sup> Tl	-22336	30			71 s	2		(2 <sup>-</sup> )	02 1970	$\beta^+ = 100$	
<sup>188</sup> Tl <sup>m</sup>	-22308	9	30	30	71 s	1		7 <sup>(+)</sup>	02 13Ba41 J	$\beta^+ = 100$	
<sup>188</sup> Tl <sup>n</sup>	-22030	40	310	30	41 ms	4		(9 <sup>-</sup> )	02 1981	IT $\approx$ 100; $\beta^+ ?$	
<sup>188</sup> Pb	-17815	11			25.1 s	0.1		0 <sup>+</sup>	02 03Va16 D	1972	$\beta^+ = ?; \alpha = 9.3$
<sup>188</sup> Pb <sup>m</sup>	-15237	11	2578.2	0.7	1.15 $\mu$ s	0.03		(8 <sup>-</sup> )	02 04Dr04 T	1999	IT=100
<sup>188</sup> Pb <sup>n</sup>	-15105	11	2709.7	0.3	94 ns	12		(12 <sup>(+)</sup> )	02 04Dr04 EJ	2004	IT=100
<sup>188</sup> Pb <sup>p</sup>	-13032	11	4783.2	0.3	437 ns	55		(19 <sup>-</sup> )	02 04Dr04 ETJ	2000	IT=100
<sup>188</sup> Bi	-7195	11			60 ms	3		3 <sup>+</sup> #	02 13La02 TD	1980	$\alpha = ?; \beta^+ = 1.1\#; \beta^+ SF = 0.0016$
<sup>188</sup> Bi <sup>m</sup>	-7130	30	66	30	> 5 $\mu$ s			7 <sup>+</sup> #	06An04 ET	1984	IT ?
<sup>188</sup> Bi <sup>n</sup>	-7040	30	153	30	265 ms	10		(10 <sup>-</sup> )	02 13La02 TD	1984	$\alpha = ?; \beta^+ = 4.9\#; \beta^+ SF = 0.0016$
<sup>188</sup> Po	-544	20			275 $\mu$ s	30		0 <sup>+</sup>	02 03Va16 T	1999	$\alpha = ?; \beta^+ ?$
* <sup>188</sup> Ta <sup>m</sup>	T : average 11St21=3.5(0.4) 09A130=4.4(1.0); other 05Ca02=5(2)										
* <sup>188</sup> Ir <sup>m</sup>	E : less than 100 keV above 923.5 level, from ENSDF										
* <sup>188</sup> Hg <sup>m</sup>	T : other 04Gl04=270(51)										
* <sup>188</sup> Tl <sup>n</sup>	E : 268.8(0.2) keV above <sup>188</sup> Tl <sup>m</sup> , from 91Va04										
* <sup>188</sup> Pb	D : also 03Va16=8.0(0.6)%										
* <sup>188</sup> Pb <sup>n</sup>	T : lifetime 99Dr10=136(18) ns										
* <sup>188</sup> Pb <sup>p</sup>	T : lifetime $\tau$ =630(80) ns										
* <sup>188</sup> Bi	T : 13La02 supersedes 06An04=66(6) 03An26=60(3)										
* <sup>188</sup> Bi	D : 13La02=0.0032 16 for both beta-delayed fission isomers										
* <sup>188</sup> Bi <sup>n</sup>	T : 13La02 supersedes 06An04=280(20) 03An26=265(15) of same group										
<sup>189</sup> Hf	-27160#	300#			2# s	(>300 ns)		3/2 <sup>-</sup> #	12 09A130 I	2009	$\beta^- ?$
<sup>189</sup> Ta	-31830#	200#			3# s	(>300 ns)		7/2 <sup>(+)</sup> #	03 99Be63 I	1999	$\beta^- ?$
<sup>189</sup> Ta <sup>m</sup>	-30230#	450#	1600#	400#	1.6 $\mu$ s	0.2			09A130 TD	2009	IT=100
<sup>189</sup> W	-35620	40			10.7 m	0.5		3/2 <sup>-</sup> #	03 1963	$\beta^- = 100$	
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...											
<sup>189</sup> Re	-37979	8				24.3 h	0.4	5/2 <sup>+</sup>	03	1963	$\beta^-$ =100
<sup>189</sup> Re <sup>m</sup>	-36208	8	1770.9	0.6		223 $\mu$ s	14	29/2 <sup>+</sup>	03	16Re02 ETJ	IT=100
<sup>189</sup> Os	-38986.7	0.7				STABLE		3/2 <sup>-</sup>	03	1931	IS=16.15 5
<sup>189</sup> Os <sup>m</sup>	-38955.9	0.7	30.812	0.015		5.81 h	0.06	9/2 <sup>-</sup>	03	1960	IT=100
<sup>189</sup> Ir	-38450	13				13.2 d	0.1	3/2 <sup>+</sup>	03	1955	$\epsilon$ =100
<sup>189</sup> Ir <sup>m</sup>	-38078	13	372.17	0.04		13.3 ms	0.3	11/2 <sup>-</sup>	03	1960	IT=100
<sup>189</sup> Ir <sup>n</sup>	-36117	13	2333.2	0.5		3.7 ms	0.2	(25/2) <sup>+</sup>	03	1975	IT=100
<sup>189</sup> Pt	-36469	10				10.87 h	0.12	3/2 <sup>-</sup>	03	1955	$\beta^+$ =100
<sup>189</sup> Pt <sup>m</sup>	-36296	10	172.80	0.06		464 ns	25	9/2 <sup>-</sup>	03	1970	IT=100
<sup>189</sup> Pt <sup>n</sup>	-36278	10	191.5	0.7		143 $\mu$ s	5	(13/2 <sup>+</sup> )	03	1976	IT=100
<sup>189</sup> Au	-33582	20				28.7 m	0.3	1/2 <sup>+</sup>	03	1955	$\beta^+$ =100; $\alpha$ <3e-5
<sup>189</sup> Au <sup>m</sup>	-33335	20	247.23	0.16		4.59 m	0.11	11/2 <sup>-</sup>	03	1966	$\beta^+$ $\approx$ 100; IT=?
<sup>189</sup> Au <sup>n</sup>	-33257	20	325.11	0.16		190 ns	15	9/2 <sup>-</sup>	03	1975	IT=100
<sup>189</sup> Au <sup>p</sup>	-31027	20	2554.7	1.2		242 ns	10	31/2 <sup>+</sup>	03	1975	IT=100
<sup>189</sup> Hg	-29630	30				7.6 m	0.1	3/2 <sup>-</sup>	03	1955	$\beta^+$ =100; $\alpha$ <3e-5
<sup>189</sup> Hg <sup>m</sup>	-29548	18	80	30	MD	8.6 m	0.1	13/2 <sup>+</sup>	03	1966	$\beta^+$ =100; $\alpha$ <3e-5
<sup>189</sup> Tl	-24616	8				2.3 m	0.2	(1/2 <sup>+</sup> )	11	1972	$\beta^+$ =100
<sup>189</sup> Tl <sup>m</sup>	-24331	8	285	6	AD	1.4 m	0.1	9/2 <sup>(-)</sup>	11	85Bo46 J	$\beta^+$ $\approx$ 100; IT<4
<sup>189</sup> Pb	-17844	14				39 s	8	3/2 <sup>-</sup>	11	09Sa09 T	$\beta^+$ $\approx$ 100; $\alpha$ $\approx$ 0.4
<sup>189</sup> Pb <sup>m</sup>	-17804	14	40	4	AD *	50.5 s	2.1	13/2 <sup>+</sup>	11	09Sa09 T	2009 $\beta^+$ $\approx$ 100; $\alpha$ <1; IT ?
<sup>189</sup> Pb <sup>n</sup>	-15370#	30#	2475#	30#		26 $\mu$ s	5	(31/2 <sup>-</sup> )	11	2005	IT=100
<sup>189</sup> Bi	-10065	21				658 ms	47	(9/2 <sup>-</sup> )	11	1973	$\alpha$ $\approx$ 100
<sup>189</sup> Bi <sup>m</sup>	-9881	21	184	5	AD	4.9 ms	0.3	(1/2 <sup>+</sup> )	11	03An26 T	1984 $\alpha$ >50; $\beta^+$ <50
<sup>189</sup> Bi <sup>n</sup>	-9707	21	357.6	0.5		880 ns	50	(13/2 <sup>+</sup> )	11	2001	IT $\approx$ 100
<sup>189</sup> Po	-1422	22				3.8 ms	0.4	(5/2 <sup>-</sup> )	07	05Va04 T	1999 $\alpha$ $\approx$ 100; $\beta^+$ ?
* <sup>189</sup> Ta <sup>m</sup>	T : other 11St21=0.58(0.22), possibly a different isomer										**
* <sup>189</sup> Tl <sup>m</sup>	J : also 13Ba41=9/2										**
* <sup>189</sup> Pb	J : 09Se13: $\alpha$ to <sup>185</sup> Hg 26.1 level										**
* <sup>189</sup> Pb <sup>m</sup>	T : average 09Sa09=50(3) 72Ga27=51(3)										**
* <sup>189</sup> Pb <sup>m</sup>	J : 09Se13: from $\alpha$ decay from <sup>193</sup> Po <sup>m</sup>										**
* <sup>189</sup> Pb <sup>n</sup>	E : 2434.50(0.18) keV above <sup>189</sup> Pb <sup>m</sup> (13/2 <sup>+</sup> )										**
* <sup>189</sup> Pb <sup>n</sup>	T : from lifetime 05Ba51 $\tau$ =32(+10-2) $\mu$ s, or T=22.2(+6.9-1.4)										**
* <sup>189</sup> Bi <sup>m</sup>	T : average 03An26=4.9(0.5) 03Ke08=4.6(+0.8-0.6) 97An09=4.8(0.5)										**
* <sup>189</sup> Bi <sup>m</sup>	T : and 97Wa05=5.2(0.6); 95Ba75=7.0(0.2), conflicting not used										**
* <sup>189</sup> Po	T : average 05Va04=3.5(0.5) 99An52=5(1) J : favored decay to (5/2 <sup>-</sup> ) level										**
<sup>190</sup> Hf	-25030#	400#				2# s	(>300 ns)	0 <sup>+</sup>	13	12Ku26 I	2012 $\beta^-$ ?
<sup>190</sup> Ta	-28510#	200#			*	5.3 s	0.7	(3)	10	09Al30 TJD	2009 $\beta^-$ =100
<sup>190</sup> Ta <sup>m</sup>	-28310#	250#	200#	150#	*	42 ns	7		10	09Al30 TD	2009 IT=100
<sup>190</sup> W	-34380	40				30.0 m	1.5	0 <sup>+</sup>	03	1976	$\beta^-$ =100
<sup>190</sup> W <sup>m</sup>	-32640	40	1742.0	2.0		111 ns	17	8 <sup>+</sup>	03	10La16 ETJ	2010 IT=100
<sup>190</sup> W <sup>n</sup>	-32540	40	1839.0	2.2		166 $\mu$ s	6	10 <sup>-</sup>	03	10La16 ETJ	2000 IT=100
<sup>190</sup> Re	-35640	70				3.1 m	0.3	(2) <sup>-</sup>	03	1955	$\beta^-$ =100
<sup>190</sup> Re <sup>m</sup>	-35440	70	204	10		3.2 h	0.2	(6 <sup>-</sup> )	03	12Re19 E	1962 $\beta^-$ =54.4 20; IT ?
<sup>190</sup> Os	-38707.8	0.6				STABLE		0 <sup>+</sup>	03	1931	IS=26.26 2
<sup>190</sup> Os <sup>m</sup>	-37002.4	0.6	1705.4	0.2		9.86 m	0.03	10 <sup>-</sup>	03	12Kr05 T	1950 IT=100
<sup>190</sup> Ir	-36753.5	1.4				11.78 d	0.10	4 <sup>-</sup>	03	1947	$\beta^+$ =100; $e^+$ <0.002
<sup>190</sup> Ir <sup>m</sup>	-36727.4	1.4	26.1	0.1		1.120 h	0.003	(1) <sup>-</sup>	03	1964	IT=100
<sup>190</sup> Ir <sup>n</sup>	-36717.3	1.4	36.154	0.025		> 2 $\mu$ s		(4) <sup>+</sup>	03	1996	IT=100
<sup>190</sup> Ir <sup>p</sup>	-36377.1	1.4	376.4	0.1		3.087 h	0.012	(11) <sup>-</sup>	03	1950	$\beta^+$ =91.4 2; IT=8.6 2
<sup>190</sup> Pt	-37306.5	0.7				650 Gy	30	0 <sup>+</sup>	03	1949	IS=0.012 2; $\alpha$ =100; 2 $\beta^+$ ?
<sup>190</sup> Au	-32834	3				42.8 m	1.0	1 <sup>-</sup>	03	1959	$\beta^+$ =100; $\alpha$ <1e-6
<sup>190</sup> Au <sup>m</sup>	-32630#	150#	200#	150#	*	125 ms	20	11 <sup>-</sup> #	03	1982	IT $\approx$ 100; $\beta^+$ ?
<sup>190</sup> Hg	-31371	16				20.0 m	0.5	0 <sup>+</sup>	03	1959	$\epsilon$ $\approx$ 100; $e^+$ <1; $\alpha$ <3.4e-7
<sup>190</sup> Tl	-24372	8				2.6 m	0.3	2 <sup>(-)</sup>	03	1970	$\beta^+$ =100
<sup>190</sup> Tl <sup>m</sup>	-24289	6	83	10	MD *	3.7 m	0.3	7 <sup>(+#)</sup>	03	1970	$\beta^+$ =100
<sup>190</sup> Tl <sup>n</sup>	-24080#	70#	290#	70#		750 $\mu$ s	40	(8 <sup>-</sup> )	03	1981	IT=100
<sup>190</sup> Tl <sup>p</sup>	-23960#	70#	410#	70#		> 1 $\mu$ s		9 <sup>-</sup>	03	91Va04 ET	1991 IT ?
<sup>190</sup> Pb	-20417	13				71 s	1	0 <sup>+</sup>	03	1972	$\beta^+$ ?; $\alpha$ =0.40 4
<sup>190</sup> Pb <sup>m</sup>	-17802	13	2614.8	0.8		150 ns	14	10 <sup>+</sup>	03	01Dr05 J	1998 IT=100
<sup>190</sup> Pb <sup>n</sup>	-17799	24	2618	20		24.3 $\mu$ s	2.1	(12 <sup>+</sup> )	03	1998	IT ?
<sup>190</sup> Pb <sup>p</sup>	-17759	13	2658.2	0.8		7.7 $\mu$ s	0.3	11 <sup>-</sup>	03	01Dr05 JT	1985 IT=100
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...										
<sup>190</sup> Bi	-10600	23			6.3 s	0.1	(3 <sup>+</sup> )	03 91Va04 J	1972	$\alpha=77$ 21; $\beta^+=?$ *
<sup>190</sup> Bi <sup>m</sup>	-10470	30	130	40	AD		(10 <sup>-</sup> )	03 91Va04 J	1988	$\alpha=70$ 9; $\beta^+ ?$ ; $\beta^+p ?$ *
<sup>190</sup> Bi <sup>n</sup>	-10479	27	121	15			(5 <sup>-</sup> )	09An11 ET	2009	IT=100 *
<sup>190</sup> Bi <sup>p</sup>	-10200	50	404	40			(8 <sup>-</sup> )	03 09An11 EJT	2001	IT=100 *
<sup>190</sup> Po	-4564	13			2.46 ms	0.05	0 <sup>+</sup>	03	1996	$\alpha \approx 100$ ; $\beta^+ = 0.1\%$ *
* <sup>190</sup> W <sup>n</sup>	T : others 11St21=108(9) 09Al30=106(18) $\mu$ s 05Ca02=60(+1500-30) $\mu$ s 00Po26<3.1ms **									
* <sup>190</sup> W <sup>m</sup>	E : other 00Po26=2381 **									
* <sup>190</sup> Os <sup>m</sup>	J : M2 + E3 to 8 <sup>+</sup> member of the ground-state band **									
* <sup>190</sup> Tl <sup>n</sup>	E : 161.9 keV above <sup>190</sup> Tl <sup>m</sup> **									
* <sup>190</sup> Tl <sup>p</sup>	E : 236.2 keV above <sup>190</sup> Tl <sup>m</sup> **									
* <sup>190</sup> Pb <sup>m</sup>	T : uncertainty from 12Dr.A **									
* <sup>190</sup> Pb <sup>n</sup>	E : above <sup>190</sup> Pb <sup>m</sup> , see 01Dr05 T : uncertainty from 12Dr.A **									
* <sup>190</sup> Pb <sup>p</sup>	T : average 01Dr05=7.2(0.6) 85St16=7.9(0.4) **									
* <sup>190</sup> Bi	D : symmetrized from $\alpha=90(+10-30)\%$ T : also 13Ny01=7.7(+1.0-0.8) not used **									
* <sup>190</sup> Bi <sup>m</sup>	T : also 13Ny01=5.9(+1.0-0.8) not used **									
* <sup>190</sup> Bi <sup>n</sup>	J : E1 and M1(+E2) $\gamma$ s in cascade to (3 <sup>+</sup> ), absence of direct $\gamma$ to (3 <sup>+</sup> ) **									
* <sup>190</sup> Bi <sup>p</sup>	E : 274(1) keV above the (10 <sup>-</sup> ) isomer J : E2 to (10 <sup>-</sup> ) **									
* <sup>190</sup> Bi <sup>p</sup>	T : symmetrized from 09An11=1.0(+1.0-0.5) **									
<sup>191</sup> Ta	-26490#	300#			3# s	(>300 ns)	7/2 <sup>+</sup> #	11 09St16 I	2009	$\beta^- ?$
<sup>191</sup> W	-31180	40			* 45# s	(>300 ns)	3/2 <sup>-</sup> #	07 99Be63 I	1999	$\beta^- ?$
<sup>191</sup> W <sup>m</sup>	-30950	60	235	50	* 340 ns	14		11St21 ETD	2009	IT=100 *
<sup>191</sup> Re	-34350	10			9.8 m	0.5	(3/2 <sup>+</sup> , 1/2 <sup>+</sup> )	07	1963	$\beta^- = 100$ *
<sup>191</sup> Re <sup>m</sup>	-32749	10	1601.5	0.4	51 $\mu$ s	3	25/2 <sup>-</sup>	16Re02 EJT	2011	IT=100 *
<sup>191</sup> Os	-36395.2	0.7			14.99 d	0.02	9/2 <sup>-</sup>	07 12Kr05 T	1940	$\beta^- = 100$
<sup>191</sup> Os <sup>m</sup>	-36320.8	0.7	74.382	0.003	13.10 h	0.05	3/2 <sup>-</sup>	07 12Kr05 T	1952	IT=100 *
<sup>191</sup> Ir	-36708.8	1.3			STABLE		3/2 <sup>+</sup>	07	1935	IS=37.3 2
<sup>191</sup> Ir <sup>m</sup>	-36537.5	1.3	171.29	0.04	4.899 s	0.023	(11/2 <sup>-</sup> )	07	1955	IT=100
<sup>191</sup> Ir <sup>n</sup>	-34607.8	1.6	2101.0	0.9	5.7 s	0.4	31/2 <sup>(+)</sup>	07 12Dr02 ETJ	1979	IT=100 *
<sup>191</sup> Pt	-35698	4			2.83 d	0.02	3/2 <sup>-</sup>	07	1948	$\epsilon = 100$
<sup>191</sup> Pt <sup>m</sup>	-35597	4	100.663	0.020	> 1 $\mu$ s		(9/2 <sup>-</sup> )	07	1976	IT=100
<sup>191</sup> Pt <sup>n</sup>	-35549	4	149.035	0.022	95 $\mu$ s	5	(13/2 <sup>+</sup> )	07	1967	IT=100
<sup>191</sup> Au	-33798	5			3.18 h	0.08	3/2 <sup>+</sup>	07	1954	$\beta^+ = 100$
<sup>191</sup> Au <sup>m</sup>	-33532	5	266.2	0.7	920 ms	110	(11/2 <sup>-</sup> )	07	1971	IT=100
<sup>191</sup> Au <sup>n</sup>	-31308	5	2489.6	0.9	402 ns	20	(31/2 <sup>+</sup> )	07	1985	IT=100
<sup>191</sup> Hg	-30592	22			49 m	10	3/2 <sup>(-)</sup>	07 86UI02 J	1954	$\beta^+ = 100$ ; $\alpha < 5e-6$
<sup>191</sup> Hg <sup>m</sup>	-30460	30	128	22	50.8 m	1.5	13/2 <sup>(+)</sup>	07 01Sc41 E	1954	$\beta^+ = 100$ ; $\alpha < 5e-6$ *
<sup>191</sup> Tl	-26283	7			20# m		1/2 <sup>(+)</sup>	07 13Ba41 J	1974	$\beta^+ ?$
<sup>191</sup> Tl <sup>m</sup>	-25986	7	297	7	BD 5.22 m	0.16	9/2 <sup>(-)</sup>	07	1970	$\beta^+ = 100$
<sup>191</sup> Pb	-20230	40			* 1.33 m	0.08	(3/2 <sup>-</sup> )	07 10Co13 JD	1974	$\beta^+ \approx 100$ ; $\alpha = 0.51$ 5
<sup>191</sup> Pb <sup>m</sup>	-20231	28	0	50	MD * 2.18 m	0.08	13/2 <sup>(+)</sup>	07 88Me.A J	1975	$\beta^+ \approx 100$ ; $\alpha \approx 0.02$
<sup>191</sup> Pb <sup>n</sup>	-17610	60	2620	50	180 ns	80	(33/2 <sup>+</sup> )	07	1999	IT=100 *
<sup>191</sup> Bi	-13239	7			11.7 s	0.4	(9/2 <sup>-</sup> )	16 13Ny01 T	1972	$\alpha = 51$ 10; $\beta^+ ?$
<sup>191</sup> Bi <sup>m</sup>	-12997	9	242	4	AD 114 ms	6	(1/2 <sup>+</sup> )	16 13Ny01 T	1981	$\alpha = 68$ 5; IT=32 5; B ?
<sup>191</sup> Bi <sup>n</sup>	-12809	7	429.7	0.5	562 ns	10	13/2 <sup>+</sup> #	16	2001	IT=100
<sup>191</sup> Bi <sup>p</sup>	-11364	26	1875	25	400 ns	40		16	2016	IT=100 *
<sup>191</sup> Po	-5069	7			22 ms	1	(3/2 <sup>-</sup> )	07	1993	$\alpha = ?$ ; $\beta^+ = 1\%$
<sup>191</sup> Po <sup>m</sup>	-5008	12	61	11	AD 93 ms	3	(13/2 <sup>+</sup> )	07	1999	$\alpha = ?$ ; $\beta^+ = 4\%$
<sup>191</sup> At	3864	16			2.1 ms	0.8	(1/2 <sup>+</sup> )	07	2003	$\alpha \approx 100$ ; $\beta^+ ?$
<sup>191</sup> At <sup>m</sup>	3922	18	58	20	AD 2.2 ms	0.4	(7/2 <sup>-</sup> )	07	2003	$\alpha \approx 100$ ; $\beta^+ ?$
* <sup>191</sup> W <sup>m</sup>	T : average 11St21=360(20) 09Al30=320(20) ns E : 68 + 167 keV $\gamma$ -rays **									
* <sup>191</sup> Re	I : also an isomer with T=77(33) $\mu$ s decaying by g of 444, 419, 225, 139 keV **									
* <sup>191</sup> Os <sup>m</sup>	T : other 12Kr05=13.6(0.2) from the decay growth J : M3 + E4 to 9/2 <sup>-</sup> **									
* <sup>191</sup> Ir <sup>n</sup>	T : average 12Dr02=5.8(0.6) 79Lu01=5.5(0.7) **									
* <sup>191</sup> Ir <sup>n</sup>	E : from least-squares fit to $\gamma$ -ray energies using 12Dr02 level scheme **									
* <sup>191</sup> Hg <sup>m</sup>	E : original uncertainty (8 keV) increased by 20 for gs+m lines in trap **									
* <sup>191</sup> Pb <sup>m</sup>	E : 2602.31(0.24) above <sup>191</sup> Pb <sup>m</sup> T : symmetrized from 150(+100-50) **									
* <sup>191</sup> Bi <sup>p</sup>	E : x keV above 1825.1 level; x=50# **									
* <sup>191</sup> At	T : symmetrized from 1.7(+1.1-0.5) **									
* <sup>191</sup> At <sup>m</sup>	T : symmetrized from 2.1(+0.4-0.3) **									
<sup>192</sup> Ta	-23060#	400#			2.2 s	0.7	(2)	12	2009	$\beta^- = 100$
<sup>192</sup> W	-29650#	200#			1# m	(>300 ns)	0 <sup>+</sup>	12	1999	$\beta^- ?$
<sup>192</sup> Re	-31590	70			16.0 s	0.9		12 12Al05 T	1965	$\beta^- = 100$
<sup>192</sup> Re <sup>m</sup>	-31430	70	159	1	88 $\mu$ s	8		12 11St21 ETD	2005	IT=100 *
<sup>192</sup> Re <sup>n</sup>	-31320	70	267	10	70 s	30		12 12Re19 ET	2012	$\beta^- = ?$ ; IT=? *
... A-group is continued on next page ...										



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>192</sup> Os	-35882.2	2.3	STABLE	(>53 Ey)	0 <sup>+</sup>	12 13Be07 T	1931	IS=40.78 19; 2β <sup>-</sup> ?; α ?	*		
<sup>192</sup> Os <sup>m</sup>	-33866.8	2.3	2015.40	0.11	5.9 s	0.1	10 <sup>-</sup>	12 13Dr05 J	1965	IT>87; β <sup>-</sup> <13	*
<sup>192</sup> Os <sup>n</sup>	-31301.9	2.5	4580.3	1.0	204 ns	7	(20 <sup>+</sup> )	12 13Dr05 ETJ	2004	IT=100	
<sup>192</sup> Ir	-34835.6	1.3			73.830 d	0.015	4 <sup>+</sup>	12 14Un01 T	1937	β <sup>-</sup> =95.24 4; ε=4.76 4	
<sup>192</sup> Ir <sup>m</sup>	-34778.9	1.3	56.720	0.005	1.45 m	0.05	1 <sup>-</sup>	12	1937	IT≈100; β <sup>-</sup> =0.0175	
<sup>192</sup> Ir <sup>n</sup>	-34667.5	1.3	168.14	0.12	241 y	9	(11 <sup>-</sup> )	12	1959	IT=100	
<sup>192</sup> Pt	-36288.5	2.6			STABLE		0 <sup>+</sup>	12	1935	IS=0.782 24	
<sup>192</sup> Pt <sup>m</sup>	-34116.1	2.6	2172.37	0.13	272 ns	23	(10 <sup>-</sup> )	12	1976	IT=100	
<sup>192</sup> Au	-32772	16			4.94 h	0.09	1 <sup>-</sup>	12	1948	β <sup>+</sup> =100	
<sup>192</sup> Au <sup>m</sup>	-32637	16	135.41	0.25	29 ms		5# <sup>+</sup>	12	1976	IT=100	
<sup>192</sup> Au <sup>n</sup>	-32340	16	431.6	0.5	160 ms	20	(11 <sup>-</sup> )	12	1976	IT=100	
<sup>192</sup> Hg	-32012	16			4.85 h	0.20	0 <sup>+</sup>	12	1952	ε=100; α<4e-6	
<sup>192</sup> Tl	-25870	30			9.6 m	0.4	2 <sup>(-)</sup>	12 13Ba41 J	1961	β <sup>+</sup> =100	
<sup>192</sup> Tl <sup>m</sup>	-25730	50	138	45	10.8 m	0.2	7 <sup>(+)</sup>	12 13Ba41 J	1961	β <sup>+</sup> =100	
<sup>192</sup> Tl <sup>n</sup>	-25480	50	388	45	296 ns	5	(8 <sup>-</sup> )	12	1980	IT=100	
<sup>192</sup> Tl <sup>p</sup>	-25695	25	180	40			(3 <sup>+</sup> )	12 91Va04 E	1991	α=100	
<sup>192</sup> Pb	-22556	13			3.5 m	0.1	0 <sup>+</sup>	12	1974	β <sup>+</sup> ≈100; α=0.0059 7	
<sup>192</sup> Pb <sup>m</sup>	-19975	13	2581.1	0.4	166 ns	6	10 <sup>+</sup>	12 07Io03 J	1985	IT=100	
<sup>192</sup> Pb <sup>n</sup>	-19931	13	2625.1	1.1	1.09 μs	0.04	12 <sup>+</sup>	12 07Io03 J	1979	IT=100	
<sup>192</sup> Pb <sup>p</sup>	-19813	13	2743.5	0.4	756 ns	14	11 <sup>-</sup>	12 07Io03 J	1991	IT=100	
<sup>192</sup> Bi	-13530	30			34.6 s	0.9	(3 <sup>+</sup> )	12	1971	β <sup>+</sup> =88 5; α=12 5	
<sup>192</sup> Bi <sup>m</sup>	-13398	9	140	30	39.6 s	0.4	(10 <sup>-</sup> )	12	1966	β <sup>+</sup> =90 3; α=10 3	
<sup>192</sup> Po	-8071	11			32.2 ms	0.3	0 <sup>+</sup>	12	1977	α=?; β <sup>+</sup> =0.5#	
<sup>192</sup> Po <sup>m</sup>	-5776	11	2294.6	1.0	580 ns	100	(11 <sup>-</sup> )	12	1999	IT=100	
<sup>192</sup> At	2926	28			11.5 ms	0.6	3 <sup>+</sup> #	12 13An03 D	2006	α=100; β <sup>+</sup> =0.6#; β <sup>+</sup> SF=0.21	*
<sup>192</sup> At <sup>m</sup>	2926	28	0	40	88 ms	6	(9 <sup>-</sup> , 10 <sup>-</sup> )	12 13An03 D	2006	α=100; β <sup>+</sup> =4.6#; β <sup>+</sup> SF=0.21	*
* <sup>192</sup> Re	T : average 12Al05=16(2) 79Ka.B=16(1)									**	
* <sup>192</sup> Re <sup>m</sup>	T : average 11St21=85(10) 09Al30=93(15); also 05Ca02=120(+210-50) μs									**	
* <sup>192</sup> Re <sup>n</sup>	E : 159.3 keV γ and X rays seen only in 11St21									**	
* <sup>192</sup> Re <sup>n</sup>	T : symmetrized from 12Re19=61(+40-20) s for q=75 <sup>+</sup>									**	
* <sup>192</sup> Os	T : lower limit is for ββ decay									**	
* <sup>192</sup> Os <sup>m</sup>	T : 15Ak02 H-like τ=15.1(+1.5-1.3) s, T=10.5(+1.0-0.9) s									**	
* <sup>192</sup> Ir	T : average 14Un01=73.831(0.074) 92Wo06=73.84(0.05) and 73.814(0.017)									**	
* <sup>192</sup> Ir	T : 80Ho17=73.831(0.074) 72La14=74.02(0.06)									**	
* <sup>192</sup> Ir	T : original unc of 80Ho17=0.008 increased to 0.1% by evaluator									**	
* <sup>192</sup> At	D : 13An03=0.42 9 for both isomers									**	
<sup>193</sup> Ta	-20870#	400#			500# ms	(>300 ns)	7/2 <sup>+</sup> #	13 12Ku26 I	2012	β <sup>-</sup> ?; β <sup>-</sup> n=0.7#	
<sup>193</sup> W	-26290#	200#			3# s	(>300 ns)	3/2 <sup>-</sup> #	11 09St16 I	2009	β <sup>-</sup> ?	
<sup>193</sup> Re	-30230	40			20# s	(>300 ns)	5/2 <sup>+</sup> #	06 99Be63 I	1999	β <sup>-</sup> ?	
<sup>193</sup> Re <sup>m</sup>	-30080	40	146.0	0.2	69 μs	6	(9/2 <sup>-</sup> )	06 11St21 ETJ	2005	IT=100	*
<sup>193</sup> Os	-33394.3	2.3			29.830 h	0.018	3/2 <sup>-</sup>	06 12Kr05 T	1940	β <sup>-</sup> =100	*
<sup>193</sup> Os <sup>m</sup>	-33152.3	2.4	242.0	0.5	132 ns	29		11St21 ETD	2011	IT=100	
<sup>193</sup> Ir	-34536.2	1.3			STABLE		3/2 <sup>+</sup>	06	1935	IS=62.7 2	
<sup>193</sup> Ir <sup>m</sup>	-34456.0	1.3	80.239	0.006	10.53 d	0.04	11/2 <sup>-</sup>	06	1957	IT=100	
<sup>193</sup> Ir <sup>n</sup>	-32258.7	1.6	2277.5	1.0	124.8 μs	2.1	31/2 <sup>+</sup>	12Dr02 ETJ	2012	IT=100	
<sup>193</sup> Pt	-34479.6	1.4			50 y	6	1/2 <sup>-</sup>	06	1948	ε=100	
<sup>193</sup> Pt <sup>m</sup>	-34329.8	1.4	149.78	0.04	4.33 d	0.03	13/2 <sup>+</sup>	06	1949	IT=100	
<sup>193</sup> Au	-33405	9			17.65 h	0.15	3/2 <sup>+</sup>	06	1948	β <sup>+</sup> =100; α<1e-5	
<sup>193</sup> Au <sup>m</sup>	-33115	9	290.19	0.03	3.9 s	0.3	11/2 <sup>-</sup>	06	1955	IT≈100; β <sup>+</sup> ≈0.03	
<sup>193</sup> Au <sup>n</sup>	-30919	9	2486.5	0.6	150 ns	50	31/2 <sup>+</sup>	06 07Ok05 J	1985	IT=100	
<sup>193</sup> Hg	-31062	16			3.80 h	0.15	3/2 <sup>(-)</sup>	06	1952	β <sup>+</sup> =100	
<sup>193</sup> Hg <sup>m</sup>	-30921	16	140.76	0.05	11.8 h	0.2	13/2 <sup>(+)</sup>	06	1973	β <sup>+</sup> =92.8 5; IT=7.2 5	
<sup>193</sup> Tl	-27477	7			21.6 m	0.8	1/2 <sup>(+)</sup> #	06	1960	β <sup>+</sup> =100	
<sup>193</sup> Tl <sup>m</sup>	-27105	8	372	4	2.11 m	0.15	9/2 <sup>(-)</sup>	06 13Ba41 J	1963	IT=75; β <sup>+</sup> =25	*
<sup>193</sup> Pb	-22190	50			5# m		(3/2 <sup>-</sup> )	06 GAu J	1974	β <sup>+</sup> ?	*
<sup>193</sup> Pb <sup>m</sup>	-22060#	90#	130#	80#	5.8 m	0.2	13/2 <sup>+</sup>	06 91Du07 J	1974	β <sup>+</sup> =100	
<sup>193</sup> Pb <sup>n</sup>	-19450#	90#	2742#	80#	180 ns	15	33/2 <sup>+</sup>	06 04Io01 J	1991	IT=100	*
<sup>193</sup> Bi	-15885	8			63.6 s	3.0	(9/2 <sup>-</sup> )	06	1971	β <sup>+</sup> ?; α=3.5 15	*
<sup>193</sup> Bi <sup>m</sup>	-15580	9	305	6	3.07 s	0.13	(1/2 <sup>+</sup> )	06 15He27 T	1970	α=84 16; β <sup>+</sup> ?	
<sup>193</sup> Bi <sup>n</sup>	-15280	8	605.5	0.5	153 ns	10	(13/2 <sup>+</sup> )	06	2004	IT=100	
<sup>193</sup> Bi <sup>p</sup>	-13535	9	2350	5	85 μs	3	(29/2 <sup>+</sup> )	06 15He27 EJT	2004	IT=100	
<sup>193</sup> Bi <sup>q</sup>	-13480	9	2405	5	3.02 μs	0.08	(29/2 <sup>-</sup> )	06 15He27 ET	2004	IT=100	
<sup>193</sup> Po	-8325	15			388 ms	40	(3/2 <sup>-</sup> )	15 13Se03 J	1967	α=?; β <sup>+</sup> =5#	
<sup>193</sup> Po <sup>m</sup>	-8225	15	100	6	245 ms	11	(13/2 <sup>+</sup> )	15 13Se03 J	1981	α=?; β <sup>+</sup> =3#	
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>193</sup> At	-67 22		29 ms 5	1/2 <sup>+</sup> #	06		2003	$\alpha \approx 100$
<sup>193</sup> At <sup>m</sup>	-59 21	8	21 ms 5	7/2 <sup>-</sup> #	06		1995	$\alpha \approx 100$
<sup>193</sup> At <sup>n</sup>	-25 21	42	28 ms 4	13/2 <sup>+</sup> #	06		2003	$\alpha \approx 24$ 10; IT=76 10
<sup>193</sup> Rn	9043 25		1.15 ms 0.27	3/2 <sup>-</sup> #	07	06An36	TD 2006	$\alpha \approx 100$
* <sup>193</sup> Re <sup>m</sup>	E : a $\gamma$ of 11St21=145.2(0.5) 09Al30=146.1(0.2) keV is observed in decay							
* <sup>193</sup> Re <sup>m</sup>	T : average 11St21=65(9) 09Al30=72(8); also 05Ca02=75(+450-40) $\mu$ s							
* <sup>193</sup> Os	T : other 92An13=30.11(0.01); large syst. unc due to large dead-time effect							
* <sup>193</sup> Os	I : also an isomer with T=132(29) decaying via a 242 keV $\gamma$ -ray							
* <sup>193</sup> Tl <sup>m</sup>	E : less than 13 keV above 365.2 level, from ENSDF							
* <sup>193</sup> Pb	J : from $\alpha$ decay from <sup>197</sup> Po							
* <sup>193</sup> Pb	T : T=4.0m reported in Karlsruhe charts 1981 and 1995. Not traceable							
* <sup>193</sup> Pb <sup>n</sup>	E : 2612.5(0.5) above <sup>193</sup> Pb <sup>m</sup>							
* <sup>193</sup> Bi	D : $\alpha=3.5$ 15 is from ENSDF'98, wrongly attributed in ENSDF'2006 to NUBASE							
* <sup>193</sup> At	T : symmetrized from 28(+5-4)							
* <sup>193</sup> At <sup>n</sup>	T : symmetrized from 27(+4-3)							
<sup>194</sup> Ta	-17300# 500#		300# ms (>300 ns)		13	12Ku26	I 2012	$\beta^- ?$ ; $\beta^- n=0.02\%$
<sup>194</sup> W	-24530# 300#		5# s (>300 ns)	0 <sup>+</sup>	11		2008	$\beta^- ?$
<sup>194</sup> Re	-27240# 200#		5 s 1	(0 <sup>+</sup> , 1)	14	12Al105	T 1999	$\beta^- =100$
<sup>194</sup> Re <sup>m</sup>	-26960# 200#	285 40	25 s 8	(11 <sup>-</sup> )	14	12Re19	E 2012	$\beta^- =100$
<sup>194</sup> Re <sup>n</sup>	-26410# 200#	833 33	100 s 10		14	12Re19	E 2012	$\beta^- =100$
<sup>194</sup> Re <sup>p</sup>	-26140# 200#	1100# 1000#	45 $\mu$ s 18		14	11St21	TD 2011	IT=100
<sup>194</sup> Re <sup>q</sup>	-25240# 200#	2000# 1000#	38 $\mu$ s 37		14		2005	IT=100
<sup>194</sup> Os	-32435.1 2.4		6.0 y 0.2	0 <sup>+</sup>	06		1951	$\beta^- =100$
<sup>194</sup> Ir	-32531.7 1.3		19.28 h 0.13	1 <sup>-</sup>	06		1937	$\beta^- =100$
<sup>194</sup> Ir <sup>m</sup>	-32384.6 1.3	147.072 0.002	31.85 ms 0.24	4 <sup>+</sup>	06		1959	IT=100
<sup>194</sup> Ir <sup>n</sup>	-32160 70	370 70	171 d 11	(10, 11) <sup>(-#)</sup>	06		1968	$\beta^- =100$
<sup>194</sup> Pt	-34760.1 0.5		STABLE	0 <sup>+</sup>	06		1935	IS=32.86 40
<sup>194</sup> Au	-32211.9 2.1		38.02 h 0.10	1 <sup>-</sup>	06		1948	$\beta^+ =100$
<sup>194</sup> Au <sup>m</sup>	-32104.5 2.2	107.4 0.5	600 ms 8	(5 <sup>+</sup> )	06		1975	IT=100
<sup>194</sup> Au <sup>n</sup>	-31736.1 2.2	475.8 0.6	420 ms 10	(11 <sup>-</sup> )	06		1953	IT=100
<sup>194</sup> Hg	-32183.9 2.9		447 y 28	0 <sup>+</sup>	06	15Do01	T 1962	$e=100$
<sup>194</sup> Tl	-26937 14		33.0 m 0.5	2 <sup>-</sup>	06		1960	$\beta^+ =100$ ; $\alpha < 1e-7$
<sup>194</sup> Tl <sup>m</sup>	-26677 4	260 14	32.8 m 0.2	7 <sup>(+)</sup>	06	13Ba41	J 1960	$\beta^+ =100$
<sup>194</sup> Pb	-24208 17		10.7 m 0.6	0 <sup>+</sup>	06		1960	$\beta^+ =100$ ; $\alpha =7.3e-6$ 29
<sup>194</sup> Pb <sup>m</sup>	-21580 17	2628.1 0.4	370 ns 13	12 <sup>+</sup>	06	FGK128	J 1972	IT=100
<sup>194</sup> Pb <sup>n</sup>	-21275 17	2933.0 0.4	133 ns 7	11 <sup>-</sup>	06		1986	IT=100
<sup>194</sup> Bi	-16029 6		95 s 3	(3 <sup>+</sup> )	06		1971	$\beta^+ \approx 100$ ; $\alpha =0.46$ 25
<sup>194</sup> Bi <sup>m</sup>	-15880 50	150 50	125 s 2	(6 <sup>+</sup> , 7 <sup>+</sup> )	06		1976	$\beta^+ \approx 100$ ; $\alpha ?$
<sup>194</sup> Bi <sup>n</sup>	-15849 8	180 10	115 s 4	(10 <sup>-</sup> )	06		1988	$\beta^+ \approx 100$ ; $\alpha =0.20$ 7
<sup>194</sup> Po	-11005 13		392 ms 4	0 <sup>+</sup>	06		1967	$\alpha \approx 100$ ; $\beta^+ ?$
<sup>194</sup> Po <sup>m</sup>	-8480 13	2525.2 0.8	15 $\mu$ s 2	(11 <sup>-</sup> )	06		1999	IT=100
<sup>194</sup> At	-720 25		286 ms 7	(4 <sup>-</sup> , 5 <sup>-</sup> )	06	13An03	TD 2009	$\alpha \approx 100$ ; $\beta^+ =8.3\%$ ; $\beta^+ SF=0.032$
<sup>194</sup> At <sup>m</sup>	-740 30	-20 40	323 ms 7	(9 <sup>-</sup> , 10 <sup>-</sup> )	06	13An03	T 1984	$\alpha \approx 100$ ; $\beta^+ =8.3\%$ ; $\beta^+ SF=0.032$
<sup>194</sup> Rn	5723 17		780 $\mu$ s 160	0 <sup>+</sup>	07		2006	$\alpha \approx 100$ ; $\beta^+ ?$
* <sup>194</sup> Re	T : other 09Ku28=1.0(0.5) withdrawn by authors in 14Ku23							
* <sup>194</sup> Re <sup>m</sup>	T : from 12Al105 two exc. isomers with 25(8)s 100(10)s; could be exchanged							
* <sup>194</sup> Re <sup>p</sup>	D : only 86.3 keV $\gamma$ is seen, but not those seen in <sup>194</sup> Re <sup>q</sup>							
* <sup>194</sup> Re <sup>q</sup>	I : decaying by delayed $\gamma$ -rays of 464, 148, 128							
* <sup>194</sup> Hg	T : average 81Ho18=477(32) 79Pr15=358(55), values corrected in 15Do01 for							
* <sup>194</sup> Hg	T : the new branching intensity of the 328.5 g ray.							
* <sup>194</sup> Pb <sup>m</sup>	J : E2 to 10 <sup>+</sup> ; magnetic moment							
* <sup>194</sup> Pb <sup>n</sup>	J : E2 to 9 <sup>-</sup> ; magnetic moment							
* <sup>194</sup> At	T : 13An03 supersedes 09An11=253(10) D : 13An03=0.065 8 for both isomers							
* <sup>194</sup> At	J : favored $\alpha$ -decay to (5 <sup>-</sup> ) isomer in <sup>190</sup> Bi							
* <sup>194</sup> At <sup>m</sup>	T : 13An03=323(7) supersedes 09An11=310(8); other 13Ny01=300(+50-40)							
* <sup>194</sup> At <sup>n</sup>	J : favored $\alpha$ -decay to (10 <sup>-</sup> ) isomer in <sup>190</sup> Bi							
<sup>195</sup> W	-21010# 300#		3# s (>300 ns)	5/2 <sup>-</sup> #		12Ku26	I 2012	$\beta^- ?$
<sup>195</sup> Re	-25580# 300#		6 s 1	5/2 <sup>+</sup> #	14		2008	$\beta^- =100$
<sup>195</sup> Os	-29510 60		6.5 m 1.1	3/2 <sup>-</sup> #	14		2004	$\beta^- ?$
<sup>195</sup> Os <sup>m</sup>	-29060 60	454 10	2.0 h 1.7	13/2 <sup>+</sup> #	14	12Re19	ED 2012	$\beta^- ?$ ; IT=?
<sup>195</sup> Ir	-31692.3 1.3		2.29 h 0.17	3/2 <sup>+</sup>	14		1952	$\beta^- =100$
<sup>195</sup> Ir <sup>m</sup>	-31592 5	100 5	3.67 h 0.08	11/2 <sup>-</sup>	14		1968	$\beta^- =95$ 5; IT=5 5
<sup>195</sup> Ir <sup>n</sup>	-29338 6	2354 6	4.4 $\mu$ s 0.6	(27/2 <sup>+</sup> )		11St21	ETJ 2011	IT=100
<sup>195</sup> Pt	-32793.8 0.5		STABLE	1/2 <sup>-</sup>	14		1935	IS=33.78 24
<sup>195</sup> Pt <sup>m</sup>	-32534.7 0.5	259.077 0.023	4.010 d 0.005	13/2 <sup>+</sup>	14		1960	IT=100
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...												
<sup>195</sup> Au	-32567.0	1.1		186.01 d	0.06		3/2 <sup>+</sup>	14	1948	$\epsilon=100$		
<sup>195</sup> Au <sup>m</sup>	-32248.4	1.1	318.58	0.04			11/2 <sup>-</sup>	14	1952	IT=100		
<sup>195</sup> Au <sup>n</sup>	-30067	20	2500	20			31/2 <sup>(-)</sup>	14	13Dr01 ET	2013	IT=100	
<sup>195</sup> Hg	-31013	23					1/2 <sup>-</sup>	14	15Do01 T	1952	$\beta^+=100$	
<sup>195</sup> Hg <sup>m</sup>	-30837	23	176.07	0.04			13/2 <sup>+</sup>	14	15Do01 T	1951	IT=54.2 20; $\beta^+=45.8$ 20	
<sup>195</sup> Tl	-28155	11					1/2 <sup>+</sup>	14	1955	$\beta^+=100$		
<sup>195</sup> Tl <sup>m</sup>	-27672	11	482.63	0.17			9/2 <sup>-</sup>	14	1957	IT=100		
<sup>195</sup> Pb	-23708	18					3/2 <sup>-</sup>	14	1957	$\beta^+=100$		
<sup>195</sup> Pb <sup>m</sup>	-23505	18	202.9	0.7			13/2 <sup>(+)</sup>	14	91Gr12 E	1957	$\beta^+=100$	
<sup>195</sup> Pb <sup>n</sup>	-21949	18	1759.0	0.7			(21/2 <sup>-</sup> )	14	1976	IT=100		
<sup>195</sup> Bi	-18026	5					9/2 <sup>(-)</sup>	14	GAu14b J	1971	$\beta^+\approx 100$ ; $\alpha=0.03$ 2	
<sup>195</sup> Bi <sup>m</sup>	-17626	8	399	6	AD		(1/2 <sup>+</sup> )	14	GAu14b J	1974	$\beta^+=67$ 17; $\alpha=33$ 17	
<sup>195</sup> Bi <sup>n</sup>	-15631	5	2395.5	0.5			(29/2 <sup>-</sup> )	14	15Ro20 J	2003	IT=100	
<sup>195</sup> Bi <sup>p</sup>	-14690	5	3336	2			(31/2 <sup>-</sup> )	15	15Ro20 ETJ	2015	IT=100	
<sup>195</sup> Po	-11060	40					(3/2 <sup>-</sup> )	15	13Se03 J	1967	$\alpha=94$ 4; $\beta^+?$	
<sup>195</sup> Po <sup>m</sup>	-10965	28	90	50	AD		(13/2 <sup>+</sup> )	15	13Se03 J	1967	$\alpha\approx 90$ ; $\beta^+\approx 10$ ; IT<0.01	
<sup>195</sup> At	-3470	10					1/2 <sup>+</sup>	14	1999	$\alpha\approx 100$ ; $\beta^+?$		
<sup>195</sup> At <sup>m</sup>	-3441	8	29	7	AD		(7/2 <sup>-</sup> )	14	1995	$\alpha=?$ ; IT=12 4; $\beta^+?$		
<sup>195</sup> At <sup>p</sup>	-3370#	40#	100#	40#			(13/2 <sup>+</sup> )	14	13Uu01 J		IT?	
<sup>195</sup> Rn	5050	50					3/2 <sup>-</sup>	14	2001	$\alpha\approx 100$		
<sup>195</sup> Rn <sup>m</sup>	5131	17	80	50	AD	*	6 ms 3	14	2001	$\alpha\approx 100$		
* <sup>195</sup> Os <sup>m</sup>	T: symmetrized from 32(+154-16) m for q=76 <sup>+</sup> (bare ion)									**		
* <sup>195</sup> Ir <sup>m</sup>	E: 268.4, 404.4, 476.4, 537.8, 566.7 $\gamma$ s in cascade to <sup>195</sup> Ir <sup>m</sup>									**		
* <sup>195</sup> Au <sup>n</sup>	E: 13Dr01=2460.9 + x, x=40#(20#) estimated by NUBASE T: $\tau=18.6(0.3)$									**		
* <sup>195</sup> Hg	T: average 15Do01=10.84(0.03) 01Li17=10.53(0.03), Birge ratio B=7.3									**		
* <sup>195</sup> Hg <sup>m</sup>	T: average 15Do01=41.6(0.2) 73Vi09=41.6(0.8)									**		
* <sup>195</sup> Pb <sup>m</sup>	J: same as <sup>199</sup> Po <sup>m</sup> and <sup>203</sup> Rn <sup>m</sup> , from $\alpha$ decay									**		
* <sup>195</sup> Bi <sup>m</sup>	J: spins of ground-state and of isomer derived from $\alpha$ decay to daughter									**		
* <sup>195</sup> Bi <sup>p</sup>	E: uncertainty estimated by NUBASE									**		
* <sup>195</sup> At <sup>m</sup>	E: ENSDF14=33.0(1.0) is erroneous									**		
* <sup>195</sup> At <sup>p</sup>	E: estimated 70#(40#) above <sup>195</sup> At <sup>m</sup> ; 13Ny01 estimated upper limit is 130 keV									**		
* <sup>195</sup> Rn	T: symmetrized from 01Uu01=6(+3-2)									**		
* <sup>195</sup> Rn <sup>m</sup>	T: symmetrized from 01Uu01=5(+3-2)									**		
<sup>196</sup> W	-18880#	400#					3# s (>300 ns)	0 <sup>+</sup>	13	12Ku26 I	2012	$\beta^-?$
<sup>196</sup> Re	-22540#	300#					2.4 s 1.5		13		2008	$\beta^-?$
<sup>196</sup> Re <sup>m</sup>	-22420#	300#	120#	40#			3.6 $\mu$ s 0.6		11St21 T	2009	IT=100	
<sup>196</sup> Os	-28280	40					34.9 m 0.2	0 <sup>+</sup>	07		1977	$\beta^-=100$
<sup>196</sup> Ir	-29440	40					52 s 1	(0 <sup>-</sup> )	07		1966	$\beta^-=100$
<sup>196</sup> Ir <sup>m</sup>	-29227	20	210	40	BD		1.40 h 0.02	(10, 11 <sup>-</sup> )	07		1959	$\beta^-\approx 100$ ; IT<0.3
<sup>196</sup> Pt	-32644.5	0.5					STABLE	0 <sup>+</sup>	07		1935	IS=25.21 34
<sup>196</sup> Au	-31138.7	3.0					6.1669 d 0.0006	2 <sup>-</sup>	07		1937	$\beta^+=92.8$ 8; $\beta^-=7.2$ 8
<sup>196</sup> Au <sup>m</sup>	-31054	3	84.656	0.020			8.1 s 0.2	(5 <sup>+</sup> )	07		1971	IT=100
<sup>196</sup> Au <sup>n</sup>	-30543	3	595.66	0.04			9.6 h 0.1	12 <sup>-</sup>	07		1960	IT=100
<sup>196</sup> Hg	-31825.9	2.9					STABLE (>2.5 Ey)	0 <sup>+</sup>	07	90Bu28 T	1930	IS=0.15 1; 2 $\beta^+?$
<sup>196</sup> Tl	-27497	12					1.84 h 0.03	2 <sup>-</sup>	07		1955	$\beta^+=100$
<sup>196</sup> Tl <sup>m</sup>	-27103	12	394.2	0.5			1.41 h 0.02	7 <sup>(+)</sup>	07	13Ba41 J	1960	$\beta^+=96.2$ 4; IT=3.8 4
<sup>196</sup> Pb	-25348	8					37 m 3	0 <sup>+</sup>	07		1957	$\beta^+=100$ ; $\alpha<3e-5$
<sup>196</sup> Pb <sup>m</sup>	-23610	8	1738.27	0.12			< 1 $\mu$ s	4 <sup>+</sup>	07		1973	IT=100
<sup>196</sup> Pb <sup>n</sup>	-23550	8	1797.51	0.14			140 ns 14	5 <sup>-</sup>	07		1973	IT=100
<sup>196</sup> Pb <sup>p</sup>	-22653	8	2694.6	0.3			270 ns 4	12 <sup>+</sup>	07		1973	IT=100
<sup>196</sup> Bi	-18009	24					5.1 m 0.2	(3 <sup>+</sup> )	07		1976	$\beta^+\approx 100$ ; $\alpha=0.00115$ 34
<sup>196</sup> Bi <sup>m</sup>	-17843	25	166.4	2.9	AD		0.6 s 0.5	(7 <sup>+</sup> )	07		1987	IT=?; $\beta^+?$
<sup>196</sup> Bi <sup>n</sup>	-17737	25	272	3	AD		4.00 m 0.05	(10 <sup>-</sup> )	07		1987	$\beta^+=74.2$ 25; IT=25.8 25; $\alpha=0.00038$ 10
<sup>196</sup> Po	-13473	14					5.56 s 0.09	0 <sup>+</sup>	07	05Uu02 T	1967	$\alpha\approx 98$ ; $\beta^+\approx 2$
<sup>196</sup> Po <sup>m</sup>	-10979	14	2493.9	0.4			856 ns 17	11 <sup>-</sup>	07		1995	IT=100
<sup>196</sup> At	-3910	30					388 ms 7	(3 <sup>+</sup> )	07	93An11 D	1967	$\alpha=?$ ; $\beta^+=5\#$ ; $\beta^+$ SF=0.088
<sup>196</sup> At <sup>m</sup>	-3950	18	-40	40	AD	*	20# ms	(10 <sup>-</sup> )	07	96En01 D	1996	$\alpha\approx 100$
<sup>196</sup> At <sup>n</sup>	-3750	30	157.9	0.1			11 $\mu$ s 2	(5 <sup>+</sup> )	07		2000	IT=100
<sup>196</sup> Rn	1971	14					4.7 ms 1.1	0 <sup>+</sup>	07		1995	$\alpha\approx 100$ ; $\beta^+=0.06\#$
* <sup>196</sup> Re	T: symmetrized from 14Ku23=3(+1-2)									**		
* <sup>196</sup> Re <sup>m</sup>	D: only Kx-rays observed; E>72 keV (K-shell binding energy)									**		
* <sup>196</sup> Po	T: average 05Uu02=5.1(+3.1-1.4) 97Pu01=5.5(0.1) 93Wu04=5.8(0.2)									**		
* <sup>196</sup> Po	T: other not used: 10He25=4.1(+5.6-1.5) ms									**		
* <sup>196</sup> At	J: same as <sup>192</sup> Bi, from $\alpha$ decay T: also 14Ka23=350(90)									**		
* <sup>196</sup> At <sup>m</sup>	I: level not accepted in ENSDF									**		
* <sup>196</sup> Rn	T: symmetrized from 4.4(+1.3-0.9)									**		

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>197</sup> W	-15140# 400#		1# s (>300 ns)	5/2 <sup>-</sup> #	13	12Ku26 I	2012	$\beta^-$ ?
<sup>197</sup> Re	-20500# 300#		300# ms (>300 ns)	5/2 <sup>+</sup> #	13		2009	$\beta^-$ ?
<sup>197</sup> Os	-25310# 200#		2.8 m 0.6	5/2 <sup>-</sup> #	09		2003	$\beta^-$ =100
<sup>197</sup> Ir	-28264 20		5.8 m 0.5	(3/2 <sup>+</sup> )	05		1952	$\beta^-$ =100
<sup>197</sup> Ir <sup>m</sup>	-28149 21 115 5		8.9 m 0.3	(11/2 <sup>-</sup> )	05		1976	$\beta^-$ ≈100; IT=0.25 10
<sup>197</sup> Ir <sup>m</sup>	-27860# 200# 400# 200#		30 μs 8			05Ca02 T	2005	IT=100
<sup>197</sup> Pt	-30419.7 0.5		19.8915 h 0.0019	1/2 <sup>-</sup>	05		1936	$\beta^-$ =100
<sup>197</sup> Pt <sup>m</sup>	-30020.1 0.5 399.59 0.20		95.41 m 0.18	13/2 <sup>+</sup>	05		1941	IT=96.7 4; $\beta^-$ =3.3 4
<sup>197</sup> Au	-31139.7 0.5		STABLE	3/2 <sup>+</sup>	05		1935	IS=100.
<sup>197</sup> Au <sup>m</sup>	-30730.6 0.5 409.15 0.08		7.73 s 0.06	11/2 <sup>-</sup>	05		1945	IT=100
<sup>197</sup> Au <sup>n</sup>	-28607.2 1.1 2532.5 1.0		150 ns 5	27/2 <sup>+</sup> #		06Wh02 ETJ	2006	IT=100
<sup>197</sup> Hg	-30540 3		64.94 h 0.07	1/2 <sup>-</sup>	05	01Li17 T	1941	$\epsilon$ =100
<sup>197</sup> Hg <sup>m</sup>	-30241 3 298.93 0.08		23.8 h 0.1	13/2 <sup>+</sup>	05		1943	IT=91.4 7; $\epsilon$ =8.6 7
<sup>197</sup> Tl	-28342 16		2.84 h 0.04	1/2 <sup>+</sup>	05		1955	$\beta^+$ =100
<sup>197</sup> Tl <sup>m</sup>	-27734 16 608.22 0.08		540 ms 10	9/2 <sup>-</sup>	05	13Ba41 J	1953	IT=100
<sup>197</sup> Pb	-24745 5		8.1 m 1.7	3/2 <sup>-</sup>	05		1955	$\beta^+$ =100
<sup>197</sup> Pb <sup>m</sup>	-24426 5 319.31 0.11		42.9 m 0.9	13/2 <sup>+</sup>	05		1957	$\beta^+$ =81 2; IT=19 2
<sup>197</sup> Pb <sup>n</sup>	-22831 5 1914.10 0.25		1.15 μs 0.20	21/2 <sup>-</sup>	05		1978	IT=100
<sup>197</sup> Bi	-19687 8		9.33 m 0.50	(9/2 <sup>-</sup> )	05		1971	$\beta^+$ =100; $\alpha$ =1e-4#
<sup>197</sup> Bi <sup>m</sup>	-19155 8 533 12 AD		5.04 m 0.16	(1/2 <sup>+</sup> )	05		1966	$\alpha$ =55 40; $\beta^+$ =45 40; IT<0.3
<sup>197</sup> Bi <sup>n</sup>	-17284 14 2403 12		263 ns 13	(29/2 <sup>-</sup> )	05	86Ch01 TJD	1986	IT=100
<sup>197</sup> Bi <sup>p</sup>	-16758 8 2929.5 0.5		209 ns 30	(31/2 <sup>-</sup> )	05	86Ch01 TJD	1986	IT=100
<sup>197</sup> Po	-13360 50		53.6 s 0.9	(3/2 <sup>-</sup> )	05	93Wa04 T	1965	$\beta^+$ ?; $\alpha$ =44 7
<sup>197</sup> Po <sup>m</sup>	-13130# 90# 230# 80#		25.8 s 0.1	(13/2 <sup>+</sup> )	05	93Wa04 T	1967	$\alpha$ =84 9; $\beta^+$ ?; IT=0.01#
<sup>197</sup> At	-6355 8		388.2 ms 5.6	(9/2 <sup>-</sup> )	05	05De01 T	1967	$\alpha$ =96.1 12; $\beta^+$ =3.9 12
<sup>197</sup> At <sup>m</sup>	-6311 9 45 8 AD *		2.0 s 0.2	(1/2 <sup>+</sup> )	05		1985	$\alpha$ ≈100; $\beta^+$ ?; IT<0.004; $\beta^+$ ?
<sup>197</sup> At <sup>p</sup>	-6044 8 310.7 0.2		1.3 μs 0.2	(13/2 <sup>+</sup> )		08An11 ETJ	1999	IT=100
<sup>197</sup> Rn	1510 16		54 ms 6	(3/2 <sup>-</sup> )	05	08An05 T	1995	$\alpha$ ≈100; $\beta^+$ ?
<sup>197</sup> Rn <sup>m</sup>	1709 16 199 11		25.6 ms 2.5	(13/2 <sup>+</sup> )	05	08An05 T	1996	$\alpha$ ≈100; $\beta^+$ ?
<sup>197</sup> Fr	10250 50		2.33 ms 1.88	7/2 <sup>-</sup> #	14	13Ka16 T	2013	$\alpha$ ≈100
* <sup>197</sup> Hg	T : 66El09=64.14(0.05) strongly conflicting: Birge ratio would be $B=9.3$							
* <sup>197</sup> Tl <sup>m</sup>	J : also in 12Bi.A							
* <sup>197</sup> Bi	I : ENSDF'05 reported an isomer at 2129.3(0.4) keV, 204(18) ns, (23/2 <sup>-</sup> ),							
* <sup>197</sup> Bi	I : not trusted by NUBASE, see fig.3 in 86Ch01							
* <sup>197</sup> Bi <sup>m</sup>	J : $\alpha$ decay to <sup>193</sup> Tl ground-state							
* <sup>197</sup> Bi <sup>n</sup>	T : more recent 95Zh36=252.6(38.7) outweighed, not used							
* <sup>197</sup> Bi <sup>n</sup>	E : 95Zh36=2383.1 + x, with x<40 keV; 86Ch01=2360.4 + x is the same level							
* <sup>197</sup> Bi <sup>n</sup>	E : but authors mis-assigned the 97 keV $\gamma$ -ray, see Fig.1 of 95Zh36							
* <sup>197</sup> Po	T : average 93Wa04=53(1) 71Ho01=60(6) 67Le21=58(3) 67Si09=52(4); other not							
* <sup>197</sup> Po	T : used 96Ta18=84(16)							
* <sup>197</sup> Po <sup>m</sup>	T : others not used 71Ho01=27(3) 67Le21=29(9) 67Si09=26(2);							
* <sup>197</sup> Po <sup>m</sup>	T : also 10He25=14.45(+14.45-4.9) ms for 3 events, strongly conflicting							
* <sup>197</sup> At	T : average 05De01=390(16) 99Sm07=388(6); also 14Ka23=354(+17-15)							
* <sup>197</sup> At <sup>m</sup>	T : also 14Ka23=2.8(+3.8-1.0)							
* <sup>197</sup> At <sup>n</sup>	T : other 99Sm07=5.5(1.4)							
* <sup>197</sup> Rn	T : symmetrized from 08An05=53(+7-5) J : from $\alpha$ decay to <sup>193</sup> Po							
* <sup>197</sup> Rn <sup>m</sup>	T : symmetrized from 08An05=25(+3-2) J : from $\alpha$ decay to <sup>193</sup> Po <sup>m</sup>							
* <sup>197</sup> Rn <sup>n</sup>	T : others 05Uu02=30(+150-15) 96En02=19(+8-4) 95Mo14=18(+9-5)							
* <sup>197</sup> Fr	T : symmetrized from 13Ka16=0.6(+30-3)							
<sup>198</sup> Re	-17140# 400#		300# ms (>300 ns)		16	09St16 I	2009	$\beta^-$ ?; $\beta^-$ -n=0#
<sup>198</sup> Os	-23840# 200#		1# m (>300 ns)	0 <sup>+</sup>	16	09Po02 I	2008	$\beta^-$ ?
<sup>198</sup> Ir	-25820# 200#		8 s 1		16		1973	$\beta^-$ =100
<sup>198</sup> Pt	-29904.0 2.1		STABLE	0 <sup>+</sup>	16		1935	IS=7.36 13; 2 $\beta^-$ ?; $\alpha$ ?
<sup>198</sup> Au	-29580.8 0.5		2.6941 d 0.0002	2 <sup>-</sup>	16		1937	$\beta^-$ =100
<sup>198</sup> Au <sup>m</sup>	-29268.6 0.5 312.2227 0.0020		124 ns 4	5 <sup>+</sup>	16		1968	IT=100
<sup>198</sup> Au <sup>n</sup>	-28768.9 1.6 811.9 1.5		2.272 d 0.016	12 <sup>-</sup>	16	FGK128 J	1972	IT=100
<sup>198</sup> Hg	-30954.3 0.5		STABLE	0 <sup>+</sup>	16		1925	IS=9.97 20
<sup>198</sup> Tl	-27529 8		5.3 h 0.5	2 <sup>-</sup>	16		1949	$\beta^+$ =100
<sup>198</sup> Tl <sup>m</sup>	-26985 8 543.6 0.4		1.87 h 0.03	7 <sup>+</sup>	16		1949	$\beta^+$ =55.9 23; IT=44.1 23
<sup>198</sup> Tl <sup>n</sup>	-26842 8 686.8 0.5		150 ns 40	(5,7,9) <sup>+</sup>	16		1977	IT=100
<sup>198</sup> Tl <sup>p</sup>	-26787 8 742.4 0.4		32.1 ms 1.0	10 <sup>-</sup>	16	FGK128 J	1975	IT=100
<sup>198</sup> Pb	-26067 9		2.4 h 0.1	0 <sup>+</sup>	16		1955	$\beta^+$ =100
<sup>198</sup> Pb <sup>m</sup>	-23926 9 2141.4 0.4		4.19 μs 0.10	7 <sup>-</sup>	16	FGK128 J	1972	IT=100
<sup>198</sup> Pb <sup>n</sup>	-23836 9 2231.4 0.5		137 ns 10	9 <sup>-</sup>	16	FGK128 J	1989	IT=100
<sup>198</sup> Pb <sup>p</sup>	-23245 9 2821.7 0.6		212 ns 4	12 <sup>+</sup>	16	FGK128 J	1973	IT=100

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>198</sup> Bi	-19369	28	10.3 m	0.3	3 <sup>(+)</sup>	16 16Ly01 J	1950	$\beta^+=100$
<sup>198</sup> Bi <sup>m</sup>	-19085	28 280	40 MD	11.6 m	0.3	7 <sup>(+)</sup>	16 16Ly01 J	1992 $\beta^+=100$
<sup>198</sup> Bi <sup>n</sup>	-18837	28 530	40 MD	7.7 s	0.5	10 <sup>(-)</sup>	16 16Ly01 J	1972 IT=100 *
<sup>198</sup> Po	-15473	17		1.760 m	0.024	0 <sup>+</sup>	16	1965 $\alpha=57.2; \beta^+=43.2$
<sup>198</sup> Po <sup>m</sup>	-12907	17 2565.92	0.20	200 ns	20	11 <sup>-</sup>	16	1990 IT=100
<sup>198</sup> Po <sup>n</sup>	-12730	50 2740	50	750 ns	50	12 <sup>+</sup>	16	1990 IT ?
<sup>198</sup> At	-6715	6		3.0 s	0.1	(3 <sup>+</sup> )	16 14Ka23 T	1967 $\alpha>94; \beta^+ ?$
<sup>198</sup> At <sup>m</sup>	-6430	8 284	10 AD	1.21 s	0.06	(10 <sup>-</sup> )	16	1967 $\alpha=84.16; \beta^+ ?$
<sup>198</sup> Rn	-1230	13		65 ms	3	0 <sup>+</sup>	16	1984 $\alpha=?; \beta^+=1\#$
<sup>198</sup> Fr	9570	30		15 ms	3	low	16	2013 $\alpha\approx 100$
<sup>198</sup> Fr <sup>m</sup>	9580	40 0	50	1.1 ms	0.7	high	16	2013 $\alpha\approx 100$
* <sup>198</sup> Re	I : other 12Ku26>300 ns **							
* <sup>198</sup> Pt	T : 52Fr23 : 0v- $\beta\beta$ >320 Ty **							
* <sup>198</sup> Au <sup>n</sup>	J : M4 to 8 <sup>+</sup> ; magnetic moment **							
* <sup>198</sup> Tl <sup>p</sup>	J : E3 to 7 <sup>+</sup> **							
* <sup>198</sup> Pb <sup>m</sup>	J : E2 to 5 <sup>-</sup> ; magnetic moment **							
* <sup>198</sup> Pb <sup>n</sup>	J : E2 to 7 <sup>-</sup> **							
* <sup>198</sup> Pb <sup>p</sup>	J : E2 to 10 <sup>+</sup> ; magnetic moment **							
* <sup>198</sup> Bi <sup>n</sup>	E : 248.5(0.5) keV above <sup>198</sup> Bi <sup>m</sup> , from 92Hu04 J : E3 to (7 <sup>+</sup> ) **							
<sup>199</sup> Re	-14860#	400#		100# ms	(>300 ns)	5/2 <sup>+</sup> #	13 12Ku26 I	2012 $\beta^- ?$
<sup>199</sup> Os	-20480#	200#		6 s	3	5/2 <sup>-</sup> #	07 14Ku23 T	2008 $\beta^-=100$ *
<sup>199</sup> Ir	-24400	40		7 s	5	3/2 <sup>+</sup> #	07 14Ku23 T	1993 $\beta^- ?$ *
<sup>199</sup> Ir <sup>m</sup>	-24270#	60# 130#	40#	235 ns	90	11/2 <sup>-</sup> #	07	2005 IT=100 *
<sup>199</sup> Pt	-27388.7	2.2		30.80 m	0.21	5/2 <sup>-</sup>	07	1937 $\beta^-=100$
<sup>199</sup> Pt <sup>m</sup>	-26964.7	3.0 424	2	13.6 s	0.4	(13/2 <sup>+</sup> )	07	1959 IT=100
<sup>199</sup> Au	-29093.7	0.5		3.139 d	0.007	3/2 <sup>+</sup>	07	1937 $\beta^-=100$
<sup>199</sup> Au <sup>m</sup>	-28544.8	0.5 548.9405	0.0021	440 $\mu$ s	30	(11/2 <sup>-</sup> )	07	1968 IT=100
<sup>199</sup> Hg	-29546.1	0.5		STABLE		1/2 <sup>-</sup>	07	1925 IS=16.87 22
<sup>199</sup> Hg <sup>m</sup>	-29013.6	0.5 532.48	0.10	42.67 m	0.09	13/2 <sup>+</sup>	07	1948 IT=100
<sup>199</sup> Tl	-28059	28		7.42 h	0.08	1/2 <sup>+</sup>	07	1949 $\beta^+=100$
<sup>199</sup> Tl <sup>m</sup>	-27310	28 748.87	0.06	28.4 ms	0.2	9/2 <sup>-</sup>	07	1963 IT=100
<sup>199</sup> Pb	-25232	10		90 m	10	3/2 <sup>-</sup>	07	1950 $\beta^+=100$
<sup>199</sup> Pb <sup>m</sup>	-24803	10 429.5	2.7	12.2 m	0.3	(13/2 <sup>+</sup> )	07	1955 IT=93; $\beta^+=7$ *
<sup>199</sup> Pb <sup>n</sup>	-22668	10 2563.8	2.7	10.1 $\mu$ s	0.2	(29/2 <sup>-</sup> )	07	1981 IT=100 *
<sup>199</sup> Bi	-20798	11		27 m	1	9/2 <sup>-</sup>	07	1950 $\beta^+=100$
<sup>199</sup> Bi <sup>m</sup>	-20131	11 667	3	24.70 m	0.15	(1/2 <sup>+</sup> )	07	1950 $\beta^+=?; IT<2; \alpha\approx 0.01$
<sup>199</sup> Bi <sup>n</sup>	-18851	18 1947	14	100 ns	30	25/2 <sup>+</sup> #	07	1974 IT=100 *
<sup>199</sup> Bi <sup>p</sup>	-18250	18 2548	14	168 ns	13	29/2 <sup>-</sup> #	07	1985 IT=100 *
<sup>199</sup> Po	-15208	18		5.47 m	0.15	(3/2 <sup>-</sup> )	07 13Se03 J	1965 $\beta^+=92.5.3; \alpha=7.5.3$
<sup>199</sup> Po <sup>m</sup>	-14897	18 311.9	2.7	4.17 m	0.05	13/2 <sup>(+)</sup>	07	1964 $\beta^+=73.5.10; \alpha=24.1; IT=2.5.10$ *
<sup>199</sup> At	-8823	5		7.02 s	0.12	9/2 <sup>(-)</sup>	07 05De01 T	1967 $\alpha=89.6; \beta^+ ?$
<sup>199</sup> At <sup>m</sup>	-8579	5 244.0	1.0	273 ms	9	(1/2 <sup>+</sup> )	14Au03 TJD	2013 IT $\approx$ 100; $\alpha=1\#$ *
<sup>199</sup> At <sup>n</sup>	-8250	5 572.9	0.1	70 ns	20	(13/2 <sup>+</sup> )	07 10Ja05 ETJ	2000 IT=100
<sup>199</sup> At <sup>p</sup>	-6530	5 2293.4	0.5	800 ns	50	(29/2 <sup>+</sup> )	07 10Ja05 ETJ	2010 IT=100
<sup>199</sup> Rn	-1500	40		590 ms	30	(3/2 <sup>-</sup> )	07 05Uu02 J	1980 $\alpha=?; \beta^+=6\#$ *
<sup>199</sup> Rn <sup>m</sup>	-1337	29 160	50	310 ms	20	(13/2 <sup>+</sup> )	07 05Uu02 J	1981 $\alpha=?; \beta^+=3\#$
<sup>199</sup> Fr	6771	14		6.6 ms	2.2	1/2 <sup>+</sup> #	07 13Ka16 T	1999 $\alpha\approx 100; \beta^+ ?$ *
<sup>199</sup> Fr <sup>m</sup>	6817	10 45	13	6.5 ms	0.9	7/2 <sup>-</sup> #	13Ka16 T	2013 $\alpha\approx 100; \beta^+ ?$ *
<sup>199</sup> Fr <sup>n</sup>	7020#	50# 250#	50#	2.2 ms	1.2		13Uu01 TD	2013 $\alpha=?; \beta^+ ?$ *
* <sup>199</sup> Os	T : symmetrized from 14Ku23=14Mo15=5(+4-2) **							
* <sup>199</sup> Ir	T : symmetrized from 14Ku23=14Mo15=6(+5-4) **							
* <sup>199</sup> Ir <sup>m</sup>	T : range 80-390 ns **							
* <sup>199</sup> Pb <sup>m</sup>	E : 424.8(0.2) + x; x < 9.3 keV D : from 78Le.A **							
* <sup>199</sup> Pb <sup>n</sup>	E : 2559.1(0.4) + x; x < 9.3 keV **							
* <sup>199</sup> Bi <sup>n</sup>	E : 1922.3 + x; x < 50 in ENSDF'07 **							
* <sup>199</sup> Bi <sup>p</sup>	E : 2523.2 + x; x < 50 in ENSDF'07 **							
* <sup>199</sup> Po <sup>m</sup>	J : same as <sup>203</sup> Rn <sup>m</sup> , from $\alpha$ decay; also 13Se03=(13/2 <sup>+</sup> ) **							
* <sup>199</sup> At	T : average 12Fo09=6.7(0.5) 05De01=6.92(0.13) 05Uu02=7.8(0.4) 67Tr06=7.2(0.5) **							
* <sup>199</sup> At	J : spins of ground-state derived from $\alpha$ decay to daughter **							
* <sup>199</sup> At	D : symmetrized from $\alpha=92(+3-8)\%$ **							
* <sup>199</sup> At <sup>m</sup>	T : 14Au03=273(9) 13Ja06=310(80) **							
* <sup>199</sup> Rn	T : others 14Ka23=340(+280-110) **							
* <sup>199</sup> Fr	T : average 13Ka16=4.5(+3.1-1.3) 99Ta20=12(+10-4) J : same as <sup>195</sup> At **							
* <sup>199</sup> Fr <sup>m</sup>	T : average 13Ka16=6.2(+1.1-0.8) 13Uu01=7(+3-2) J : same as <sup>195</sup> Atm **							
* <sup>199</sup> Fr <sup>n</sup>	T : symmetrized from 13Uu01=1.6(+1.6-6) **							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)					
<sup>200</sup> Os	-18780#	300#			7 s	4	0 <sup>+</sup>	08 14Ku23	T	2005	$\beta^- = 100$	*			
<sup>200</sup> Ir	-21610#	200#			43 s	6	(2 <sup>-</sup> , 3 <sup>-</sup> )	11 14Mo15	T	2008	$\beta^- = 100$	*			
<sup>200</sup> Pt	-26599	20			12.6 h	0.3	0 <sup>+</sup>	07		1957	$\beta^- = 100$				
<sup>200</sup> Au	-27240	27			48.4 m	0.3	1 <sup>(-)</sup>	07		1951	$\beta^- = 100$				
<sup>200</sup> Au <sup>m</sup>	-26233	26	1010	40	BD		18.7 h	0.5		07	1968	$\beta^- = 82.2$ ; IT=18.2			
<sup>200</sup> Hg	-29503.3	0.5			STABLE		0 <sup>+</sup>	07		1925	IS=23.10 19				
<sup>200</sup> Tl	-27047	6			26.1 h	0.1	2 <sup>-</sup>	07		1949	$\beta^+ = 100$				
<sup>200</sup> Tl <sup>m</sup>	-26293	6	753.6	0.24			34.0 ms	0.9		07	1963	IT=100			
<sup>200</sup> Tl <sup>n</sup>	-26285	6	762.00	0.24			330 ns	50		07	1972	IT=100			
<sup>200</sup> Pb	-26251	11			21.5 h	0.4	0 <sup>+</sup>	07		1950	$\epsilon = 100$				
<sup>200</sup> Pb <sup>m</sup>	-24068	11	2183.3	1.1			448 ns	12		07	1972	IT=100			
<sup>200</sup> Pb <sup>n</sup>	-23245	11	3005.8	1.2			199 ns	3		07	1975	IT=100			
<sup>200</sup> Bi	-20371	22			*		36.4 m	0.5		07	1950	$\beta^+ = 100$			
<sup>200</sup> Bi <sup>m</sup>	-20270#	70#	100#	70#	*		31 m	2		07	1978	$\beta^+ < 100$ ; IT ?			
<sup>200</sup> Bi <sup>n</sup>	-19943	22	428.20	0.10			400 ms	50		07	1972	IT=100			
<sup>200</sup> Po	-16942	8					11.51 m	0.08		07	1951	$\beta^+ = 88.9$ 3; $\alpha = 11.1$ 3			
<sup>200</sup> Po <sup>m</sup>	-14346	8	2596.1	0.3			100 ns	10		07	1985	IT=100			
<sup>200</sup> Po <sup>n</sup>	-14125	11	2817	8			268 ns	3		07	1985	IT=100	*		
<sup>200</sup> At	-8988	24					43.2 s	0.9		07	1963	$\alpha = 52$ 3; $\beta^+ = 48$ 3	*		
<sup>200</sup> At <sup>m</sup>	-8875	25	112.9	2.9	AD		47 s	1		07	1967	$\alpha = 43$ 7; $\beta^+ = ?$ ; IT ?			
<sup>200</sup> At <sup>n</sup>	-8644	25	343.8	3.0	AD		8.0 s	2.1		07	1967	IT < 89.5 3; $\alpha \approx 10.5$ 3; $\beta^+ ?$	*		
<sup>200</sup> Rn	-4005	14					1.09 s	0.16		07	1971	$\alpha = 92$ 8; $\beta^+ ?$	*		
<sup>200</sup> Rn <sup>m</sup>	-1685	24	2320	20			28 $\mu$ s	9		07	2002	IT=100	*		
<sup>200</sup> Fr	6130	30			*		47.5 ms	2.8		07	14Ka23	TD	1995	$\alpha = 100$ ; $\beta^- = 2.5$ %; $\beta^- \text{SF} > 1.4$	*
<sup>200</sup> Fr <sup>m</sup>	6180	50	50	60	AD	*	190 ms	120		07	96En01	TD	1996	$\alpha \approx 100$ ; IT ?	*
<sup>200</sup> Fr <sup>n</sup>	6280#	60#	150#	50#			790 ns	360		07	14Ka23	T	2014	IT ?	*
* <sup>200</sup> Os	T : symmetrized from 14Ku23=14Mo15=6(+4-3); other 05Ku.A=4.6(1.3) same group											**			
* <sup>200</sup> Ir	J : from 13Mo20=(2-,3-)											**			
* <sup>200</sup> Po <sup>n</sup>	E : Ex < 25 keV above 2804.5(0.6) level											**			
* <sup>200</sup> At	T : average 96Ta18=44(2) 92Hu04=43(1)											**			
* <sup>200</sup> At <sup>n</sup>	E : 230.9(0.2) keV above <sup>200</sup> At <sup>m</sup> , from ENSDF											**			
* <sup>200</sup> At <sup>m</sup>	T : symmetrized from 7.3(+2.6-1.5)											**			
* <sup>200</sup> Rn	T : symmetrized from 1.03(+0.20-0.11) D : symmetrized from 86(+14-4)%											**			
* <sup>200</sup> Rn <sup>m</sup>	E : Estimated 20#(20#) keV above 2300.5(0.5) level											**			
* <sup>200</sup> Rn <sup>n</sup>	T : symmetrized from 25(+11-6)											**			
* <sup>200</sup> Fr	T : average 14Ka23=46(4) 05De01=49(4)											**			
* <sup>200</sup> Fr <sup>m</sup>	I : two events with 100 ms and E(a)=7550 correlated with E(a)=6880											**			
* <sup>200</sup> Fr <sup>n</sup>	I : assigned by evaluators to <sup>200</sup> Fr <sup>m</sup> and <sup>196</sup> At <sup>m</sup>											**			
* <sup>200</sup> Fr <sup>m</sup>	I : existence under discussion, level not accepted in ENSDF											**			
* <sup>200</sup> Fr <sup>n</sup>	T : symmetrized from 100(+180-40) (2 evts with half-life=100ms), see 84Sc13											**			
* <sup>200</sup> Fr <sup>n</sup>	E : 14Ka23 > 101.13 keV T : symmetrized from 14Ka23=600(+500-200)											**			
<sup>201</sup> Os	-15240#	300#			1# s	(>300 ns)	1/2 <sup>-</sup> #	13			2009	$\beta^- ?$			
<sup>201</sup> Ir	-19900#	200#			21 s	5	(3/2 <sup>+</sup> )	11	14Mo15	T	2008	$\beta^- = 100$	*		
<sup>201</sup> Pt	-23740	50			2.5 m	0.1	(5/2 <sup>-</sup> )	07			1962	$\beta^- = 100$			
<sup>201</sup> Au	-26401	3			26.0 m	0.8	3/2 <sup>+</sup>	07			1952	$\beta^- = 100$			
<sup>201</sup> Au <sup>m</sup>	-25807	6	594	5	730 $\mu$ s	630	(11/2 <sup>-</sup> )	07	11St21	ETJ	1981	IT=100	*		
<sup>201</sup> Au <sup>n</sup>	-24791	6	1610	5	5.6 $\mu$ s	2.4			11St21	ETD	2011	IT=100	*		
<sup>201</sup> Hg	-27662.5	0.7			STABLE		3/2 <sup>-</sup>	07			1925	IS=13.18 9			
<sup>201</sup> Hg <sup>m</sup>	-26896.3	0.7	766.22	0.15			94.0 $\mu$ s	2.0		07	1961	IT=100			
<sup>201</sup> Tl	-27181	14			3.0442 d	0.0019	1/2 <sup>+</sup>	07	14Un01	T	1950	$\epsilon = 100$	*		
<sup>201</sup> Tl <sup>m</sup>	-26262	14	919.16	0.21			2.01 ms	0.07		07	1962	IT=100			
<sup>201</sup> Pb	-25271	14			9.33 h	0.03	5/2 <sup>-</sup>	07			1950	$\beta^+ = 100$			
<sup>201</sup> Pb <sup>m</sup>	-24642	14	629.1	0.3			60.8 s	1.8		07	1952	IT $\approx$ 100; $\beta^+ ?$			
<sup>201</sup> Pb <sup>n</sup>	-22333	24	2938	20			508 ns	3		07	1981	IT=100	*		
<sup>201</sup> Bi	-21416	15			103 m	3	9/2 <sup>-</sup>	07			1950	$\beta^+ = 100$			
<sup>201</sup> Bi <sup>m</sup>	-20570	15	846.35	0.18			57.5 m	2.1		07	1950	$\beta^+ > 91.1$ %; IT < 8.6; $\alpha = ?$	*		
<sup>201</sup> Bi <sup>n</sup>	-19443	27	1973	23			118 ns	28		07	1982	IT=100	*		
<sup>201</sup> Bi <sup>p</sup>	-19404	27	2012	23			105 ns	75		07	1985	IT=100	*		
<sup>201</sup> Bi <sup>q</sup>	-18635	27	2781	23			124 ns	4		07	1982	IT=100	*		
<sup>201</sup> Po	-16521	5			15.6 m	0.1	3/2 <sup>-</sup>	07	13Se03	J	1951	$\beta^+ = 98.87$ 3; $\alpha = 1.13$ 3			
<sup>201</sup> Po <sup>m</sup>	-16097	5	423.8	2.4	AD		8.96 m	0.12		07	13Se03	J	1962	IT=56.2 12; $\beta^+ = 41.4$ 7; $\alpha = 2.4$ 5	
<sup>201</sup> At	-10789	8			85.2 s	1.6	(9/2 <sup>-</sup> )	07			1963	$\alpha = 71$ 7; $\beta^+ = 29$ 7			
<sup>201</sup> At <sup>m</sup>	-10330	8	459	1			45 ms	3		07	14Au03	ETJ	2015	IT=100	
<sup>201</sup> At <sup>n</sup>	-8470	8	2319	1			3.39 $\mu$ s	0.09		07	15Au01	ETJ	2015	IT=100	*

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...									
<sup>201</sup> Rn	-4070	50		7.0 s	0.4	(3/2 <sup>-</sup> )	07	1967	$\alpha=?; \beta^+=49\#$
<sup>201</sup> Rn <sup>m</sup>	-3790#	90#	280#	3.8 s	0.1	(13/2 <sup>+</sup> )	07	1967	$\beta^+=66\#; \alpha=?$
<sup>201</sup> Fr	3589	9		62.8 ms	1.9	(9/2 <sup>-</sup> )	07	14Ka23 T	$\alpha=100$
<sup>201</sup> Fr <sup>m</sup>	3715	11	127	17 ms	7	(1/2 <sup>+</sup> )	07	14Ka23 T	$\alpha=100$
<sup>201</sup> Fr <sup>n</sup>	3790	60	200	890 ns	360	(13/2 <sup>+</sup> )		14Ka23 ETJ	IT=100
<sup>201</sup> Ra	11937	20		20 ms	30	(3/2 <sup>-</sup> )		14Ka23 TJ	$\alpha=100$
<sup>201</sup> Ra <sup>m</sup>	12200	26	263	6 ms	5	(13/2 <sup>+</sup> )	07	2005	$\alpha=100$
* <sup>201</sup> Ir	J : from 13Mo20=(1/2 <sup>+</sup> , 3/2 <sup>+</sup> , 5/2 <sup>+</sup> ), 3/2 <sup>+</sup> agrees with systematics of odd-A Z=77								
* <sup>201</sup> Au <sup>m</sup>	T : symmetrized from 340(+900-290) $\mu$ s								
* <sup>201</sup> Au <sup>n</sup>	E : 378.2, 638 $\gamma$ s above <sup>201</sup> Au <sup>m</sup>								
* <sup>201</sup> Tl	T : average 14Un01=3.046(0.006) 04Sc04=3.0486(0.0030) 94Si26=3.0400(0.0028)								
* <sup>201</sup> Pb <sup>n</sup>	E : estimated 20#(20#) keV above 2917.6(0.9) level								
* <sup>201</sup> Bi <sup>m</sup>	D : $\alpha$ decay is observed. Its branching ratio is estimated 0.3%# in ENSDF								
* <sup>201</sup> Bi <sup>n</sup>	E : 1933.3(0.4) + x ; x<80								
* <sup>201</sup> Bi <sup>p</sup>	E : 1972.3(0.4) + x ; x<80								
* <sup>201</sup> Bi <sup>q</sup>	E : 2741.0(0.3) + x ; x<80								
* <sup>201</sup> At <sup>n</sup>	E : error estimated by evaluator								
* <sup>201</sup> Rn <sup>m</sup>	T : other 10He25=3.24(+3.24-1.08) ms								
* <sup>201</sup> Fr	T : average 14Ka23=64(3) 05Uu02=53(4) 05De01=67(3)								
* <sup>201</sup> Fr <sup>m</sup>	T : average 14Ka23=8(+12-3) 05Uu02=19(+19-6)								
* <sup>201</sup> Fr <sup>n</sup>	E : derived from range in 14Ka23 101 to 300 keV								
* <sup>201</sup> Fr <sup>n</sup>	T : symmetrized from 14Ka23=700(+500-200)								
* <sup>201</sup> Ra	T : symmetrized from 14Ka23=8(+40-4)								
* <sup>201</sup> Ra <sup>m</sup>	T : symmetrized from 1.6(+7.7-0.7)								
<sup>202</sup> Os	-13090#	400#		200# ms	(>300 ns)	0 <sup>+</sup>	13	2009	$\beta^-?$
<sup>202</sup> Ir	-16780#	300#		11 s	3	(2 <sup>-</sup> )	08	14Ku23 T	$\beta^-=100$
<sup>202</sup> Ir <sup>m</sup>	-14780#	300#	2000#	3.4 $\mu$ s	0.6			11St21 TD	IT=100
<sup>202</sup> Pt	-22692	25		44 h	15	0 <sup>+</sup>	08	1992	$\beta^-=100$
<sup>202</sup> Pt <sup>m</sup>	-20904	25	1788.5	141 $\mu$ s	7	(7 <sup>-</sup> )	08	11St21 T	IT $\approx$ 100
<sup>202</sup> Au	-24353	23		28.4 s	1.2	(1 <sup>-</sup> )	08	1967	$\beta^-=100$
<sup>202</sup> Hg	-27345.3	0.7		STABLE		0 <sup>+</sup>	08	1920	IS=29.86 26
<sup>202</sup> Tl	-25980.2	1.6		12.31 d	0.08	2 <sup>-</sup>	08	1940	$\epsilon=100$
<sup>202</sup> Tl <sup>m</sup>	-25030.0	1.6	950.19	591 $\mu$ s	3	7 <sup>+</sup>	08	1958	IT=100
<sup>202</sup> Pb	-25941	4		52.5 ky	2.8	0 <sup>+</sup>	08	1954	$\epsilon=100$
<sup>202</sup> Pb <sup>m</sup>	-23771	4	2169.85	3.54 h	0.02	9 <sup>-</sup>	08	1954	IT=90.5 5; $\beta^+=9.5$ 5
<sup>202</sup> Pb <sup>n</sup>	-21800	50	4140	110 ns	5	16 <sup>+</sup> #	08	1986	IT=100
<sup>202</sup> Pb <sup>p</sup>	-20640	50	5300	107 ns	3	19 <sup>+</sup> #	08	1987	IT=100
<sup>202</sup> Bi	-20741	15		1.72 h	0.05	5 <sup>(+)</sup> #	08	1951	$\beta^+=100; \alpha<1e-5$
<sup>202</sup> Bi <sup>m</sup>	-20116	19	625	3.04 $\mu$ s	0.06	10 <sup>-</sup> #	08	1981	IT=100
<sup>202</sup> Bi <sup>n</sup>	-18124	19	2617	310 ns	50	(17 <sup>+</sup> )	08	1981	IT=100
<sup>202</sup> Po	-17942	9		44.6 m	0.4	0 <sup>+</sup>	08	1951	$\beta^+=?; \alpha=1.92$ 7
<sup>202</sup> Po <sup>m</sup>	-16230	15	1712	110 ns	15	8 <sup>+</sup>	08	1971	IT=100
<sup>202</sup> At	-10591	28		184 s	1	3 <sup>(+)</sup>	08	16Ly01 JD	$\beta^+=?; \alpha=12$ 7
<sup>202</sup> At <sup>m</sup>	-10401	28	190	182 s	2	7 <sup>(+)</sup>	08	16Ly01 J	IT ?; $\beta^+ ?; \alpha=8.7$ 15
<sup>202</sup> At <sup>n</sup>	-10010	28	580	460 ms	50	10 <sup>(-)</sup>	08	16Ly01 J	IT $\approx$ 100; $\alpha=0.096$ 11; $\beta^+=0.033\#$
<sup>202</sup> Rn	-6275	18		9.7 s	0.1	0 <sup>+</sup>	08	1967	$\alpha=78$ 8; $\beta^+ ?$
<sup>202</sup> Rn <sup>m</sup>	-3970#	50#	2310#	2.22 $\mu$ s	0.07	11 <sup>-</sup> #		02Do19 T	IT=100
<sup>202</sup> Fr	3096	7		372 ms	12	3 <sup>+</sup>	08	14Ka23 T	$\alpha=?; \beta^+=14\#$
<sup>202</sup> Fr <sup>m</sup>	3370	9	274	286 ms	13	10 <sup>-</sup>	08	14Ka23 T	$\alpha=?; \beta^+=14\#$
<sup>202</sup> Ra	9075	15		4.1 ms	1.1	0 <sup>+</sup>	08	14Ka23 T	$\alpha=100$
* <sup>202</sup> Ir	T : 14Ku23=11(3) supersedes 14Mo15=15(3) J : from 13Mo20=(2 <sup>-</sup> )								
* <sup>202</sup> Ir <sup>m</sup>	D : 311.5, 655.9, 737.2, 889.2, 967.6 $\gamma$ rays seen in decay								
* <sup>202</sup> Hg	D : lower half-life limit for <sup>24</sup> Ne decay $T>3.7$ Zy, from 90Bu28								
* <sup>202</sup> Pb <sup>n</sup>	E : 4091.0(0.7) + x ; x estimated 50(50)								
* <sup>202</sup> Pb <sup>p</sup>	E : 5251.0(0.5) + u ; u estimated 50(50)								
* <sup>202</sup> Bi	J : re-evaluation to a possible 6 <sup>+</sup> is discussed in 96Ca02								
* <sup>202</sup> Bi <sup>m</sup>	E : 605 + x with x<40 keV								
* <sup>202</sup> Bi <sup>n</sup>	E : 2597.07(0.25) + x, with x<40 keV								
* <sup>202</sup> Po <sup>m</sup>	E : 1691.5(0.4) + x, with x<40 keV								
* <sup>202</sup> At <sup>n</sup>	E : 391.7(0.5) keV above <sup>202</sup> At <sup>m</sup>								
* <sup>202</sup> Fr	J : from 13Fl09=3 <sup>+</sup> (see their Fig.2)								
* <sup>202</sup> Fr <sup>m</sup>	J : from 13Fl09=10 <sup>-</sup> (see their Fig.2)								
* <sup>202</sup> Ra	T : symmetrized from 14Ka23=3.8(+1.3-0.8)								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^{\pi}$	Ens	Reference	Year of discovery	Decay modes and intensities (%)					
<sup>203</sup> Os	-7640#	400#	100# ms (>300 ns)	9/2 <sup>+</sup> #	13	12Ku26	I	2012	$\beta^-$ ?; $\beta^- n=7\#$				
<sup>203</sup> Ir	-14690#	400#	6# s (>300 ns)	3/2 <sup>+</sup> #	13	09St16	I	2009	$\beta^-$ ?				
<sup>203</sup> Ir <sup>m</sup>	-12550#	400#	798 ns	350 (23/2 <sup>+</sup> )		11St21	TJD	2011	IT=100				
<sup>203</sup> Pt	-19630#	200#	22 s	4 (1/2 <sup>-</sup> )	06	14Mo15	T	2008	$\beta^- =100$				
<sup>203</sup> Pt <sup>m</sup>	-16530#	200#	3100#	1000#	641 ns	55	33/2 <sup>+</sup> #	11St21	TJD	2011	IT=100		
<sup>203</sup> Au	-23143	3	60 s	6	3/2 <sup>+</sup>	05		1952	$\beta^- =100$				
<sup>203</sup> Au <sup>m</sup>	-22502	4	641	3	140 $\mu$ s	44	11/2 <sup>-</sup> #	05	11St21	TJ	2005	IT=100	
<sup>203</sup> Hg	-25269.3	1.6	46.613 d	0.018	5/2 <sup>-</sup>	05	14Un01	T	1943	$\beta^- =100$			
<sup>203</sup> Hg <sup>m</sup>	-24336.2	1.6	933.14	0.23	21.9 $\mu$ s	1.0	(13/2 <sup>+</sup> )	05	11St21	T	1964	IT=100	
<sup>203</sup> Hg <sup>n</sup>	-16988.3	1.7	8281.0	0.5	146 ns	30	(53/2 <sup>+</sup> )		11Sz01	EJT	2011	IT=100	
<sup>203</sup> Tl	-25761.4	1.2	STABLE				1/2 <sup>+</sup>	05			1931	IS=29.52 1	
<sup>203</sup> Tl <sup>m</sup>	-22200	50	3565	50	7.7 $\mu$ s	0.5	(25/2 <sup>+</sup> )	05			1998	IT=100	
<sup>203</sup> Pb	-24787	7	51.916 h	0.015	5/2 <sup>-</sup>	05	14Un01	T	1942	$\epsilon =100$			
<sup>203</sup> Pb <sup>m</sup>	-23962	7	825.2	0.3	6.21 s	0.11	13/2 <sup>+</sup>	05			1955	IT=100	
<sup>203</sup> Pb <sup>n</sup>	-21838	7	2949.2	0.4	480 ms	7	29/2 <sup>-</sup>	05			1977	IT=100	
<sup>203</sup> Pb <sup>p</sup>	-21820	50	2970	50	122 ns	4	25/2 <sup>-</sup> #	05			1988	IT=100	
<sup>203</sup> Bi	-21525	13	11.76 h	0.05	11.76 h	0.05	9/2 <sup>-</sup>	05			1950	$\beta^+ =100$	
<sup>203</sup> Bi <sup>m</sup>	-20427	13	1098.12	0.12	305 ms	5	1/2 <sup>+</sup>	05			1984	IT=100	
<sup>203</sup> Bi <sup>n</sup>	-19484	13	2041.5	0.6	194 ns	30	25/2 <sup>+</sup>	05			1978	IT=100	
<sup>203</sup> Po	-17311	9	36.7 m	0.5	36.7 m	0.5	5/2 <sup>-</sup>	05	13Se03	J	1951	$\beta^+ \approx 100$ ; $\alpha=0.11 2$	
<sup>203</sup> Po <sup>m</sup>	-16669	9	641.68	0.17	45 s	2	13/2 <sup>+</sup>	05	13Se03	J	1969	IT $\approx$ 100; $\alpha=0.04\#$	
<sup>203</sup> Po <sup>n</sup>	-15153	9	2158.5	0.6	> 200 ns			05			1986	IT=100	
<sup>203</sup> At	-12163	11	7.4 m	0.2	7.4 m	0.2	9/2 <sup>-</sup>	05			1951	$\beta^+ =69 3$ ; $\alpha=31 3$	
<sup>203</sup> Rn	-6154	18	44 s	2	44 s	2	3/2 <sup>-</sup> #	05			1967	$\alpha=66 9$ ; $\beta^+ ?$	
<sup>203</sup> Rn <sup>m</sup>	-5791	18	363	4	AD	26.9 s	0.5	13/2 <sup>(+)</sup>	05	87Bo29	J	1967	$\alpha=75 10$ ; $\beta^+ ?$
<sup>203</sup> Fr	876	6	550 ms	10	9/2 <sup>-</sup>	05	13Fi09	J	1967	$\alpha \approx 100$ ; $\beta^+ =5\#$			
<sup>203</sup> Fr <sup>m</sup>	1237	7	361	6	43 ms	4	(1/2 <sup>+</sup> )		13Ja06	TJD	2013	IT=?; $\alpha=20 4$	
<sup>203</sup> Fr <sup>n</sup>	1300	100	426	100	370 ns	50	(13/2 <sup>+</sup> )		13Ja06	TJD	2013	IT $\approx$ 100	
<sup>203</sup> Ra	8660	40	36 ms	13	(3/2 <sup>-</sup> )	05	96Le09	J	1996	$\alpha \approx 100$ ; $\beta^+ ?$			
<sup>203</sup> Ra <sup>m</sup>	8851	29	190	50	AD	25 ms	5	(13/2 <sup>+</sup> )	05	96Le09	J	1996	$\alpha \approx 100$ ; $\beta^+ ?$
* <sup>203</sup> Ir <sup>m</sup>	E : 207.0, 841.3, 894.7 $\gamma$ s in cascade to 11/2 <sup>-</sup> estimated at 200(50) keV								**				
* <sup>203</sup> Pt	J : from 13Mo20=(1/2 <sup>-</sup> )								**				
* <sup>203</sup> Hg	T : average 14Un01=46.62(0.06) 83Wa26=46.612(0.019)								**				
* <sup>203</sup> Tl <sup>m</sup>	E : 3514.6 + x and x estimated 50(50) keV								**				
* <sup>203</sup> Pb	T : average 14Un01=51.923(0.036) 01Li17=51.99(0.03) 80Ho17=51.88(0.02)								**				
* <sup>203</sup> Pb <sup>p</sup>	E : 2923.4(0.7) + x ; x estimated 50(50)								**				
* <sup>203</sup> Rn	J : not yet known, will be same as <sup>195</sup> Pb and <sup>199</sup> Po, from $\alpha$ decay								**				
* <sup>203</sup> Ra	T : symmetrized from 05Uu02=31(+17-9); other 14Ka23=50(+40-15)								**				
* <sup>203</sup> Ra <sup>m</sup>	T : symmetrized from 05Uu02=24(+6-4); other 14Ka23=37(+37-12)								**				
<sup>204</sup> Ir	-9690#	400#	1# s (>300 ns)		13	12Ku26	I		$\beta^- ?$ ; $\beta^- n=0.01\#$				
<sup>204</sup> Pt	-17920#	200#	10.3 s	1.4	0 <sup>+</sup>	10		2008	$\beta^- =100$				
<sup>204</sup> Pt <sup>m</sup>	-15930#	200#	5.5 $\mu$ s	0.7	(5 <sup>-</sup> )	10	11St21	E	2009	IT=100			
<sup>204</sup> Pt <sup>n</sup>	-15890#	200#	2035	23	55 $\mu$ s	3	(7 <sup>-</sup> )	10		2009	IT ?		
<sup>204</sup> Pt <sup>p</sup>	-14730#	200#	3193	23	146 ns	14	(10 <sup>+</sup> )	10		2009	IT=100		
<sup>204</sup> Au	-20650#	200#	38.3 s	1.3	(2 <sup>-</sup> )	10	14Mo15	T	1972	$\beta^- =100$			
<sup>204</sup> Au <sup>m</sup>	-16830#	200#	3816#	1000#	2.1 $\mu$ s	0.3	16 <sup>+</sup> #	10	11St21	JD	2008	IT=100	
<sup>204</sup> Hg	-24690.1	0.5	STABLE				0 <sup>+</sup>	10			1920	IS=6.87 15; 2 $\beta^- ?$	
<sup>204</sup> Hg <sup>m</sup>	-20301.4	0.7	4388.7	0.5	29 ns	2.1	4 <sup>+</sup>		15Wr02	ETJ	2015	IT=100	
<sup>204</sup> Hg <sup>n</sup>	-17464.0	0.7	7226.1	0.5	> 480 ns	2	2 <sup>+</sup>		15Wr02	ETJ	2015	IT=100	
<sup>204</sup> Tl	-24346.1	1.2	3.783 y	0.012	3.783 y	0.012	2 <sup>-</sup>	10			1953	$\beta^- =97.08 7$ ; $\epsilon + \beta^+ =2.92 7$	
<sup>204</sup> Tl <sup>m</sup>	-23242.0	1.2	1104.1	0.2	61.7 $\mu$ s	1.0	7 <sup>+</sup>	10	11Br12	EJ	1972	IT=100	
<sup>204</sup> Tl <sup>n</sup>	-22027.1	1.2	2319.0	0.3	2.6 $\mu$ s	0.2	12 <sup>-</sup>	10	11Br12	EJ	1998	IT=100	
<sup>204</sup> Tl <sup>p</sup>	-19954.5	1.3	4391.6	0.5	420 ns	30	18 <sup>+</sup>	10	11Br12	ETJ	1998	IT=100	
<sup>204</sup> Tl <sup>q</sup>	-18106.7	1.3	6239.4	0.5	90 ns	3	22 <sup>-</sup>	10	11Br12	ETJ	2011	IT=100	
<sup>204</sup> Pb	-25109.9	1.1	STABLE				(>140 Py)	0 <sup>+</sup>	10		1932	IS=1.4 1; $\alpha ?$	
<sup>204</sup> Pb <sup>m</sup>	-23835.8	1.1	1274.13	0.05	265 ns	6	4 <sup>+</sup>	10			1963	IT=100	
<sup>204</sup> Pb <sup>n</sup>	-22924.0	1.1	2185.88	0.08	66.93 m	0.10	9 <sup>-</sup>	10			1956	IT=100	
<sup>204</sup> Pb <sup>p</sup>	-22845.5	1.1	2264.42	0.06	490 ns	70	7 <sup>-</sup>	10			1978	IT=100	
<sup>204</sup> Bi	-20646	9	11.22 h	0.10	11.22 h	0.10	6 <sup>+</sup>	10			1947	$\beta^+ =100$	
<sup>204</sup> Bi <sup>m</sup>	-19841	9	805.5	0.3	13.0 ms	0.1	10 <sup>-</sup>	10			1974	IT=100	
<sup>204</sup> Bi <sup>n</sup>	-17813	9	2833.4	1.1	1.07 ms	0.03	17 <sup>+</sup>	10			1974	IT=100	
<sup>204</sup> Po	-18341	11	3.519 h	0.012	3.519 h	0.012	0 <sup>+</sup>	10			1951	$\beta^+ =99.33 3$ ; $\alpha=0.67 3$	
<sup>204</sup> Po <sup>m</sup>	-16702	11	1639.03	0.06	158.6 ns	1.8	8 <sup>+</sup>	10	10Ka29	T	1970	IT=100	
<sup>204</sup> At	-11875	22	9.12 m	0.11	9.12 m	0.11	7 <sup>+</sup>	10			1961	$\beta^+ =96.2 2$ ; $\alpha=3.8 2$	
<sup>204</sup> At <sup>m</sup>	-11288	22	587.30	0.20	108 ms	10	10 <sup>-</sup>	10			1969	IT=100	
<sup>204</sup> Rn	-7970	7	1.242 m	0.023	1.242 m	0.023	0 <sup>+</sup>	10			1967	$\alpha=72.4 9$ ; $\beta^+ ?$	

... A-group is continued on next page ...



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>204</sup> Fr	607 25		1.75 s 0.26	3 <sup>+</sup>	10	95Bi.A D	1964	$\alpha=96\ 2; \beta^+ ?$ *
<sup>204</sup> Fr <sup>m</sup>	658 25 50	4 AD	2.41 s 0.19	7 <sup>(+)</sup>	10	95Bi.A D	1967	$\alpha=90\ 2; \beta^+ ?$ *
<sup>204</sup> Fr <sup>n</sup>	934 25 326	4 AD	1.65 s 0.15	10 <sup>(-)</sup>	10	13Ja06 T	1992	$\alpha=53\ 10; IT=47\ 10$ *
<sup>204</sup> Ra	6057 15		60 ms 9	0 <sup>+</sup>	10	05Uu02 T	1995	$\alpha\approx 100; \beta^+=0.3\#$ *
* <sup>204</sup> Pt	T: other 14Mo15=16(+6-5) **							
* <sup>204</sup> Pt <sup>m</sup>	E: 872.4(0.5), 1122.7(0.5) $\gamma$ s to 0 <sup>+</sup> **							
* <sup>204</sup> Pt <sup>n</sup>	E: 1995.1(0.7) + x; x < 80 keV **							
* <sup>204</sup> Pt <sup>p</sup>	E: 1157.5(0.5) $\gamma$ to <sup>204</sup> Pt <sup>n</sup> **							
* <sup>204</sup> Au	T: average 14Mo15=37.2(0.8) 84Cr01=39.8(0.9); other 72Pa06=40(3) **							
* <sup>204</sup> Au <sup>m</sup>	E: 839.0, 976.6 $\gamma$ s in cascade to 12 <sup>-</sup> # estimated at 2000#(1000#) keV **							
* <sup>204</sup> Pb	T: also 13Be16>140Ey **							
* <sup>204</sup> Pb <sup>p</sup>	T: symmetrized from 450(+100-30) **							
* <sup>204</sup> Po <sup>m</sup>	T: average 10Ka29=161(4) 87Ra04=158(2); others 90Fa03=150(10) 83He08=150(10) **							
* <sup>204</sup> Po <sup>m</sup>	T: 71Ha01=140(5) 70Ya03=190(20) 70Br.A=143(5) **							
* <sup>204</sup> At	T: other 10Ka29=9.6(2) **							
* <sup>204</sup> Fr	T: average 05Uu02=1.9(0.5) 92Hu04=1.7(0.3) J: 14Ly01=3 13Vo10=3 **							
* <sup>204</sup> Fr <sup>m</sup>	T: average 13Ja06=2.6(0.3) 05Uu02=1.6(+0.5-0.3) 92Hu04=2.6(0.3) **							
* <sup>204</sup> Fr <sup>m</sup>	J: 15Vo05=7 **							
* <sup>204</sup> Fr <sup>n</sup>	E: 276.1 keV above <sup>204</sup> Fr <sup>m</sup> , from 95Bi.A D: from 14Ly01 **							
* <sup>204</sup> Fr <sup>n</sup>	T: 13Ja06=1.65(0.15) supersedes 05Uu02=0.8(0.2) same group J: 15Vo05=10 **							
* <sup>204</sup> Ra	T: average 05Uu02=54(+19-11) 96Le09=59(+12-9); other 10He25=44(+44-15) **							
* <sup>204</sup> Ra	T: 95Le04=45(+55-21) **							
<sup>205</sup> Ir	-5960# 500#		300# ms (>300 ns)	3/2 <sup>+</sup> #	13	12Ku26 I	2012	$\beta^- ?; \beta^-n=10\#$
<sup>205</sup> Pt	-12970# 300#		5# s (>300 ns)	9/2 <sup>+</sup> #	11	10Al24 I	2009	$\beta^- ?$
<sup>205</sup> Au	-18770# 200#		32.5 s 1.4	3/2 <sup>+</sup> #	04	09Po01 T	1994	$\beta^- =100$ *
<sup>205</sup> Au <sup>m</sup>	-17860# 200# 907	5	6 s 2	11/2 <sup>-</sup> #		09Po01 ETJ	2009	IT=?; $\beta^- =?$
<sup>205</sup> Au <sup>n</sup>	-15920# 200# 2850	5	163 ns 5	19/2 <sup>+</sup> #		11St21 ET	2011	IT=100
<sup>205</sup> Hg	-22288 4		5.14 m 0.09	1/2 <sup>-</sup>	04		1940	$\beta^- =100$ *
<sup>205</sup> Hg <sup>m</sup>	-20732 4 1556.40	0.17	1.09 ms 0.04	13/2 <sup>+</sup>	04		1985	IT=100
<sup>205</sup> Hg <sup>n</sup>	-18972 4 3315.8	0.9	5.89 $\mu$ s 0.18	(23/2 <sup>-</sup> )		11St21 ETJ	2011	IT=100 *
<sup>205</sup> Tl	-23820.9 1.2		STABLE	1/2 <sup>+</sup>	04		1931	IS=70.48 1
<sup>205</sup> Tl <sup>m</sup>	-20530.3 1.2 3290.60	0.17	2.6 $\mu$ s 0.2	25/2 <sup>+</sup>	04		1976	IT=100
<sup>205</sup> Tl <sup>n</sup>	-18985.3 1.9 4835.6	1.5	235 ns 10	(35/2 <sup>-</sup> )	04		2004	IT=100
<sup>205</sup> Pb	-23770.2 1.1		17.3 My 0.7	5/2 <sup>-</sup>	04		1954	$\epsilon=100$
<sup>205</sup> Pb <sup>m</sup>	-23767.9 1.1 2.329	0.007	24.2 $\mu$ s 0.4	1/2 <sup>-</sup>	04		1994	IT=100
<sup>205</sup> Pb <sup>n</sup>	-22756.4 1.1 1013.85	0.03	5.55 ms 0.02	13/2 <sup>+</sup>	04		1960	IT=100
<sup>205</sup> Pb <sup>p</sup>	-20574.5 1.2 3195.7	0.5	217 ns 5	25/2 <sup>-</sup>	04		1973	IT=100
<sup>205</sup> Bi	-21065 5		15.31 d 0.04	9/2 <sup>-</sup>	04		1951	$\beta^+ =100$
<sup>205</sup> Bi <sup>m</sup>	-19568 5 1497.17	0.09	7.9 $\mu$ s 0.7	1/2 <sup>+</sup>	04		1972	IT=100
<sup>205</sup> Bi <sup>n</sup>	-18926 5 2139.0	0.7	220 ns 25	25/2 <sup>+</sup>	04		1978	IT=100
<sup>205</sup> Po	-17521 10		1.74 h 0.08	5/2 <sup>-</sup>	04		1951	$\beta^+ \approx 100; \alpha=0.04\ 1$
<sup>205</sup> Po <sup>m</sup>	-17378 10 143.166	0.017	310 ns 60	1/2 <sup>-</sup>	04		1960	IT=100
<sup>205</sup> Po <sup>n</sup>	-16641 10 880.31	0.07	645 $\mu$ s 20	13/2 <sup>+</sup>	04		1962	IT=100
<sup>205</sup> Po <sup>p</sup>	-16060 10 1461.21	0.21	57.4 ms 0.9	19/2 <sup>-</sup>	04		1973	IT=100
<sup>205</sup> Po <sup>q</sup>	-14434 10 3087.2	0.4	115 ns 10	29/2 <sup>-</sup>	04		1985	IT=100
<sup>205</sup> At	-12972 15		33.8 m 0.2	9/2 <sup>-</sup>	04	10Ka29 T	1951	$\beta^+ ?; \alpha=10\ 2$
<sup>205</sup> At <sup>m</sup>	-10632 15 2339.65	0.23	7.76 $\mu$ s 0.14	29/2 <sup>+</sup>	04		1982	IT=100
<sup>205</sup> Rn	-7710 5		2.83 m 0.07	5/2 <sup>-</sup>	04		1967	$\beta^+ ?; \alpha=24.6\ 9$
<sup>205</sup> Rn <sup>m</sup>	-7053 5 657.1	0.5	> 10 s	13/2 <sup>+</sup> #	04	10De04 ED	2010	IT $\approx$ 100; $\alpha ?; \beta^+ ?$
<sup>205</sup> Fr	-1310 8		3.82 s 0.06	9/2 <sup>-</sup>	04	10De04 T	1964	$\alpha\approx 100; \beta^+ < 1$ *
<sup>205</sup> Fr <sup>m</sup>	-766 8 544.0	1.0	80 ns 20	(13/2 <sup>+</sup> )		12Ja01 EJT	2012	IT=100
<sup>205</sup> Fr <sup>n</sup>	-701 9 609	5	1.15 ms 0.04	(1/2 <sup>+</sup> )		12Ja01 ETJ	2012	IT=100
<sup>205</sup> Ra	5840 70		220 ms 50	(3/2 <sup>-</sup> )	04		1987	$\alpha=?; \beta^+ ?$ *
<sup>205</sup> Ra <sup>m</sup>	6140# 100# 300# 100#		180 ms 50	(13/2 <sup>+</sup> )	04		1995	$\alpha=?; IT ?; \beta^+ ?$ *
<sup>205</sup> Ac	14110 50		80 ms 60	9/2 <sup>-</sup> #	14	14Zh03 T	2014	$\alpha\approx 100; \beta^+=0.2\#$ *
* <sup>205</sup> Au	T: average 09Po01=34(2) 94We02=31(2); other 16Ca25=35(17) **							
* <sup>205</sup> Hg	T: other 10Ku02=5.61(0.38) for q=80 <sup>+</sup> (bare ion) **							
* <sup>205</sup> Hg <sup>n</sup>	E: least-squares fit to $\gamma$ -ray energies 227.6(0.5), 722.6(0.5), 810.0(0.5) 1014.7(0.5) **							
* <sup>205</sup> Fr	T: unweighed average 10De04=4.03(0.08) 05De01=3.80(0.03) 81Ri04=3.96(0.04) **							
* <sup>205</sup> Fr	T: 74Ho27=3.7(0.1) 67Va20=3.7(0.2) 64Gr04=3.7(0.4) **							
* <sup>205</sup> Fr	J: from 14Ly01=9/2 13Vo10=9/2 13Fl09=9/2; parity from mag. moment **							
* <sup>205</sup> Ra	T: symmetrized from 210(+60-40) **							
* <sup>205</sup> Ra <sup>m</sup>	T: symmetrized from 170(+60-40); other 10He25=68(+68-23) ms **							
* <sup>205</sup> Ac	T: symmetrized from 14Zh03=20(+97-9) **							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>206</sup> Pt	-9630# 300#		5# s (>300 ns)	0 <sup>+</sup>	13	12Ku26	I	2012	$\beta^-$ ?; $\beta^-n=0\#$
<sup>206</sup> Au	-14220# 300#		47 s 11	(5 <sup>+</sup> , 6 <sup>+</sup> )	16	16Ca25	TJ	2009	$\beta^-$ =100 *
<sup>206</sup> Hg	-20946 20		8.32 m 0.07	0 <sup>+</sup>	08			1961	$\beta^-$ =100
<sup>206</sup> Hg <sup>m</sup>	-18844 20 2102.4 0.3		2.09 $\mu$ s 0.02	5 <sup>-</sup>	08	11St21	T	1982	IT=100 *
<sup>206</sup> Hg <sup>n</sup>	-17224 20 3722.3 1.0		106 ns 6	(10 <sup>+</sup> )	08	11St21	ETJ	2001	IT=100 *
<sup>206</sup> Tl	-22253.4 1.3		4.202 m 0.011	0 <sup>-</sup>	08			1935	$\beta^-$ =100
<sup>206</sup> Tl <sup>m</sup>	-19610.3 1.3 2643.10 0.18		3.74 m 0.03	(12 <sup>-</sup> )	08			1976	IT=100
<sup>206</sup> Pb	-23785.6 1.1		STABLE (>2.5 Zy)	0 <sup>+</sup>	08	13Be16	T	1927	IS=24.1 1; $\alpha$ ?
<sup>206</sup> Pb <sup>m</sup>	-21585.4 1.1 2200.16 0.04		125 $\mu$ s 2	7 <sup>-</sup>	08			1953	IT=100
<sup>206</sup> Pb <sup>n</sup>	-19758.3 1.3 4027.3 0.7		202 ns 3	12 <sup>+</sup>	08			1971	IT=100
<sup>206</sup> Bi	-20028 8		6.243 d 0.003	6 <sup>(+)</sup>	08			1947	$\beta^+$ =100
<sup>206</sup> Bi <sup>m</sup>	-19968 8 59.897 0.017		7.7 $\mu$ s 0.2	(4 <sup>+</sup> )	08			1957	IT=100
<sup>206</sup> Bi <sup>n</sup>	-18983 8 1044.8 0.7		890 $\mu$ s 10	(10 <sup>-</sup> )	08			1974	IT=100
<sup>206</sup> Po	-18189 4		8.8 d 0.1	0 <sup>+</sup>	08			1947	$\beta^+$ =94.55 5; $\alpha$ =5.45 5
<sup>206</sup> Po <sup>m</sup>	-16603 4 1585.90 0.11		232 ns 4	8 <sup>+</sup>	08	FGK145	J	1970	IT=100 *
<sup>206</sup> Po <sup>n</sup>	-15927 4 2262.09 0.12		1.05 $\mu$ s 0.06	9 <sup>-</sup>	08	FGK145	J	1970	IT=100 *
<sup>206</sup> At	-12430 15		30.6 m 0.8	(5 <sup>+</sup> )	08			1961	$\beta^+$ =99.10 8; $\alpha$ =0.90 8
<sup>206</sup> At <sup>m</sup>	-11620 15 810 3		813 ns 21	(10 <sup>-</sup> )	08	09Dr08	T	1999	IT=100 *
<sup>206</sup> Rn	-9133 9		5.67 m 0.17	0 <sup>+</sup>	08			1954	$\alpha$ =62 3; $\beta^+$ =38 3
<sup>206</sup> Fr	-1242 28		16 s	3 <sup>+</sup>	08	16Ly01	D	1964	$\beta^+$ ?; $\alpha$ =88.4 33
<sup>206</sup> Fr <sup>m</sup>	-1048 28 190 40		16 s	7 <sup>(+)</sup>	08	16Ly01	D	1964	$\alpha$ =84.7 15; $\beta^+$ ?; IT?
<sup>206</sup> Fr <sup>n</sup>	-517 28 730 40 MD		700 ms 100	10 <sup>(-)</sup>	08	16Ly01	D	1983	IT=?; $\alpha$ =13 2 *
<sup>206</sup> Fr <sup>x</sup>	-1140 100 100 100 MD		R=?	spmix					
<sup>206</sup> Ra	3566 18		240 ms 20	0 <sup>+</sup>	08			1967	$\alpha$ =?; $\beta^+$ =2.5#
<sup>206</sup> Ac	13480 50		25 ms 7	(3 <sup>+</sup> )	08			1998	$\alpha$ ≈100; $\beta^+$ =0.2# *
<sup>206</sup> Ac <sup>m</sup>	13700 30 220 60 AD		41 ms 16	(10 <sup>-</sup> )	08			1996	$\alpha$ ≈100; $\beta^+$ ? *
* <sup>206</sup> Au	T: average 16Ca25=56(17) 15Mo20=40(15)								**
* <sup>206</sup> Hg <sup>m</sup>	T: average 11St21(=09Si35)=2.09(0.02) 82Be38=2.15(0.21)								**
* <sup>206</sup> Hg <sup>n</sup>	T: average 11St21(=09Si35)=112(4) 09Al29=96(15) 01Fo08=92(8) 01La09=90(10)								**
* <sup>206</sup> Po <sup>m</sup>	J: measured magnetic moment and non-observation of $\gamma$ s to 3 <sup>+</sup> and 4 <sup>+</sup> levels								**
* <sup>206</sup> Po <sup>n</sup>	J: E1 $\gamma$ s to 8 <sup>+</sup> levels								**
* <sup>206</sup> At <sup>m</sup>	T: others 10Ka29=377(44) 99Fe10=410(80)								**
* <sup>206</sup> At <sup>n</sup>	E: from ENSDF'08 806.7(1.4) + x; x<6 estimated by NUBASE								**
* <sup>206</sup> Fr	J: 14Ly01=3 13Vo10=3								**
* <sup>206</sup> Fr <sup>m</sup>	T: 92Hu04=15.9(0.3) J: 15Vo05=7								**
* <sup>206</sup> Fr <sup>n</sup>	E: 81Ri04=531(2) keV above <sup>206</sup> Fr <sup>m</sup> J: 15Vo05=10								**
* <sup>206</sup> Ac	T: symmetrized from 98Es02=22(+9-5); also 14Zh03=41(+56-15)								**
* <sup>206</sup> Ac <sup>m</sup>	T: symmetrized from 98Es02=33(+22-9)								**
<sup>207</sup> Pt	-4540# 400#		1# s (>300 ns)	9/2 <sup>+</sup> #	13	12Ku26	I	2012	$\beta^-$ ?; $\beta^-n=2\#$
<sup>207</sup> Au	-10810# 300#		10# s (>300 ns)	3/2 <sup>+</sup> #	11			2010	$\beta^-$ ?; $\beta^-n=0.4\#$
<sup>207</sup> Hg	-16487 30		2.9 m 0.2	9/2 <sup>+</sup> #	11			1982	$\beta^-$ =100
<sup>207</sup> Tl	-21034 5		4.77 m 0.02	1/2 <sup>+</sup>	11			1908	$\beta^-$ =100 *
<sup>207</sup> Tl <sup>m</sup>	-19686 5 1348.18 0.16		1.33 s 0.11	11/2 <sup>-</sup>	11			1965	IT≈100; $\beta^- < 0.1\#$
<sup>207</sup> Pb	-22452.0 1.1		STABLE (>1.9 Zy)	1/2 <sup>-</sup>	11	13Be16	T	1927	IS=22.1 1; $\alpha$ ?
<sup>207</sup> Pb <sup>m</sup>	-20818.6 1.1 1633.356 0.004		806 ms 5	13/2 <sup>+</sup>	11			1951	IT=100
<sup>207</sup> Bi	-20054.6 2.4		31.20 y 0.03	9/2 <sup>-</sup>	11	14Un01	T	1950	$\beta^+$ =100
<sup>207</sup> Bi <sup>m</sup>	-17953.0 2.4 2101.61 0.16		182 $\mu$ s 6	21/2 <sup>+</sup>	11			1967	IT=100
<sup>207</sup> Po	-17146 7		5.80 h 0.02	5/2 <sup>-</sup>	11			1947	$\beta^+$ ≈100; $\alpha$ =0.021 2
<sup>207</sup> Po <sup>m</sup>	-17077 7 68.557 0.014		205 ns 10	1/2 <sup>-</sup>	11			1963	IT=100
<sup>207</sup> Po <sup>n</sup>	-16031 7 1115.076 0.017		49 $\mu$ s 4	13/2 <sup>+</sup>	11			1962	IT=100
<sup>207</sup> Po <sup>n</sup>	-15763 7 1383.16 0.07		2.79 s 0.08	19/2 <sup>-</sup>	11			1961	IT=100
<sup>207</sup> At	-13227 12		1.81 h 0.03	9/2 <sup>-</sup>	11			1951	$\beta^+$ ?; $\alpha$ ≈10
<sup>207</sup> At <sup>m</sup>	-11110 12 2117.3 0.6		108 ns 2	25/2 <sup>+</sup>	11			1981	IT=100
<sup>207</sup> Rn	-8635 8		9.25 m 0.17	5/2 <sup>-</sup>	11			1954	$\beta^+$ =79 3; $\alpha$ =21 3
<sup>207</sup> Rn <sup>m</sup>	-7736 8 899.1 1.0		184.5 $\mu$ s 0.9	13/2 <sup>+</sup>	11			1974	IT=100
<sup>207</sup> Fr	-2844 18		14.8 s 0.1	9/2 <sup>-</sup>	11	85Co24	J	1964	$\alpha$ =95 2; $\beta^+$ ?
<sup>207</sup> Ra	3540 50		1.38 s 0.18	5/2 <sup>-</sup> #	11			1967	$\alpha$ ≈86; $\beta^+$ ?
<sup>207</sup> Ra <sup>m</sup>	4102 20 560 50 AD		57 ms 8	13/2 <sup>+</sup> #	11	96Le09	T	1987	IT=85#; $\alpha$ =?; $\beta^+$ =0.55# *
<sup>207</sup> Ac	11150 50		31 ms 8	9/2 <sup>-</sup> #	11	98Es02	T	1994	$\alpha$ ≈100 *
* <sup>207</sup> Tl	T: other 05Oh08=4.25(0.14) 10Ku02=4.70(0.19) for q=81 <sup>+</sup> (bare ion)								**
* <sup>207</sup> Ra	T: average 95Uu01=1.1(+0.9-0.3) 68Lo15=1.8(0.5) 67Va22=1.3(0.2)								**
* <sup>207</sup> Ra <sup>m</sup>	T: average 96Le09=63(16) 87He10=55(10)								**
* <sup>207</sup> Ac	T: average 98Es02=27(+11-6) 94Le05=22(+40-9)								**
* <sup>207</sup> Ac	J: unhindered $\alpha$ decay to <sup>203</sup> Fr 9/2 <sup>-</sup> #								**

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>208</sup> Pt	-990# 400#		1# s (>300 ns)	0 <sup>+</sup>	13	12Ku26 I	2012	$\beta^- ?$ ; $\beta^- n=90\#$
<sup>208</sup> Au	-6100# 300#		10# s (>300 ns)		11	10A124 I	2010	$\beta^- ?$ ; $\beta^- n=5\#$
<sup>208</sup> Hg	-13270 30		42 m 5	0 <sup>+</sup>	10		1994	$\beta^- =100$
<sup>208</sup> Hg <sup>m</sup>	-11930 40	1338 24	99 ns 14	(8 <sup>+</sup> )	10		2009	IT=100
<sup>208</sup> Tl	-16750.1 1.9		3.053 m 0.004	5 <sup>+</sup>	07		1909	$\beta^- =100$
<sup>208</sup> Pb	-21748.6 1.1		STABLE (>2.6 Zy)	0 <sup>+</sup>	07	13Be16 T	1927	IS=52.4 1; $\alpha ?$
<sup>208</sup> Pb <sup>m</sup>	-16853.4 1.1	4895.23 0.05	500 ns 10	10 <sup>+</sup>	07	98Pf02 T	1998	IT=100
<sup>208</sup> Bi	-18870.2 2.3		368 ky 4	5 <sup>+</sup>	07		1953	$\beta^+ =100$
<sup>208</sup> Bi <sup>m</sup>	-17299.1 2.3	1571.1 0.4	2.58 ms 0.04	10 <sup>-</sup>	07		1961	IT=100
<sup>208</sup> Po	-17469.6 1.7		2.898 y 0.002	0 <sup>+</sup>	07		1947	$\alpha \approx 100$ ; $\beta^+ =0.0040 4$
<sup>208</sup> Po <sup>m</sup>	-15941.4 1.7	1528.22 0.04	350 ns 20	8 <sup>+</sup>	07		1968	IT=100
<sup>208</sup> At	-12470 9		1.63 h 0.03	6 <sup>+</sup>	07		1950	$\beta^+ =99.45 6$ ; $\alpha =0.55 6$
<sup>208</sup> At <sup>m</sup>	-10194 9	2276.4 1.8	1.5 $\mu$ s 0.2	16 <sup>-</sup>	07		1991	IT=100
<sup>208</sup> Rn	-9656 11		24.35 m 0.14	0 <sup>+</sup>	07		1955	$\alpha =62 7$ ; $\beta^+ =38 7$
<sup>208</sup> Rn <sup>m</sup>	-7828 11	1828.3 0.4	487 ns 12	8 <sup>+</sup>	07		1979	IT=100
<sup>208</sup> Fr	-2666 12		59.1 s 0.3	7 <sup>+</sup>	07	78Ek02 J	1964	$\alpha =89 3$ ; $\beta^+ =11 3$
<sup>208</sup> Fr <sup>m</sup>	-1839 22	827 18	432 ns 11	(10 <sup>-</sup> )	07	09Dr08 T	2009	IT=100
<sup>208</sup> Ra	1728 9		1.110 s 0.045	0 <sup>+</sup>	07	10He25 TD	1967	$\alpha =87 3$ ; $\beta^+ ?$
<sup>208</sup> Ra <sup>m</sup>	3875 9	2147.4 0.4	263 ns 17	(8 <sup>+</sup> )	07	05Re02 T	1998	IT=100
<sup>208</sup> Ac	10750 60		97 ms 15	(3 <sup>+</sup> )	07	14Ya19 T	1994	$\alpha =?$ ; $\beta^+ =1\#$
<sup>208</sup> Ac <sup>m</sup>	11258 28	500 50 AD	28 ms 7	(10 <sup>-</sup> )	07	96lk01 T	1994	$\alpha =?$ ; IT<10#; $\beta^+ =1\#$
<sup>208</sup> Th	16680 30		2.4 ms 1.2	0 <sup>+</sup>	11		2010	$\alpha \approx 100$
* <sup>208</sup> Hg	T : symmetrized from 98Zh22=41(+5-4); other 16Ca25=132(50) s							
* <sup>208</sup> Hg <sup>m</sup>	E : 1296.9(0.9) + x and x<83 keV							
* <sup>208</sup> Rn <sup>m</sup>	T : other 10Ka29=590(144) ns							
* <sup>208</sup> Fr <sup>m</sup>	T : from lifetime 09Dr08=623(16); other 10Ka29=233(18), not trusted							
* <sup>208</sup> Fr <sup>m</sup>	T : also 06Me03=446(14) originally assigned to <sup>209</sup> Fr, see 09Dr04							
* <sup>208</sup> Ra	T : other 68Lo15=1.8(0.5) 67Va22=1.2(0.2)							
* <sup>208</sup> Ra <sup>m</sup>	T : average 05Re02=250(30) 99Co13=270(21)							
* <sup>208</sup> Ac	T : average 14Ya19=93(+40-22) 96lk01=83(+34-19) 94Le05=95(+24-16)							
* <sup>208</sup> Ac <sup>m</sup>	E : if $\alpha$ decay goes to (7 <sup>+</sup> ) <sup>204</sup> Fr <sup>m</sup> , instead of (10 <sup>-</sup> ) as assumed in AME, then							
* <sup>208</sup> Ac <sup>m</sup>	E : E will become 234(22) keV							
* <sup>208</sup> Ac <sup>m</sup>	T : average 96lk01=21(+28-8) 94Le05=25(+9-5)							
* <sup>208</sup> Th	T : symmetrized from 10He25=1.7(+1.7-0.6)							
<sup>209</sup> Au	-2540# 400#		1# s (>300 ns)	3/2 <sup>+</sup> #	15	10A124 I	2010	$\beta^- ?$ ; $\beta^- n=90\#$
<sup>209</sup> Hg	-8640# 150#		38 s 6	9/2 <sup>+</sup> #	15		1998	$\beta^- =100$ ; $\beta^- n=0\#$
<sup>209</sup> Tl	-13645 6		2.162 m 0.007	1/2 <sup>+</sup>	15		1950	$\beta^- =100$ ; $\beta^- n=0\#$
<sup>209</sup> Pb	-17614.6 1.7		3.234 h 0.007	9/2 <sup>+</sup>	15		1940	$\beta^- =100$
<sup>209</sup> Bi	-18258.7 1.4		20.1 Ey 0.8	9/2 <sup>-</sup>	15		1924	IS=100.; $\alpha =100$
<sup>209</sup> Po	-16366.1 1.8		124 y 3	1/2 <sup>-</sup>	15	13Se03 J	1949	$\alpha \approx 100$ ; $\beta^+ =0.454 7$
<sup>209</sup> Po <sup>m</sup>	-12100.7 1.8	4265.4 0.3	119 ns 4	31/2 <sup>-</sup>	15		1974	IT=100
<sup>209</sup> At	-12883 5		5.42 h 0.05	9/2 <sup>-</sup>	15		1951	$\beta^+ =95.9 5$ ; $\alpha =4.1 5$
<sup>209</sup> At <sup>m</sup>	-10454 5	2429.32 0.22	916 ns 10	29/2 <sup>+</sup>	15		1975	IT=100
<sup>209</sup> Rn	-8941 10		28.8 m 1.0	5/2 <sup>-</sup>	15		1952	$\beta^+ =83 2$ ; $\alpha =17 2$
<sup>209</sup> Rn <sup>m</sup>	-7767 10	1174.01 0.13	13.4 $\mu$ s 1.3	13/2 <sup>+</sup>	15		1985	IT=100
<sup>209</sup> Rn <sup>n</sup>	-5304 10	3636.81 0.23	3.0 $\mu$ s 0.3	35/2 <sup>+</sup>	15		1985	IT=100
<sup>209</sup> Fr	-3770 15		50.5 s 0.7	9/2 <sup>-</sup>	15	78Ek02 J	1964	$\alpha =89 3$ ; $\beta^+ =11 3$
<sup>209</sup> Fr <sup>m</sup>	890 15	4659.8 0.7	420 ns 18	45/2 <sup>-</sup>	15		2006	IT=100
<sup>209</sup> Ra	1858 6		4.71 s 0.08	5/2 <sup>-</sup>	15	08Ha12 T	1967	$\alpha \approx 100$ ; $\beta^+ ?$
<sup>209</sup> Ra <sup>m</sup>	2740 6	882.4 0.7	117 $\mu$ s 5	13/2 <sup>+</sup>	15	08Ha12 D	2008	$\alpha \approx 90$ ; $\beta^+ \approx 10$
<sup>209</sup> Ac	8840 50		94 ms 10	(9/2 <sup>-</sup> )	15	14Ya19 T	1968	$\alpha =?$ ; $\beta^+ =1\#$
<sup>209</sup> Th	16370# 140#		60# ms	5/2 <sup>-</sup> #				$\alpha ?$ ; $\beta^+ ?$
<sup>209</sup> Th <sup>m</sup>	16840# 100#	470# 100#	3.1 ms 1.2	(13/2 <sup>+</sup> )	15		1996	$\alpha \approx 100$ ; $\beta^+ ?$
* <sup>209</sup> Hg	T : symmetrized from ENSDF2015=36(+7-4); other 16Ca25=6(1)							
* <sup>209</sup> Fr <sup>m</sup>	T : from lifetime 09Dr04=606(26);							
* <sup>209</sup> Ac	T : average 14Ya19=98(22) 00He17=98(+59-27) 96lk01=82(+18-13)							
* <sup>209</sup> Ac	T : 94Le05=91(+21-14) and 68Va04=100(50)							
* <sup>209</sup> Th <sup>m</sup>	T : symmetrized from ENSDF2015=2.5(+1.7-0.7)							
<sup>210</sup> Au	2330# 400#		1# s (>300 ns)		14	10A124 I	2010	$\beta^- ?$ ; $\beta^- n=10\#$
<sup>210</sup> Hg	-5370# 200#		64 s 10	0 <sup>+</sup>	14	16Ca25 TD	1998	$\beta^- =100$ ; $\beta^- n=2.2 22$
<sup>210</sup> Hg <sup>m</sup>	-4710# 200#	663 2	2.1 $\mu$ s 0.7	(3 <sup>-</sup> )	14		2013	IT=100
<sup>210</sup> Hg <sup>n</sup>	-3960# 200#	1406 23	2 $\mu$ s 1	8 <sup>+</sup> #	14	13Go10 E	2013	IT=100
<sup>210</sup> Tl	-9247 12		1.30 m 0.03	5 <sup>+</sup> #	14		1909	$\beta^- =100$ ; $\beta^- n=0.009 6$
<sup>210</sup> Pb	-14728.5 1.4		22.20 y 0.22	0 <sup>+</sup>	14		1900	$\beta^- =100$ ; $\alpha =1.9e-6 4$
<sup>210</sup> Pb <sup>m</sup>	-13451 5	1278 5	201 ns 17	8 <sup>+</sup>	14		1980	IT=100
<sup>210</sup> Bi	-14792.0 1.4		5.012 d 0.005	1 <sup>-</sup>	14		1905	$\beta^- =100$ ; $\alpha =13.2e-5 10$
<sup>210</sup> Bi <sup>m</sup>	-14520.7 1.4	271.31 0.11	3.04 My 0.06	9 <sup>-</sup>	14		1953	$\alpha =100$

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)					
... A-group continued ...													
<sup>210</sup> Po	-15953.1	1.1	138.376 d	0.002	0 <sup>+</sup>	14	1898	$\alpha=100$					
<sup>210</sup> Po <sup>m</sup>	-14396.1	1.1	1556.97	0.03	98.9 ns	2.5	1968	IT=100					
<sup>210</sup> Po <sup>n</sup>	-10895.5	1.1	5057.65	0.05	263 ns	5	1985	IT=100					
<sup>210</sup> At	-11972	8	8.1 h	0.4	(5) <sup>+</sup>	14	1949	$\beta^+\approx 100; \alpha=0.175$ 20					
<sup>210</sup> At <sup>m</sup>	-9422	8	2549.6	0.2	482 ns	6	1970	IT=100					
<sup>210</sup> At <sup>n</sup>	-7944	8	4027.7	0.2	5.66 $\mu$ s	0.07	1975	IT=100					
<sup>210</sup> Rn	-9605	5	2.4 h	0.1	0 <sup>+</sup>	14	1952	$\alpha=96$ 1; $\beta^+$ ?					
<sup>210</sup> Rn <sup>m</sup>	-7900	30	1710	30	AD	644 ns	40	1979	IT ?				
<sup>210</sup> Rn <sup>n</sup>	-5750	30	3857	30	1.06 $\mu$ s	0.05	(17) <sup>-</sup>	14	1979	IT=100			
<sup>210</sup> Rn <sup>p</sup>	-3090	30	6514	30	1.04 $\mu$ s	0.07	(23) <sup>+</sup>	14	1986	IT=100			
<sup>210</sup> Fr	-3333	15	3.18 m	0.06	6 <sup>+</sup>	14	05Ku06	D	1964	$\alpha=71$ 4; $\beta^+$ ?			
<sup>210</sup> Ra	443	9	4.0 s	0.1	0 <sup>+</sup>	14	08Ha12	TJ	1967	$\alpha=?; \beta^+=4\#$			
<sup>210</sup> Ra <sup>m</sup>	2494	9	2050.9	0.7	2.29 $\mu$ s	0.03	8 <sup>+</sup>	14	04Re04	TJ	1998	IT=100	
<sup>210</sup> Ac	8790	60	350 ms	40	7 <sup>+</sup> #	14	00He17	T	1968	$\alpha=?; \beta^+=9\#$			
<sup>210</sup> Th	14059	19	16.0 ms	3.6	0 <sup>+</sup>	14	1995	$\alpha=?; \beta^+=1\#$					
* <sup>210</sup> Hg <sup>n</sup>	E : from 13Go10; stated to be less than 80 keV above 1366 level												
* <sup>210</sup> Tl	D : symmetrized from $\beta^-$ n=0.007(+7-4)%												
* <sup>210</sup> Rn <sup>n</sup>	E : ENSDF2014: 2147.4(0.2) keV above the 8 <sup>+</sup> level, quoted 3812.40(0.16) + x												
* <sup>210</sup> Rn <sup>p</sup>	E : ENSDF2014: 4803.7(0.4) keV above the 8 <sup>+</sup> level, quoted 6469.02(0.21) + x												
* <sup>210</sup> Ra	T : also 07Le14=2.5(+1.4-0.7) and 3.5(+4.8-1.3)												
* <sup>210</sup> Ra <sup>m</sup>	T : average 13Ba29=2.1(0.1) 06Ha17=2.28(0.08) 04Re04=2.1(0.1) 04He25=2.36(0.04)												
* <sup>210</sup> Ac	T : average 00He17=335(+64-46) 68Va04=350(50)												
<sup>211</sup> Hg	-620#	200#	26 s	8	9/2 <sup>+</sup> #	13	16Ca25	TD	2010	$\beta^-$ =100; $\beta^-$ n=6.3 63			
<sup>211</sup> Tl	-6080	40	80 s	16	1/2 <sup>+</sup>	13	14Mo02	J	1998	$\beta^-$ =100; $\beta^-$ n=2.2 22			
<sup>211</sup> Pb	-10492.9	2.3	36.164 m	0.012	9/2 <sup>+</sup>	13	16Ai01	T	1904	$\beta^-$ =100			
<sup>211</sup> Pb <sup>m</sup>	-8789	15	1704	15	159 ns	28	(27/2 <sup>+</sup> )	13	05La01	ET	2005	IT=100	
<sup>211</sup> Bi	-11859	5	2.14 m	0.02	9/2 <sup>-</sup>	13	1905	$\alpha\approx 100; \beta^-$ =0.276 4					
<sup>211</sup> Bi <sup>m</sup>	-10602	11	1257	10	1.4 $\mu$ s	0.3	(25/2 <sup>-</sup> )	13	1998	IT=100			
<sup>211</sup> Po	-12432.6	1.3	516 ms	3	9/2 <sup>+</sup>	15	13Se03	J	1913	$\alpha=100$			
<sup>211</sup> Po <sup>m</sup>	-10970	5	1462	5	AD	25.2 s	6	(25/2 <sup>+</sup> )	15	1954	$\alpha\approx 100; IT=0.016$ 4		
<sup>211</sup> Po <sup>n</sup>	-10298	5	2135	5	243 ns	21	(31/2 <sup>-</sup> )	15	1998	IT $\approx$ 100; $\alpha$ ?			
<sup>211</sup> Po <sup>p</sup>	-7561	6	4872	6	2.8 $\mu$ s	0.7	(43/2 <sup>+</sup> )	15	1998	IT $\approx$ 100; $\alpha$ ?			
<sup>211</sup> At	-11647.3	2.7	7.214 h	0.007	9/2 <sup>-</sup>	13	1940	$\epsilon=58.20$ 8; $\alpha=41.80$ 8					
<sup>211</sup> At <sup>m</sup>	-6832.8	2.7	4814.5	0.5	4.23 $\mu$ s	0.07	(39/2 <sup>-</sup> )	13	1971	IT=100			
<sup>211</sup> Rn	-8755	7	14.6 h	0.2	1/2 <sup>-</sup>	13	1952	$\beta^+=72.6$ 17; $\alpha=27.4$ 17					
<sup>211</sup> Rn <sup>m</sup>	-7152	16	1603	14	596 ns	28	(17/2 <sup>-</sup> )	13	1981	IT=100			
<sup>211</sup> Rn <sup>n</sup>	125	16	8880	14	201 ns	4	(63/2 <sup>-</sup> )	13	1981	IT=100			
<sup>211</sup> Fr	-4140	12	3.10 m	0.02	9/2 <sup>-</sup>	13	05Ku06	D	1964	$\alpha=87$ 3; $\beta^+$ ?			
<sup>211</sup> Fr <sup>m</sup>	-1717	12	2423.16	0.24	146 ns	14	(29/2 <sup>+</sup> )	13	1986	IT=100			
<sup>211</sup> Fr <sup>n</sup>	517	12	4657.3	0.4	123 ns	14	(45/2 <sup>-</sup> )	13	1986	IT=100			
<sup>211</sup> Ra	832	8	13.2 s	1.4	5/2 <sup>(-)</sup>	13	07Le14	T	1967	$\alpha>93; \beta^+<7$			
<sup>211</sup> Ra <sup>m</sup>	2030	8	1198.1	0.8	9.5 $\mu$ s	0.3	13/2 <sup>+</sup>	13	13Ba29	T	2004	IT=100	
<sup>211</sup> Ac	7200	50	213 ms	25	9/2 <sup>-</sup>	13	00He17	T	1968	$\alpha\approx 100; \beta^+<0.2$			
<sup>211</sup> Th	13910	70	48 ms	20	5/2 <sup>-</sup> #	13	1995	$\alpha=?; \beta^+=0.5\#$					
<sup>211</sup> Pa	22080#	100#	3# ms	(>300 ns)	9/2 <sup>-</sup> #	13	2006	$\alpha ?; \beta^+ ?; p ?$					
* <sup>211</sup> Tl	T : average 16Ca25=76(18) 12Be28=88(+46-29) D : $\beta^-$ n 16Ca25=2.2 22												
* <sup>211</sup> Pb	T : average 16Ai01=36.164(0.013) 15Ko09=36.165(0.037)												
* <sup>211</sup> Pb <sup>m</sup>	E : E=1679.1 + x in 05La01, where x<50 keV												
* <sup>211</sup> Rn <sup>m</sup>	E : 1577.8 + x ; x<50												
* <sup>211</sup> Rn <sup>n</sup>	E : 8854.5(0.4) + y ; y<50												
* <sup>211</sup> Ra	T : average 07Le14=9(5) 68Lo15=12(2) 67Va22=15(2)												
* <sup>211</sup> Ra <sup>m</sup>	T : average 13Ba29=9.4(0.4) 06Ha17=9.7(0.6); other 04He25=4.0(0.5)												
* <sup>211</sup> Ac	T : average 00He17=200(29) 68Va04=250(50)												
* <sup>211</sup> Th	T : symmetrized from 95Uu01=37(+28-11); other 15Ya13=20.8(+37.9-8.2)(2 evts)												
<sup>212</sup> Hg	2760#	300#	1# m	(>300 ns)	0 <sup>+</sup>	11	10Al24	I	2010	$\beta^-$ ?; $\beta^-$ n=8#			
<sup>212</sup> Tl	-1550#	200#	31 s	8	(5 <sup>+</sup> )	12	16Ca25	TD	1998	$\beta^-$ =100; $\beta^-$ n=1.8 18			
<sup>212</sup> Pb	-7548.8	1.8	10.64 h	0.01	0 <sup>+</sup>	05	1905	$\beta^-$ =100					
<sup>212</sup> Pb <sup>m</sup>	-6213.8	2.7	1335	2	6.0 $\mu$ s	0.8	8 <sup>+</sup> #	05	12Re.B	E	1998	IT=100	
<sup>212</sup> Bi	-8118.0	1.9	60.55 m	0.06	1 <sup>(-)</sup>	05	89Ha.A	D	1905	$\beta^-$ =64.06 6; $\alpha=35.94$ 6; $\beta^-$ $\alpha=0.014$			
<sup>212</sup> Bi <sup>m</sup>	-7870	30	250	30	AD	25.0 m	0.2	(8 <sup>-</sup> , 9 <sup>-</sup> )	05	1978	$\alpha=67$ 1; $\beta^-$ =33 1; $\beta^-$ $\alpha=30$ 1		
<sup>212</sup> Bi <sup>n</sup>	-6639	30	1479	30	MD	7.0 m	0.3	> 16	05	13Ch12	D	1978	$\beta^-$ =?; IT>75
<sup>212</sup> Po	-10369.5	1.2	294.7 ns	1.0	0 <sup>+</sup>	05	13Be31	T	1906	$\alpha=100$			
<sup>212</sup> Po <sup>m</sup>	-7446	5	2923	4	AD	45.1 s	0.6	(18 <sup>+</sup> )	05	1962	$\alpha\approx 100; IT=0.07$ 2		
... A-group is continued on next page ...													

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)						
... A-group continued ...														
<sup>212</sup> At	-8628.2	2.4	314 ms	2	(1 <sup>-</sup> )	05	1954	$\alpha \approx 100$ ; $\beta^+ < 0.03$ ; $\beta^- < 2e-6$						
<sup>212</sup> At <sup>m</sup>	-8405.3	2.4	119 ms	3	9 <sup>-</sup> #	05	1970	$\alpha > 99$ ; IT < 1						
<sup>212</sup> At <sup>l</sup>	-3856.6	2.6	4771.6	1.1	152 $\mu$ s	5	1998	IT=100						
<sup>212</sup> Rn	-8660	3	23.9 m	1.2	0 <sup>+</sup>	05	1950	$\alpha = 100$ ; $2\beta^+ ?$						
<sup>212</sup> Rn <sup>m</sup>	-7020	3	1639.8	0.3	118 ns	14	1971	IT=100						
<sup>212</sup> Rn <sup>n</sup>	-6966	3	1694.0	0.4	910 ns	30	1971	IT=100						
<sup>212</sup> Rn <sup>p</sup>	-2486	3	6174.0	0.4	104.0 ns	2.8	1977	IT=100						
<sup>212</sup> Rn <sup>q</sup>	-81	3	8579.0	0.5	154 ns	14	1977	IT=100						
<sup>212</sup> Fr	-3516	9	20.0 m	0.6	5 <sup>+</sup>	05	1950	$\beta^+ = 57.2$ ; $\alpha = 43.2$						
<sup>212</sup> Fr <sup>m</sup>	-1965	9	1551.4	0.3	31.9 $\mu$ s	0.7	1977	IT=100						
<sup>212</sup> Fr <sup>n</sup>	-1024	9	2492.2	0.4	604 ns	28	1977	IT=100						
<sup>212</sup> Fr <sup>p</sup>	2339	9	5854.7	0.6	312 ns	21	1986	IT=100						
<sup>212</sup> Fr <sup>q</sup>	5017	9	8533.4	1.1	23.6 $\mu$ s	2.1	1990	IT=100						
<sup>212</sup> Ra	-199	11	13.0 s	0.2	0 <sup>+</sup>	05	1967	$\alpha = ?$ ; $\beta^+ = 15\%$						
<sup>212</sup> Ra <sup>m</sup>	1759	11	1958.4	0.5	8.1 $\mu$ s	0.7	1986	IT=100						
<sup>212</sup> Ra <sup>n</sup>	2414	11	2613.4	0.5	512 ns	104	1986	IT=100						
<sup>212</sup> Ac	7280	50	895 ms	28	6 <sup>+</sup> #	05	1968	$\alpha = ?$ ; $\beta^+ = 3\%$						
<sup>212</sup> Th	12111	10	31.7 ms	1.3	0 <sup>+</sup>	15	1980	$\alpha \approx 100$ ; $\beta^- = 0.3\%$						
<sup>212</sup> Pa	21590	70	7.5 ms	2.8	7 <sup>+</sup> #	05	1997	$\alpha = 100$						
* <sup>212</sup> Pb <sup>m</sup>	T : 12Go19=6.0(0.8) supersedes 12Re.B=5.0(0.3); other 98Pf02=5(1)													
* <sup>212</sup> Bi <sup>n</sup>	D : IT not observed, deduced from half-life > 30 m for highly charged ions													
* <sup>212</sup> Rn <sup>m</sup>	J : E2 to 4 <sup>+</sup> for <sup>212</sup> Rn <sup>m</sup> ; E2 to 6 <sup>+</sup> for <sup>212</sup> Rn <sup>n</sup> ; magnetic moment measurement													
* <sup>212</sup> Ra <sup>m</sup>	T : average 13Ba29=7.1(0.2) 06Ha17=9.7(0.6) 04He25=8.31(0.25) 86Ko01=10.9(0.4)													
* <sup>212</sup> Ra <sup>n</sup>	J : 63.3 keV $\gamma$ to 6 <sup>+</sup> ; no $\gamma$ to 2 <sup>+</sup> and 4 <sup>+</sup> ; measured magnetic moment													
* <sup>212</sup> Ra <sup>l</sup>	T : average 13Ba29=480(40) 86Ko01=850(130)													
* <sup>212</sup> Ra <sup>l</sup>	J : 655 keV $\gamma$ E3 to 8 <sup>+</sup> ; measured magnetic moment													
* <sup>212</sup> Ac	T : average 14Ya19=880(35) 00He17=880(110) 68Va04=930(50)													
* <sup>212</sup> Ac	J : ENSDF proposes to assign 7 <sup>+</sup> , if the observed $\alpha$ feeds the <sup>208</sup> Fr 7 <sup>+</sup> ground-state													
* <sup>212</sup> Pa	T : average 14Ya10=5.1(+5.1-1.7) 97Mi03=5.1(+6.1-1.9)													
<sup>213</sup> Hg	7670#	300#	1#	s	(>300 ns)	5/2 <sup>+</sup> #	11	10Al24	I	2010	$\beta^- ?$ ; $\beta^- n = 30\%$			
<sup>213</sup> Tl	1784	27	24	s	4	1/2 <sup>+</sup>	12	16Ca25	TD	2010	$\beta^- = 100$ ; $\beta^- n = 7.6$ 34			
<sup>213</sup> Pb	-3204	7	10.2	m	0.3	(9/2 <sup>+</sup> )	07			1964	$\beta^- = 100$			
<sup>213</sup> Bi	-5232	5	45.61	m	0.04	9/2 <sup>-</sup>	07	13Ma13	T	1947	$\beta^- = 97.91$ 3; $\alpha = 2.09$ 3			
<sup>213</sup> Bi <sup>m</sup>	-3930#	200#	1300#	200#	> 168	s	25/2 <sup>-</sup> #	08Ch.A	T	2008				
<sup>213</sup> Po	-6654	3	3.708	$\mu$ s	0.008	9/2 <sup>+</sup>	07	13Su13	T	1947	$\alpha = 100$			
<sup>213</sup> At	-6580	5	125	ns	6	9/2 <sup>-</sup>	07			1968	$\alpha = 100$			
<sup>213</sup> At <sup>m</sup>	-5210	50	110	ns	17		07			1980	IT=100			
<sup>213</sup> At <sup>l</sup>	-3600	50	2980	50	45	$\mu$ s	4	(49/2 <sup>+</sup> )	07	2003	IT=100			
<sup>213</sup> Rn	-5696	3	19.5	ms	0.1	9/2 <sup>+</sup> #	07			1967	$\alpha = 100$			
<sup>213</sup> Rn <sup>m</sup>	-3990	50	1710	50	1.00	$\mu$ s	0.21	(25/2 <sup>+</sup> )	07	1988	IT=100			
<sup>213</sup> Rn <sup>n</sup>	-3460	50	2240	50	1.36	$\mu$ s	0.07	(31/2 <sup>-</sup> )	07	1988	IT=100			
<sup>213</sup> Rn <sup>p</sup>	280	50	5980	50	164	ns	11	(55/2 <sup>+</sup> )	07	1988	IT=100			
<sup>213</sup> Fr	-3553	5	34.14	s	0.06	9/2 <sup>-</sup>	07	13Fi08	T	1964	$\alpha = 99.44$ 5; $\beta^+ = 0.56$ 5			
<sup>213</sup> Fr <sup>m</sup>	-1963	5	1590.41	0.18	505	ns	14	21/2 <sup>-</sup>	07	1971	IT=100			
<sup>213</sup> Fr <sup>n</sup>	-1015	5	2537.62	0.23	238	ns	6	29/2 <sup>+</sup>	07	1971	IT=100			
<sup>213</sup> Fr <sup>p</sup>	4542	5	8094.8	0.7	3.1	$\mu$ s	0.2	(65/2 <sup>-</sup> )	07	1989	IT=100			
<sup>213</sup> Ra	346	10	2.73	m	0.05	1/2 <sup>-</sup>	07			1955	$\alpha = 80$ 5; $\beta^+ ?$			
<sup>213</sup> Ra <sup>m</sup>	2114	11	1768	4	AD	2.20	ms	0.05	(17/2 <sup>-</sup> )	07	06Ku26	TD	1976	IT $\approx$ 99; $\alpha = 0.6$ 4
<sup>213</sup> Ac	6155	15	738	ms	16	9/2 <sup>-</sup> #	07			1968	$\alpha = ?$ ; $\beta^+ ?$			
<sup>213</sup> Th	12120	9	144	ms	21	5/2 <sup>-</sup> #	07			1968	$\alpha = ?$ ; $\beta^+ = 1.4\%$			
<sup>213</sup> Th <sup>m</sup>	13300	9	1180	3	1.4	$\mu$ s	0.4	13/2 <sup>+</sup> #	07	07Kh22	TD	2007	IT=100	
<sup>213</sup> Th <sup>p</sup>	12380#	50#	260#	50#										
<sup>213</sup> Pa	19660	70	7	ms	3	9/2 <sup>-</sup> #	07	95Ni05	TD	1995	$\alpha = 100$			
* <sup>213</sup> Bi	T : average 13Ma13=45.62(0.06) 73Po16=45.59(0.06)													
* <sup>213</sup> At <sup>m</sup>	E : 1318.1(0.6) + x ; x estimated 50(50) by NUBASE													
* <sup>213</sup> At <sup>l</sup>	E : 2926 + y ; y estimated 50(50) by NUBASE													
* <sup>213</sup> Rn <sup>m</sup>	E : 1664.0(1.0) + x ; x=50(50) estimated by NUBASE													
* <sup>213</sup> Rn <sup>n</sup>	E : 2186.7 + x ; x=50(50) estimated by NUBASE													
* <sup>213</sup> Rn <sup>p</sup>	E : 5929 + y ; y=50(50) estimated by NUBASE													
* <sup>213</sup> Fr	T : see discussion of previous results in 13Fi08													
* <sup>213</sup> Ra <sup>m</sup>	E : derived from difference in $\alpha$ decay energy in the AME evaluation.													
* <sup>213</sup> Ra <sup>n</sup>	E : <sup>76</sup> Ra37 less than 10 keV above 1769.7 level, thus 1775(3) keV													
* <sup>213</sup> Ra <sup>m</sup>	J : 17/2 <sup>-</sup> or 13/2 <sup>+</sup> as proposed in <sup>76</sup> Ra37													
* <sup>213</sup> Th <sup>m</sup>	E : uncertainty estimated by NUBASE													
* <sup>213</sup> Pa	T : symmetrized from 5.3(+4.0-1.6)													

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
$^{214}\text{Hg}$	11180# 400#		1# s (>300 ns)	$0^+$	11	10Al24 I	2010	$\beta^-$ ?; $\beta^-n=10\#$	
$^{214}\text{Tl}$	6470# 200#		11 s 2	$5^+\#$	11	16Ca25 TD	2010	$\beta^-n=100$ ; $\beta^-n=34$ 12	
$^{214}\text{Pb}$	-182.8 2.0		27.06 m 0.07	$0^+$	15		1904	$\beta^-n=100$	
$^{214}\text{Pb}^m$	1237 20	1420 20	6.2 $\mu\text{s}$ 0.3	$8^+\#$	15		2012	IT=100 *	
$^{214}\text{Bi}$	-1201 11		19.9 m 0.4	$1^-$	09	89Ha.A D	1904	$\beta^-n\approx 100$ ; $\alpha=0.021$ 1; $\beta^-n\alpha=0.003$	
$^{214}\text{Bi}^m$	-1000# 100#	200# 100#	> 93 s	$8^-\#$		08Ch.A T	2008		
$^{214}\text{Po}$	-4470.0 1.4		163.72 $\mu\text{s}$ 0.27	$0^+$	09	13Be31 T	1912	$\alpha=100$ *	
$^{214}\text{At}$	-3380 4		558 ns 10	$1^-$	09		1949	$\alpha=100$	
$^{214}\text{At}^m$	-3321 8	59 9	AD	265 ns 30	09		1982	$\alpha<100$	
$^{214}\text{At}^n$	-3146 5	234 6	AD	760 ns 15	09		1982	$\alpha<100$	
$^{214}\text{Rn}$	-4320 9		270 ns 20	$0^+$	09		1970	$\alpha=100$	
$^{214}\text{Rn}^m$	275 9	4595.4 1.8		245 ns 30	(22 <sup>+</sup> )	09	1983	IT=100	
$^{214}\text{Fr}$	-959 9		5.18 ms 0.16	(1 <sup>-</sup> )	09	15Kh09 T	1967	$\alpha=100$ *	
$^{214}\text{Fr}^m$	-837 8	122 5	AD	3.35 ms 0.05	(8 <sup>-</sup> )	09	1962	$\alpha=100$	
$^{214}\text{Fr}^n$	-321 10	638 5		103 ns 4	(11 <sup>+</sup> )	09	1993	IT=100	
$^{214}\text{Fr}^p$	5620 100	6580 100		108 ns 7	(33 <sup>+</sup> )	09	1994	IT ? *	
$^{214}\text{Ra}$	93 5		2.437 s 0.016	$0^+$	09	15Kh09 T	1967	$\alpha\approx 100$ ; $\beta^+=0.059$ 4 *	
$^{214}\text{Ra}^m$	1913 5	1819.7 1.8		118 ns 7	$6^+$	09	2004	IT=100	
$^{214}\text{Ra}^n$	1958 5	1865.2 1.8		67.3 $\mu\text{s}$ 1.5	$8^+$	09	1971	IT $\approx 100$ ; $\alpha=0.09$ 7	
$^{214}\text{Ra}^p$	2776 5	2683.2 1.8		295 ns 7	$11^-$	09	1979	IT=100	
$^{214}\text{Ra}^q$	3571 5	3478.4 1.8		279 ns 4	$14^+$	09	1979	IT=100	
$^{214}\text{Ra}^r$	4240 5	4146.8 1.8		225 ns 4	$17^-$	09	1979	IT=100	
$^{214}\text{Ra}^s$	6670 5	6577.0 1.8		128 ns 4	(25 <sup>-</sup> )	09	1992	IT=100	
$^{214}\text{Ac}$	6444 15		8.2 s 0.2	$5^+\#$	09		1968	$\alpha>89$ 3; $\beta^+<11$ 3	
$^{214}\text{Th}$	10695 11		87 ms 10	$0^+$	09		1968	$\alpha\approx 100$ ; $\beta^+=0.1\#$	
$^{214}\text{Th}^m$	12876 11	2181.0 2.7	1.24 $\mu\text{s}$ 0.12	$8^+\#$	09		2007	IT=100	
$^{214}\text{Pa}$	19490 80		17 ms 3		09	95Ni05 D	1995	$\alpha\approx 100$	
* $^{214}\text{Pb}^m$	E : 1365 + x ; x=20-90 keV **								
* $^{214}\text{Po}$	T : average 13Be31=163.6(0.3) 12Su11=164.2(0.6) **								
* $^{214}\text{Fr}$	T : average 15Kh09=5.9(0.4) 05Li17=4.6(0.7) 68To10=5.0(0.2) 68Va18=5.5(0.5) **								
* $^{214}\text{Fr}^p$	E : 6477 + y ; y=100(100) estimated by NUBASE **								
* $^{214}\text{Ra}$	T : average 15Kh09=2.36(0.06) 12No08=2.435(0.020) 73Be33=2.46(0.03) **								
$^{215}\text{Hg}$	16210# 400#		1# s (>300 ns)	$3/2^+\#$	13	10Al24 I	2010	$\beta^-$ ?; $\beta^-n=4\#$	
$^{215}\text{Tl}$	9910# 300#		10 s 4	$1/2^+\#$	13	16Ca25 TD	2010	$\beta^-n=100$ ; $\beta^-n=4.6$ 46	
$^{215}\text{Pb}$	4340 50		2.34 m 0.19	$9/2^+\#$	13	16Ca25 T	1998	$\beta^-n=100$ *	
$^{215}\text{Bi}$	1629 6		7.6 m 0.2	(9/2 <sup>-</sup> )	13		1953	$\beta^-n=100$	
$^{215}\text{Bi}^m$	2996 21	1367 20	36.9 s 0.6	(25/2 <sup>-</sup> )	13		2001	IT=76.9 5; $\beta^-n=23.1$ 5 *	
$^{215}\text{Po}$	-541.7 2.1		1.781 ms 0.005	$9/2^+$	13		1911	$\alpha=100$ ; $\beta^-n=2.3e-4$ 2	
$^{215}\text{At}$	-1256 7		100 $\mu\text{s}$ 20	$9/2^-$	13		1944	$\alpha=100$	
$^{215}\text{Rn}$	-1169 8		2.30 $\mu\text{s}$ 0.10	$9/2^+$	13		1952	$\alpha=100$	
$^{215}\text{Fr}$	318 7		86 ns 5	$9/2^-$	13		1970	$\alpha=100$	
$^{215}\text{Ra}$	2534 8		1.67 ms 0.01	$9/2^+\#$	13		1967	$\alpha=100$	
$^{215}\text{Ra}^m$	4412 8	1877.8 0.3	7.31 $\mu\text{s}$ 0.13	(25/2 <sup>+</sup> )	13	04He25 T	1983	IT=100 *	
$^{215}\text{Ra}^n$	4781 8	2246.9 0.4	1.39 $\mu\text{s}$ 0.07	(29/2 <sup>-</sup> )	13		1998	IT=100	
$^{215}\text{Ra}^p$	6340 50	3810 50	555 ns 10	(43/2 <sup>-</sup> )	13		1987	IT=100 *	
$^{215}\text{Ac}$	6031 12		170 ms 10	$9/2^-$	13		1968	$\alpha\approx 100$ ; $\beta^+=0.09$ 2	
$^{215}\text{Ac}^m$	7827 12	1796.0 0.9	185 ns 30	(21/2 <sup>-</sup> )	13		1983	IT=100	
$^{215}\text{Ac}^n$	8520 50	2490 50	335 ns 10	(29/2 <sup>+</sup> )	13		1983	IT=100 *	
$^{215}\text{Th}$	10922 9		1.2 s 0.2	(1/2 <sup>-</sup> )	13		1968	$\alpha=100$ *	
$^{215}\text{Th}^m$	12390 50	1470 50	770 ns 60	$9/2^+\#$	13		2005	IT=100 *	
$^{215}\text{Pa}$	17860 70		14 ms 2	$9/2^-\#$	13		1979	$\alpha=100$	
$^{215}\text{U}$	24920 90		1.4 ms 0.9	$5/2^-\#$	15	15Ya13 T	2015	$\alpha>0$ ; $\beta^+ ?$ *	
* $^{215}\text{Pb}$	T : average 16Ca25=98(30)s 13De20=147(12)s; others 14Mo02=160(40) 96Ry.B=36(1) **								
* $^{215}\text{Bi}^m$	E : 1347.5(0.2) + x ; x=20(20) estimated by NUBASE **								
* $^{215}\text{Ra}^m$	T : average 04He25=7.6(0.2) 98St24=6.9(0.3) 88Fu10=7.2(0.2) **								
* $^{215}\text{Ra}^p$	E : 3756.6(0.4) + x ; x=50(50) estimated by NUBASE **								
* $^{215}\text{Ac}^n$	E : 2438 + x ; x=50(50) from ENSDF'2001 **								
* $^{215}\text{Th}$	T : also 07Le14=0.63(+1.26-0.21) **								
* $^{215}\text{Th}^m$	E : 1421.3(0.3) + x ; x=50(50) estimated by NUBASE **								
* $^{215}\text{U}$	T : symmetrized from 15Ya13=0.73(+1.33-0.29) ms **								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>216</sup> Hg	19860#	400#	100# ms	(>300 ns)	0 <sup>+</sup>	11 10Al24 I	2010	$\beta^-$ ?; $\beta^-$ -n=6#	
<sup>216</sup> Tl	14720#	300#	6 s	3	5 <sup>+</sup> #	11 16Ca25 TD	2010	$\beta^-$ =100; $\beta^-$ -n<11.5	
<sup>216</sup> Pb	7480#	200#	1.65 m	0.2	0 <sup>+</sup>	15 16Ca25 TD	2010	$\beta^-$ =100	
<sup>216</sup> Pb <sup>m</sup>	8990#	200#	1514 20	400 ns	40	8 <sup>+</sup> #	15	IT=100 *	
<sup>216</sup> Bi	5874	11	2.25 m	0.05	(6 <sup>-</sup> , 7 <sup>-</sup> )	07	1989	$\beta^-$ ≈100	
<sup>216</sup> Bi <sup>m</sup>	5898	15	24 19	6.6 m	2.1	(3) <sup>(-#)</sup>	07	$\beta^-$ ≈100	
<sup>216</sup> Po	1782.4	1.8	145 ms	2	0 <sup>+</sup>	07	1910	$\alpha$ =100; 2 $\beta^-$ ?	
<sup>216</sup> At	2257	4	300 $\mu$ s	30	1 <sup>(-)</sup>	07	1948	$\alpha$ ≈100; $\beta^-$ <0.006; $\epsilon$ <3e-7	
<sup>216</sup> At <sup>m</sup>	2417	10	161 11	100# $\mu$ s	9 <sup>-</sup> #	07	1971	$\alpha$ =100	
<sup>216</sup> Rn	253	6	45 $\mu$ s	5	0 <sup>+</sup>	07	1949	$\alpha$ =100	
<sup>216</sup> Fr	2971	4	700 ns	20	(1 <sup>-</sup> )	07	1970	$\alpha$ =100; $\beta^+$ <2e-7#	
<sup>216</sup> Fr <sup>m</sup>	3190	6	219 6	850 ns	30	(9 <sup>-</sup> )	07	$\alpha$ =?; $\beta^+$ ?	
<sup>216</sup> Ra	3291	9	182 ns	10	0 <sup>+</sup>	07	1972	$\alpha$ =100; $\epsilon$ <1e-8	
<sup>216</sup> Ac	8144	11	440 $\mu$ s	16	(1 <sup>-</sup> )	07	1967	$\alpha$ =100; $\beta^+$ =7e-5#	
<sup>216</sup> Ac <sup>m</sup>	8188	10	44 8	441 $\mu$ s	7	(9 <sup>-</sup> )	07	$\alpha$ =100; $\beta^+$ =7e-5#	
<sup>216</sup> Ac <sup>n</sup>	8560#	100#	420# 100#	300 ns	ns	07	2006	IT=100 *	
<sup>216</sup> Th	10298	12	26.0 ms	0.2	0 <sup>+</sup>	07	1968	$\alpha$ ≈100; $\beta^+$ =0.01#	
<sup>216</sup> Th <sup>m</sup>	12342	14	2043 9	134 $\mu$ s	4	(8 <sup>+</sup> )	07	IT ?; $\alpha$ =2.8 9	
<sup>216</sup> Th <sup>n</sup>	12945	12	2646.8 0.1	580 ns	30	(11 <sup>-</sup> )	07	IT=100	
<sup>216</sup> Th <sup>p</sup>	13979	12	3681.4 0.7	740 ns	70	(14 <sup>+</sup> )	07	IT=100	
<sup>216</sup> Pa	17800	50	105 ms	12	0 <sup>+</sup>	07	96An21 T	1972	$\alpha$ =?; $\beta^+$ =2# *
<sup>216</sup> U	23066	28	6.9 ms	2.9	0 <sup>+</sup>	15 15Ma37 T	2015	$\alpha$ =100 *	
<sup>216</sup> U <sup>m</sup>	25320	30	2250 40	1.4 ms	0.9	8 <sup>+</sup> #	15 15Ma37 T	2015	$\alpha$ =100 *
* <sup>216</sup> Pb <sup>m</sup>	E : 1459 + x ; x=20-90 keV **								
* <sup>216</sup> Ac <sup>n</sup>	E : 322 + x, x=100#100 **								
* <sup>216</sup> Pa	T : others 98Ik01=150(70-40), 140(50-30) 79Sc09=170(100-40) 71Su14=200(40) **								
* <sup>216</sup> U	T : average 15Ma37=4.72(+4.72-1.57) 15De22=3.8(+8.8-3.2) **								
* <sup>216</sup> U <sup>m</sup>	T : symmetrized from 15Ma37=0.74(+1.34-0.29) **								
<sup>217</sup> Tl	18310#	400#	1# s	(>300 ns)	1/2 <sup>+</sup> #	11 10Al24 I	2010	$\beta^-$ ?; $\beta^-$ -n=100#	
<sup>217</sup> Pb	12240#	300#	20 s	5	9/2 <sup>+</sup> #	11 16Ca25 TD	2010	$\beta^-$ =100	
<sup>217</sup> Bi	8730	18	98.5 s	1.3	9/2 <sup>-</sup> #	14	1998	$\beta^-$ =100	
<sup>217</sup> Bi <sup>m</sup>	10210	40	1480 40	2.70 $\mu$ s	0.06	25/2 <sup>-</sup> #	14 14Mo02 T	2012	IT=100 *
<sup>217</sup> Po	5884	7	1.514 s	0.026	(9/2 <sup>+</sup> )	03 04Li28 TJ	1956	$\alpha$ >95; $\beta^-$ <5 *	
<sup>217</sup> At	4395	5	32.62 ms	0.24	9/2 <sup>-</sup>	03 13Su13 T	1947	$\alpha$ ≈100; $\beta^-$ =0.008 2 *	
<sup>217</sup> Rn	3659	4	540 $\mu$ s	50	9/2 <sup>+</sup>	03	1949	$\alpha$ =100	
<sup>217</sup> Fr	4315	7	16.8 $\mu$ s	1.9	9/2 <sup>-</sup>	03 90An19 T	1968	$\alpha$ =100 *	
<sup>217</sup> Ra	5890	7	1.63 $\mu$ s	0.17	(9/2 <sup>+</sup> )	03 90An19 T	1970	$\alpha$ =100 *	
<sup>217</sup> Ac	8704	11	69 ns	4	9/2 <sup>-</sup>	03	1972	$\alpha$ =100; $\beta^+$ =6.9e-9	
<sup>217</sup> Ac <sup>m</sup>	10716	18	2012 20	740 ns	40	(29/2 <sup>+</sup> )	03	IT=95.7 10; $\alpha$ =4.3 10	
<sup>217</sup> Th	12206	11	247 $\mu$ s	4	9/2 <sup>+</sup> #	03 05Ku31 T	1968	$\alpha$ =100 *	
<sup>217</sup> Th <sup>m</sup>	12880	11	673.8 1.8	141 ns	50	(15/2 <sup>-</sup> )	03	1989	IT=100 *
<sup>217</sup> Th <sup>n</sup>	14510#	60#	2307# 55#	71 $\mu$ s	14	(25/2 <sup>+</sup> )	03 05Ku31 ETJ	2002	IT=100 *
<sup>217</sup> Pa	17068	16	3.48 ms	0.09	9/2 <sup>-</sup> #	03 02He29 T	1968	$\alpha$ =100; B=0.0024# *	
<sup>217</sup> Pa <sup>m</sup>	18929	16	1860 7	1.08 ms	0.03	(23/2 <sup>-</sup> )	03 02He29 TD	1979	$\alpha$ =73 4; IT ? *
<sup>217</sup> U	22970#	70#	800 $\mu$ s	700	1/2 <sup>-</sup> #	03 05Le42 T	2000	$\alpha$ ≈100; $\beta^-$ =0.05# *	
* <sup>217</sup> Bi <sup>m</sup>	E : 1436 + y ; y=40(40) estimated by NUBASE **								
* <sup>217</sup> Po	T : average 03Ku25=1.53(0.03) 96Ry.B=1.47(0.05); other 04Li28=1.6(0.2) **								
* <sup>217</sup> At	T : average 13Su13=32.8(0.3) 63Di05=32.3(0.4) **								
* <sup>217</sup> At	D : average $\beta^-$ 97Ch53=0.0067(24)% 69Le.A=0.012(4)% **								
* <sup>217</sup> Fr	T : average 90An19=16(2) 70Bo13=22(5) **								
* <sup>217</sup> Ra	T : average 90An19=1.7(0.3) 70Bo13=1.6(0.2) **								
* <sup>217</sup> Th	T : unweighed average 05Ku31=257(2) 02He29=237(2) 00He17=247(3) 73Ha32=252(7) **								
* <sup>217</sup> Th	T : others 15Kh09=259(12) 05Li17=310(70) **								
* <sup>217</sup> Th <sup>m</sup>	E : uncertainty estimated by NUBASE **								
* <sup>217</sup> Th <sup>n</sup>	T : symmetrized from 05Ku31=67(+17-11); other 02Mu.A=20(5) **								
* <sup>217</sup> Th <sup>n</sup>	E : weak Kx rays placed it less than 110 keV above 21/2 <sup>+</sup> at 2252 keV **								
* <sup>217</sup> Pa	T : average 02He29=3.8(0.2) 00He17=3.4(0.1) **								
* <sup>217</sup> Pa <sup>m</sup>	J : from 13As01 **								
* <sup>217</sup> U	T : symmetrized from 0.19(+1.13-0.10) ms; other 00Ma65=15.6(+21.3-5.7) ms **								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{218}\text{Tl}$	23180#	400#	200# ms	$5^+\#$				$\beta^-$ ?; $\beta^-_{n=70}\#$
$^{218}\text{Pb}$	15450#	300#	15 s 7	$0^+$	11	16Ca25 TD	2009	$\beta^-_{n=100}$
$^{218}\text{Bi}$	13216	27	33 s 1	$(6^-, 7^-, 8^-)$	06	04De16 J	1998	$\beta^-_{n=100}$
$^{218}\text{Po}$	8356.9	2.0	3.098 m 0.012	$0^+$	06		1904	$\alpha \approx 100$ ; $\beta^- = 0.02\#$
$^{218}\text{At}$	8098	12	1.5 s 0.3	$1^- \#$	06		1943	$\alpha \approx 100$ ; $\beta^- = 0.1\#$
$^{218}\text{Rn}$	5217.3	2.3	33.75 ms 0.15	$0^+$	06	12Su11 T	1948	$\alpha = 100$
$^{218}\text{Fr}$	7059	5	1.0 ms 0.6	$1^-$	06		1949	$\alpha = 100$
$^{218}\text{Fr}^m$	7146	6	86 4 AD	$(8^-)$	06	99Sh03 J	1982	$\alpha \approx 100$ ; IT ?
$^{218}\text{Fr}^p$	7260#	150#	200# 150#	high				
$^{218}\text{Ra}$	6651	11	25.2 $\mu\text{s}$ 0.3	$0^+$	06		1970	$\alpha = 100$ ; $2\beta^+$ ?
$^{218}\text{Ac}$	10840	50	1.00 $\mu\text{s}$ 0.04	$1^- \#$	06	15Kh09 T	1970	$\alpha = 100$
$^{218}\text{Ac}^m$	10990#	70#	150# 50#	32 ns 9		94De04 ET	1994	*
$^{218}\text{Ac}^n$	11370#	70#	530# 50#	103 ns 11		06	1994	IT=100
$^{218}\text{Th}$	12367	11	117 ns 9	$0^+$	06		1973	$\alpha = 100$
$^{218}\text{Pa}$	18684	18	113 $\mu\text{s}$ 10		06		1979	$\alpha = 100$
$^{218}\text{U}$	21895	14	550 $\mu\text{s}$ 140	$0^+$	06		1992	$\alpha = 100$
$^{218}\text{U}^m$	24004	18	2109 17 AD	$(8^+)$	06	15Ma37 T	2005	$\alpha = 100$
* $^{218}\text{Ac}$	T : average 15Kh09=0.96(0.05) 89Mi17=1.06(0.09) 83Sc23=1.12(0.11)							
* $^{218}\text{Ac}^m$	E : at least 122.5 in 94De04							
* $^{218}\text{Ac}^n$	E : 384.49(0.13) keV above $^{218}\text{Ac}^m$ , from ENSDF							
* $^{218}\text{Th}$	T : also 15Kh09=160(40)							
* $^{218}\text{U}$	T : symmetrized from 510(+170-100)							
* $^{218}\text{U}^m$	T : average 15Ma37=280(+1300-120) 05Le42=560(+260-140)							
$^{219}\text{Pb}$	20280#	400#	10# s (>300 ns)	$9/2^+\#$	11	10A124 I	2009	$\beta^-$ ?
$^{219}\text{Bi}$	16280#	200#	8.7 s 2.9	$9/2^- \#$	12	16Ca25 T	2009	$\beta^-_{n=100}$
$^{219}\text{Po}$	12681	16	10.3 m 1.0	$9/2^+\#$	15	15Fi07 T	1998	$\beta^-$ ?; $\alpha = 28.2$ 20
$^{219}\text{At}$	10396	3	56 s 3	$(9/2^-)$	16		1953	$\alpha = 93.6$ 10; $\beta^- = ?$
$^{219}\text{Rn}$	8829.4	2.1	3.96 s 0.01	$5/2^+$	01		1903	$\alpha = 100$
$^{219}\text{Fr}$	8618	7	20 ms 2	$9/2^-$	01		1948	$\alpha = 100$
$^{219}\text{Ra}$	9394	8	10 ms 3	$(7/2^+)$	01		1952	$\alpha = 100$
$^{219}\text{Ac}$	11570	50	11.8 $\mu\text{s}$ 1.5	$9/2^-$	01		1970	$\alpha = 100$ ; $\beta^+ = 1e-6\#$
$^{219}\text{Th}$	14470	50	1.021 $\mu\text{s}$ 0.024	$9/2^+\#$	12	15Kh09 T	1973	$\alpha = 100$ ; $\beta^+ = 1e-7\#$
$^{219}\text{Pa}$	18540	50	53 ns 10	$9/2^-$	01		2005	$\alpha = 100$ ; $\beta^+ = 5e-9\#$
$^{219}\text{U}$	23290	50	55 $\mu\text{s}$ 25	$9/2^+\#$	01		1993	$\alpha = 100$ ; $\beta^+ = 1.4e-5\#$
$^{219}\text{Np}$	29460	90	< 5 $\mu\text{s}$	$9/2^- \#$	16	15De22 D		$\alpha = 100$
* $^{219}\text{Bi}$	T : other 12Be28t=22(7)							
* $^{219}\text{Po}$	T : from 15Fi07=620(59) s							
* $^{219}\text{Th}$	T : 15Kh09=0.97(0.04) 73Ha32=1.05(0.03)							
* $^{219}\text{U}$	T : symmetrized from 42(+34-13); also 05Le42=80(+100-30)							
$^{220}\text{Pb}$	23670#	400#	30# s (>300 ns)	$0^+$	11	10A124 I	2010	$\beta^-$ ?
$^{220}\text{Bi}$	20820#	300#	9.5 s 5.7	$1^- \#$	11	16Ca25 TD	2010	$\beta^-_{n=100}$ ; $\beta^-_{n=0.04}\#$
$^{220}\text{Po}$	15263	18	40# s (>300 ns)	$0^+$	11	98Pf02 I	1998	$\beta^-$ ?
$^{220}\text{At}$	14376	14	3.71 m 0.04	$3^{\text{(-#)}}$	11		1989	$\beta^-_{n=92} 2$ ; $\alpha = 8$ 2
$^{220}\text{Rn}$	10612.1	1.8	55.6 s 0.1	$0^+$	11		1900	$\alpha = 100$ ; $2\beta^-$ ?
$^{220}\text{Fr}$	11482	4	27.4 s 0.3	$1^+$	11	78Ek02 J	1948	$\alpha \approx 100$ ; $\beta^- = 0.35$ 5
$^{220}\text{Ra}$	10270	8	17.9 ms 1.4	$0^+$	11	00He17 T	1949	$\alpha = 100$
$^{220}\text{Ac}$	13744	6	26.36 ms 0.19	$(3^-)$	11	90An19 T	1970	$\alpha = 100$ ; $\beta^+ = 5e-4\#$
$^{220}\text{Th}$	14669	22	9.7 $\mu\text{s}$ 0.6	$0^+$	11		1973	$\alpha = 100$ ; $\epsilon = 2e-7\#$
$^{220}\text{Pa}$	20220#	50#	780 ns 160	$1^- \#$	11		2005	$\alpha = 100$ ; $\beta^+ = 3e-7\#$
$^{220}\text{U}$	22930#	100#	60# ns	$0^+$				$\alpha$ ?; $\beta^+$ ?
$^{220}\text{Np}$	30310#	200#	30# ns	$1^- \#$				$\alpha$ ?
* $^{220}\text{Ra}$	T : average 00He17=18(2) 90An19=17(2) 61Ru06=23(5)							
* $^{220}\text{Ac}$	T : average 90An19=26.4(0.2) 70Bo13=26.1(0.5)							
$^{221}\text{Bi}$	24100#	300#	5# s (>300 ns)	$9/2^- \#$	11	10A124 I	2009	$\beta^-$ ?; $\beta^-_{n=2}\#$
$^{221}\text{Po}$	19774	20	2.2 m 0.7	$9/2^+\#$	13		2010	$\beta^-$ ?
$^{221}\text{At}$	16783	14	2.3 m 0.2	$3/2^- \#$	07		1989	$\beta^-_{n=100}$
$^{221}\text{Rn}$	14471	6	25.7 m 0.5	$7/2^+$	07	97Li23 T	1956	$\beta^-_{n=78} 1$ ; $\alpha = 22$ 1
$^{221}\text{Fr}$	13277	5	4.801 m 0.005	$5/2^-$	07	13Su13 T	1947	$\alpha \approx 100$ ; $\beta^- = 0.0048$ 15; 14C=8.8e-11 11
$^{221}\text{Ra}$	12964	5	28 s 2	$5/2^+$	07	94Bo28 D	1949	$\alpha = 100$ ; 14C=1.2e-10 9
$^{221}\text{Ac}$	14520	50	52 ms 2	$9/2^- \#$	07		1968	$\alpha = 100$
$^{221}\text{Th}$	16940	8	1.78 ms 0.03	$7/2^+\#$	07	14Lo10 T	1970	$\alpha = 100$
$^{221}\text{Pa}$	20380	50	5.9 $\mu\text{s}$ 1.7	$9/2^-$	07		1983	$\alpha = 100$

... A-group is continued on next page ...



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
... A-group continued ...										
<sup>221</sup> U	24520	50	660 ns	140	(9/2 <sup>+</sup> )	15	2015	$\alpha \approx 100$ ; $\beta^+$ ?		
<sup>221</sup> Np	29850#	200#	30#	ns	9/2 <sup>-</sup> #			$\alpha$ ?		
* <sup>221</sup> Po	T : symmetrized from 10Ch19=112(+58-28) s									
* <sup>221</sup> Fr	D : $\beta^-$ intensity is from 97Ch53; <sup>14</sup> C intensity is from 94Bo28									
* <sup>221</sup> Fr	T : average 13Su13=4.806(0.006) 10Wa42=4.768(0.017) 07Je07=4.79(0.02)									
<sup>222</sup> Bi	28730#	300#	2#	s (>300 ns)	1 <sup>-</sup> #	10Al24	I	2009	$\beta^-$ ?; $\beta^- n=1\#$	
<sup>222</sup> Po	22490	40	9.1 m	7.2	0 <sup>+</sup>	11		2010	$\beta^-$ ?	
<sup>222</sup> At	20953	16	54 s	10		11		1989	$\beta^- =100$	
<sup>222</sup> Rn	16372.2	1.9	3.8215 d	0.0002	0 <sup>+</sup>	11	15Be07	T	1899	$\alpha=100$
<sup>222</sup> Fr	16378	7	14.2 m	0.3	2 <sup>-</sup>	11	78Ek02	J	1975	$\beta^- =100$
<sup>222</sup> Ra	14320	4	33.6 s	0.4	0 <sup>+</sup>	11	12Po13	T	1948	$\alpha=100$ ; 14C=3.0e-8 10
<sup>222</sup> Ac	16621	5	5.0 s	0.5	1 <sup>-</sup>	11		1949	$\alpha=99$ 1; $\beta^+=1$ 1	
<sup>222</sup> Ac <sup>m</sup>	16820#	150#	200#	150#	1.05 m	0.05	high	11	1972	$\alpha=?$ ; IT<10; $\beta^+=1.4$ 4
<sup>222</sup> Th	17203	12	2.24 ms	0.03	0 <sup>+</sup>	11		1970	$\alpha=100$ ; $\epsilon < 1.3e-8\#$	
<sup>222</sup> Pa	22160#	70#	3.2 ms	0.3		11	95Ni.A	T	1970	$\alpha=100$
<sup>222</sup> U	24270	50	4.7 $\mu$ s	0.7	0 <sup>+</sup>	15		1983	$\alpha=100$ ; $\beta^+ < 1e-6\#$	
<sup>222</sup> Np	31020#	200#	700#	ns	1 <sup>-</sup> #				$\alpha$ ?	
* <sup>222</sup> Po	T : symmetrized from 10Ch19=145(+694-66) s									
* <sup>222</sup> Rn	T : rounded from 15Be07=3.82146(16stat,4syst)									
* <sup>222</sup> Ra	T : others not used 95Ko54=36.17(0.10) 82Bo04=43(4)									
* <sup>222</sup> Ac <sup>m</sup>	D : derived from 0.7% < $\beta^+$ < 2%, in ENSDF									
* <sup>222</sup> Pa	T : average 95Ni.A=3.3(0.3) 79Sc09=2.9(+0.6-0.4)									
* <sup>222</sup> Pa	T : 70Bo13=5.7(0.5) conflicting, not used									
<sup>223</sup> Bi	32140#	400#	1#	s (>300 ns)	9/2 <sup>-</sup> #	11	10Al24	I	2009	$\beta^-$ ?; $\beta^- n=5\#$
<sup>223</sup> Po	27080#	200#	1#	m (>300 ns)	9/2 <sup>+</sup> #	11	10Al24	I	2010	$\beta^-$ ?
<sup>223</sup> At	23428	14	50 s	7	3/2 <sup>-</sup> #	01		1989	$\beta^- \approx 100$ ; $\alpha=0.008\#$	
<sup>223</sup> Rn	20390	8	24.3 m	0.4	7/2 <sup>(+)</sup> #	01		1964	$\beta^- =100$ ; $\alpha=0.0004\#$	
<sup>223</sup> Fr	18382.4	1.9	22.00 m	0.07	3/2 <sup>-</sup>	01	85Co24	J	1939	$\beta^- \approx 100$ ; $\alpha=0.006$
<sup>223</sup> Ra	17233.3	2.1	11.4377 d	0.0022	3/2 <sup>+</sup>	01	15Ko06	T	1905	$\alpha=100$ ; 14C=8.9e-8 4
<sup>223</sup> Ac	17826	7	2.10 m	0.05	(5/2 <sup>-</sup> )	01		1948	$\alpha=99$ ; $\epsilon=1$	
<sup>223</sup> Th	19386	9	600 ms	20	(5/2 <sup>+</sup> )	01		1952	$\alpha=100$	
<sup>223</sup> Pa	22320	70	5.1 ms	0.3	9/2 <sup>-</sup> #	01	99Ho28	T	1970	$\alpha=100$ ; $\beta^+ < 0.001\#$
<sup>223</sup> U	25840	70	21 $\mu$ s	8	7/2 <sup>+</sup> #	01		1991	$\alpha \approx 100$ ; $\beta^+ =0.2\#$	
<sup>223</sup> Np	30600#	200#	1#	$\mu$ s	9/2 <sup>-</sup> #				$\alpha$ ?	
* <sup>223</sup> Ra	T : average 15Ko06=11.4362(0.0050) 15Be13=11.447(0.007) 15Be13=11.445(0.013)									
* <sup>223</sup> Ra	T : 15Co02=11.4358(0.0028)									
* <sup>223</sup> Pa	T : average 99Ho28=4.9(0.4) 95Ni.A=5.0(1.0) 70Bo13=6.5(1.0)									
* <sup>223</sup> U	T : symmetrized from 18(+10-5)									
<sup>224</sup> Bi	36830#	400#	300#	ms (>300 ns)	1 <sup>-</sup> #	15	10Al24	I	2010	$\beta^-$ ?; $\beta^- n=10\#$
<sup>224</sup> Po	29910#	200#	1#	m (>300 ns)	0 <sup>+</sup>	15	10Al24	I	2010	$\beta^-$ ?
<sup>224</sup> At	27711	22	2.5 m	1.5		15		2010	$\beta^-$ ?	
<sup>224</sup> Rn	22445	10	107 m	3	0 <sup>+</sup>	15		1964	$\beta^- =100$	
<sup>224</sup> Fr	21749	11	3.33 m	0.10	1 <sup>-</sup>	15	85Co24	J	1969	$\beta^- =100$
<sup>224</sup> Fr <sup>x</sup>	21850#	100#	100#	100#	MD					
<sup>224</sup> Ra	18825.9	1.8	3.6319 d	0.0023	0 <sup>+</sup>	15		1902	$\alpha=100$ ; 14C=4.0e-9 12	
<sup>224</sup> Ac	20234	4	2.78 h	0.16	(0 <sup>-</sup> )	15		1948	$\beta^+ =90.6$ 17; $\alpha=9.4$ 17; $\beta^- < 1.6\#$	
<sup>224</sup> Th	19994	10	1.04 s	0.02	0 <sup>+</sup>	15		1949	$\alpha=100$ ; 2 $\beta^+$ ?	
<sup>224</sup> Pa	23862	8	846 ms	20	5 <sup>-</sup> #	15		1958	$\alpha \approx 100$ ; $\beta^+ =0.1\#$	
<sup>224</sup> U	25722	23	396 $\mu$ s	17	0 <sup>+</sup>	15	14Lo10	T	1991	$\alpha=100$ ; $\beta^+ < 1.2e-4\#$
<sup>224</sup> Np	31880#	200#	100#	$\mu$ s	1 <sup>-</sup> #				$\alpha$ ?	
* <sup>224</sup> At	T : symmetrized from 10Ch19=76(+138-23) s									
* <sup>224</sup> Ac	D : symmetrized from 51Me10 $\beta^+ =90.9(+1.4-2.0)\%$ ; $\alpha=9.1(+2.0-1.4)\%$									
<sup>225</sup> Po	34530#	300#	20#	s (>300 ns)	9/2 <sup>+</sup> #	11	10Al24	I	2010	$\beta^-$ ?
<sup>225</sup> At	30400#	300#	2#	m (>300 ns)	1/2 <sup>+</sup> #	11	10Al24	I	2010	$\beta^-$ ?
<sup>225</sup> Rn	26534	11	4.66 m	0.04	7/2 <sup>-</sup>	09		1969	$\beta^- =100$	
<sup>225</sup> Fr	23821	12	3.95 m	0.14	3/2 <sup>-</sup>	09	85Co24	J	1969	$\beta^- =100$
<sup>225</sup> Ra	21993.1	2.6	14.9 d	0.2	1/2 <sup>+</sup>	09		1947	$\beta^- =100$	
<sup>225</sup> Ac	21637	5	9.920 d	0.003	3/2 <sup>-</sup> #	09	12Po14	T	1947	$\alpha=100$ ; 14C=4.5e-12 14
<sup>225</sup> Th	22310	5	8.75 m	0.04	(3/2 <sup>+</sup> )	09		1949	$\alpha \approx 90$ ; $\epsilon \approx 10$	
<sup>225</sup> Pa	24340	70	1.7 s	0.2	5/2 <sup>-</sup> #	09		1958	$\alpha=100$	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
... A-group continued ...											
<sup>225</sup> U	27380	11	61 ms	4		5/2 <sup>+</sup> #	09	00He17 T	1989	$\alpha=100$	*
<sup>225</sup> Np	31590	70	6 ms	5		9/2 <sup>-</sup> #	09	15De22 T	1994	$\alpha=100; \beta^+ ?$	*
* <sup>225</sup> U	T : symmetrized from 00He17=59(+5-2); others not used 03Ni10=135(+93-39)									**	
* <sup>225</sup> U	T : 01Ku07=84(4) 94An02=68(+45-20) 92To02=95(15) and 89He13=80(+40-10)									**	
* <sup>225</sup> Np	T : symmetrized from 15De22=3.3(+7.6-2.7); also 15De22=3.8(+7.6-2.7)									**	
<sup>226</sup> Po	37550#	400#	20#	s	(>300 ns)	0 <sup>+</sup>	11	10Al24 I	2010	$\beta^- ?$	
<sup>226</sup> At	34610#	300#	20#	s	(>300 ns)		11	10Al24 I	2010	$\beta^- ?; \beta^- n=0\#$	
<sup>226</sup> Rn	28747	10	7.4	m	0.1	0 <sup>+</sup>	96		1969	$\beta^- =100$	
<sup>226</sup> Fr	27521	6	49	s	1	1 <sup>-</sup>	96	85Co24 J	1969	$\beta^- =100$	
<sup>226</sup> Ra	23667.8	1.9	1.600	ky	0.007	0 <sup>+</sup>	96	90We01 D	1898	$\alpha=100; 14C=2.6e-9 6; 2\beta^- ?$	*
<sup>226</sup> Ac	24309	3	29.37	h	0.12	(1) <sup>(-#)</sup>	96		1950	$\beta^- =83 3; \epsilon=17 3; \alpha=0.006 2$	
<sup>226</sup> Th	23198	4	30.70	m	0.03	0 <sup>+</sup>	96	01Bo11 D	1948	$\alpha=100; ^{18}O<3.2e-12$	*
<sup>226</sup> Pa	26033	11	1.8	m	0.2		96		1949	$\alpha=74 5; \beta^+=26 5$	
<sup>226</sup> U	27329	13	269	ms	6	0 <sup>+</sup>	14	01Ca.B T	1973	$\alpha=100$	*
<sup>226</sup> Np	32780#	90#	35	ms	10		96		1990	$\alpha=100; \beta^+=0.003\#$	
* <sup>226</sup> Ra	D : <sup>14</sup> C : average 90We01=2.3(0.8)e-9% 86Ba26=2.9(1.0)e-9% 85Ho21=3.2(1.6)e-9%									**	
* <sup>226</sup> Th	T : from 12Po13; other 87Mi10=30.57(0.10)									**	
* <sup>226</sup> U	T : average 01Ca.B=258(13) 00He17=281(9) 99Gr28=260(10)									**	
<sup>227</sup> Po	42280#	400#	5#	s	(>300 ns)	9/2 <sup>+</sup> #	16		2010	$\beta^- ?$	
<sup>227</sup> At	37480#	300#	20#	s	(>300 ns)	1/2 <sup>+</sup> #	16		2010	$\beta^- ?; \beta^- n=0.2\#$	
<sup>227</sup> Rn	32886	14	20.2	s	0.4	(5/2) <sup>(+#)</sup>	16		1986	$\beta^- =100$	
<sup>227</sup> Fr	29682	6	2.47	m	0.03	1/2 <sup>+</sup>	16	85Co24 J	1972	$\beta^- =100$	*
<sup>227</sup> Ra	27177.7	2.0	42.2	m	0.5	3/2 <sup>+</sup>	16		1953	$\beta^- =100$	
<sup>227</sup> Ac	25849.6	1.9	21.772	y	0.003	3/2 <sup>-</sup>	16		1902	$\beta^- =98.62 36; \alpha=1.38 36$	
<sup>227</sup> Th	25804.8	2.1	18.697	d	0.007	(1/2 <sup>+</sup> )	16		1906	$\alpha=100$	
<sup>227</sup> Pa	26831	7	38.3	m	0.3	(5/2 <sup>-</sup> )	16		1948	$\alpha=85 2; \epsilon=15 2$	
<sup>227</sup> U	29045	10	1.1	m	0.1	(3/2 <sup>+</sup> )	16		1952	$\alpha=100; \beta^+ <0.001\#$	
<sup>227</sup> Np	32560	70	510	ms	60	5/2 <sup>-</sup> #	16		1990	$\alpha \approx 100; \beta^+ =0.05\#$	
<sup>227</sup> Pu	36770#	100#	20#	ms		5/2 <sup>+</sup> #				$\alpha ?$	
<sup>228</sup> At	41680#	400#	5#	s	(>300 ns)		14	10Al24 I	2010	$\beta^- ?; \beta^- n=0.6\#$	
<sup>228</sup> Rn	35243	18	65	s	2	0 <sup>+</sup>	14		1989	$\beta^- =100$	
<sup>228</sup> Fr	33384	7	38	s	1	2 <sup>-</sup>	14	85Co24 J	1972	$\beta^- =100$	*
<sup>228</sup> Ra	28940.3	2.0	5.75	y	0.03	0 <sup>+</sup>	14		1907	$\beta^- =100$	
<sup>228</sup> Ac	28894.7	2.1	6.15	h	0.02	3 <sup>+</sup>	14		1908	$\beta^- =100$	*
<sup>228</sup> Th	26771.0	1.8	1.9124	y	0.0008	0 <sup>+</sup>	14	93Bo20 D	1905	$\alpha=100; ^{20}O=1.13e-11 22$	*
<sup>228</sup> Pa	28924	4	22	h	1	3 <sup>+</sup>	14		1948	$\beta^+=98.15 17; \alpha=1.85 17$	
<sup>228</sup> U	29222	14	9.1	m	0.2	0 <sup>+</sup>	14		1949	$\alpha>95; \epsilon<5$	
<sup>228</sup> Np	33600	50	61.4	s	1.4		14	94Kr13 D	1994	$\epsilon=60 7; \alpha=40 7; \beta^+ SF=0.012 6$	*
<sup>228</sup> Pu	36087	29	2.1	s	1.3	0 <sup>+</sup>	14	03Ni10 T	1994	$\alpha \approx 100; \beta^+ <7\#$	*
* <sup>228</sup> Fr	I : 08Ch.A reports an excited isomer with half-life=94(+170-29) s									**	
* <sup>228</sup> Ac	I : 08Ch.A reports an excited isomer with half-life=149(+95-42) s									**	
* <sup>228</sup> Th	T : average 14Un01=698.3(0.6) 71Jo14=698.77(0.32) 56Ki16=697.6(0.7)									**	
* <sup>228</sup> Np	D : $\beta^+ SF=0.020(9)\%$ defined in 94Kr13 relative to $\epsilon$ , thus 0.012(6)% of total									**	
* <sup>228</sup> Pu	T : symmetrized from 03Ni10=1.1(+2.0-0.5)									**	
<sup>229</sup> At	44820#	400#	5#	s	(>300 ns)	1/2 <sup>+</sup> #	11	10Al24 I	2010	$\beta^- ?; \beta^- n=4\#$	
<sup>229</sup> Rn	39362	13	11.9	s	1.3	5/2 <sup>+</sup> #	09		2009	$\beta^- ?$	*
<sup>229</sup> Fr	35668	5	50.2	s	0.4	(1/2 <sup>+</sup> )	08	14Bu06 J	1975	$\beta^- =100$	*
<sup>229</sup> Ra	32562	15	4.0	m	0.2	5/2 <sup>+</sup>	08		1975	$\beta^- =100$	
<sup>229</sup> Ac	30690	12	62.7	m	0.5	(3/2 <sup>+</sup> )	08		1952	$\beta^- =100$	
<sup>229</sup> Th	29585.6	2.4	7.920	ky	0.017	5/2 <sup>+</sup>	08	14Va04 T	1947	$\alpha=100$	*
<sup>229</sup> Th <sup>m</sup>	29585.6	2.4	< 1	s		(3/2 <sup>+</sup> )	08	16We07 ETD	1994	$\Gamma T=100; \alpha ?$	*
<sup>229</sup> Pa	29897	3	1.50	d	0.05	(5/2 <sup>+</sup> )	08		1949	$\epsilon \approx 100; \alpha=0.48 5$	
<sup>229</sup> Pa <sup>m</sup>	29909	3	12.20	ns	0.04	3/2 <sup>-</sup>	08	15Ah04 EJD	1982	$\epsilon ?; \Gamma T ?$	*
<sup>229</sup> U	31211	6	57.8	m	0.5	(3/2 <sup>+</sup> )	08	15Ah04 T	1949	$\beta^+ \approx 80; \alpha \approx 20$	
<sup>229</sup> Np	33780	90	4.00	m	0.18	5/2 <sup>+</sup> #	08	04Sa05 TD	1968	$\alpha=68 11; \beta^+ ?$	*
<sup>229</sup> Np <sup>p</sup>	33940#	100#	160#	50#		5/2 <sup>-</sup> #					
<sup>229</sup> Pu	37400	50	91	s	26	3/2 <sup>+</sup> #	08	10Kh06 TD	1994	$\alpha=50 20; \beta^+=50 20; SF<7$	*
... A-group is continued on next page ...											

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>229</sup> Am	42150 90		1.8 s	1.5	5/2 <sup>-</sup> #	15	2015	$\alpha \approx 91; \beta^+ ?$
<sup>229</sup> Am <sup>p</sup>	42530#	220# 380# 200#						IT ?
* <sup>229</sup> Rn	T : symmetrized from 09Ne03=12.0(+1.2-1.3)							
* <sup>229</sup> Fr	T : 92Bo05=50.2(0.4); ENSDF2008 50.2 S 20 is misprint							
* <sup>229</sup> Th	T : as evaluated by 14Va04							
* <sup>229</sup> Th <sup>m</sup>	T : >60 s for 2 <sup>+</sup> charge state from 16We07; others 09In01(1m<T<3m); 09Ki14<2h							
* <sup>229</sup> Th <sup>m</sup>	E : 0.0063<Ex<0.0183 from 16We07; other 94He08=0.0035(0.0010)							
* <sup>229</sup> Pa <sup>m</sup>	D : 98Le15 IT=100 rejected by 15Ah04							
* <sup>229</sup> Np	T : average 04Sa05=4.0(0.4) 68Ha14=4.0(0.2)							
* <sup>229</sup> Pu	T : average 10Kh06=67(+41-19) 01Ca.B=90(+71-27)							
* <sup>229</sup> Pu	D : from ENSDF'97							
* <sup>229</sup> Am	T : symmetrized from 15De22=0.9(+2.1-0.7); also 15De22=6.4(+14.9-5.4)							
<sup>230</sup> Rn	42050#	200#	10#	s	(>300 ns)	0 <sup>+</sup>	12 10Al24 I	2010 $\beta^- ?$
<sup>230</sup> Fr	39487	7	19.1	s	0.5		12	1987 $\beta^- =100$
<sup>230</sup> Ra	34516	10	93	m	2	0 <sup>+</sup>	12	1978 $\beta^- =100$
<sup>230</sup> Ac	33838	16	122	s	3	(1 <sup>+</sup> )	12	1973 $\beta^- =100; \beta^- SF=1.2e-6 4$
<sup>230</sup> Th	30862.6	1.2	75.4	ky	0.3	0 <sup>+</sup>	12	1907 $\alpha =100; SF<4e-12; ^{24}Ne=5.8e-11 13$
<sup>230</sup> Pa	32174	3	17.4	d	0.5	2 <sup>-</sup>	14	1948 $\beta^+ =92.2 7; \beta^- =7.8 7; \alpha =0.0032 1$
<sup>230</sup> U	31615	5	20.23	d	0.02	0 <sup>+</sup>	12 12Po12 T	1948 $\alpha =100; ^{22}Ne=4.8e-12 20; SF<1.4e-10#; \dots$
<sup>230</sup> Np	35240	50	4.6	m	0.3		12	1968 $\beta^+ <97; \alpha >3$
<sup>230</sup> Pu	36934	15	1.70	m	0.17	0 <sup>+</sup>	12 01Ca.B T	1990 $\alpha \approx 100; \beta^+ ?$
<sup>230</sup> Am	42930#	130#	40	s	16		12 16Ka13 TD	2003 $\beta^+ \approx 100; \beta^+ SF=?; SF=?$
* <sup>230</sup> U	D : ... ; 2 $\beta^+ ?$							
* <sup>230</sup> Am	T : symmetrized from 16Ka13=32(+22-9) s							
<sup>231</sup> Rn	46450#	300#	300#	ms	(>300 ns)	1/2 <sup>+</sup> #	13 10Al24 I	2010 $\beta^- ?$
<sup>231</sup> Fr	42081	8	17.6	s	0.6	(1/2 <sup>+</sup> )	13 14Bu06 J	1985 $\beta^- =100$
<sup>231</sup> Ra	38216	11	104	s	1	(5/2 <sup>+</sup> )	13 06Bo33 T	1983 $\beta^- =100$
<sup>231</sup> Ra <sup>m</sup>	38282	11	53	$\mu$ s		(1/2 <sup>+</sup> )	13	2001 IT=100
<sup>231</sup> Ac	35763	13	7.5	m	0.1	1/2 <sup>+</sup>	13	1973 $\beta^- =100$
<sup>231</sup> Th	33815.9	1.2	25.52	h	0.01	5/2 <sup>+</sup>	13	1911 $\beta^- =100; \alpha =4e-11#$
<sup>231</sup> Pa	33424.4	1.8	32.76	ky	0.11	3/2 <sup>-</sup>	13	1918 $\alpha =100; SF \leq 3e-10; ^{24}Ne=13.4e-10 17; \dots$
<sup>231</sup> U	33806.0	2.7	4.2	d	0.1	(5/2) <sup>(+)</sup>	13	1949 $\epsilon \approx 100; \alpha =0.004 1$
<sup>231</sup> Np	35620	50	48.8	m	0.2	(5/2) <sup>(+)</sup>	13	1950 $\beta^+ =98 1; \alpha =2 1$
<sup>231</sup> Pu	38309	23	8.6	m	0.5	(3/2 <sup>+</sup> )	13	1999 $\beta^+ =87 5; \alpha =13 5$
<sup>231</sup> Am	42410#	300#	1#	m		5/2 <sup>-</sup> #		$\beta^+ ?; \alpha ?$
<sup>231</sup> Cm	47270#	300#	20#	s		3/2 <sup>+</sup> #		$\beta^+ ?; \alpha ?$
* <sup>231</sup> Pa	D : ... ; <sup>23</sup> F=9.9e-13							
* <sup>231</sup> Pu	D : symmetrized from 99La14=90(+3-7)% and 10(+7-3)%							
<sup>232</sup> Fr	46073	14	5.5	s	0.6	(5)	06	1990 $\beta^- =100; \beta^- SF<2e-4$
<sup>232</sup> Ra	40497	9	4.0	m	0.3	0 <sup>+</sup>	06 08Ch.A T	1983 $\beta^- =100$
<sup>232</sup> Ac	39154	13	1.98	m	0.08	(1 <sup>+</sup> )	06	1986 $\beta^- =100$
<sup>232</sup> Th	35446.8	1.4	14.0	Gy	0.1	0 <sup>+</sup>	06	1898 IS=100; $\alpha =100; SF=1.1e-9 4; \dots$
<sup>232</sup> Pa	35947	8	1.32	d	0.02	(2 <sup>-</sup> )	06	1949 $\beta^- \approx 100; \epsilon =0.003 1$
<sup>232</sup> U	34609.5	1.8	68.9	y	0.4	0 <sup>+</sup>	06	1949 $\alpha =100; ^{24}Ne=8.9e-10 7; SF=2.7e-12 6; \dots$
<sup>232</sup> Np	37360#	100#	14.7	m	0.3	(4 <sup>+</sup> )	06	1950 $\beta^+ \approx 100; \alpha \approx 0.0002#$
<sup>232</sup> Pu	38363	18	33.7	m	0.5	0 <sup>+</sup>	06	1973 $\epsilon =90#; \alpha =11 6$
<sup>232</sup> Am	43340#	300#	1.31	m	0.04	1 <sup>-</sup> #	06	1967 $\beta^+ =?; \alpha =3#; \beta^+ SF=0.069 10$
<sup>232</sup> Cm	46310#	200#	10#	s		0 <sup>+</sup>		$\beta^+ ?; \alpha ?$
* <sup>232</sup> Ra	T : average 08Ch.A=4.00(0.33) 86Gi08=4.2(0.8)							
* <sup>232</sup> Th	D : ... ; <sup>24</sup> Ne+ <sup>26</sup> Ne<2.78e-10; 2 $\beta^- ?$							
* <sup>232</sup> U	D : ... ; <sup>28</sup> Mg<5e-12							
* <sup>232</sup> Pu	T : average 00La25=33.1(0.8) 73Ja06=34.1(0.7) D : 52Or.A $\alpha >1.6\%$ 73Ja06<20%							
<sup>233</sup> Fr	48920	20	900	ms	100	1/2 <sup>+</sup> #	14	2010 $\beta^- =100; \beta^- n=0#$
<sup>233</sup> Ra	44334	9	30	s	5	1/2 <sup>+</sup> #	05	1990 $\beta^- =100$
<sup>233</sup> Ac	41308	13	145	s	10	(1/2 <sup>+</sup> )	05	1983 $\beta^- =100$
<sup>233</sup> Th	38731.7	1.4	21.83	m	0.04	(1/2) <sup>+</sup>	05	1935 $\beta^- =100$
<sup>233</sup> Pa	37489.5	1.3	26.975	d	0.013	3/2 <sup>-</sup>	05	1938 $\beta^- =100$
<sup>233</sup> U	36919.2	2.3	159.2	ky	0.2	5/2 <sup>+</sup>	05	1947 $\alpha =100; SF<6e-11; ^{24}Ne=7.2e-11 9; \dots$
<sup>233</sup> Np	37950	50	36.2	m	0.1	5/2 <sup>+</sup> #	05 50Ma14 D	1950 $\beta^+ \approx 100; \alpha =0.0007$
<sup>233</sup> Np <sup>p</sup>	38000#	60# 50# 30#				(5/2 <sup>-</sup> )	05	
... A-group is continued on next page ...								

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>233</sup> Pu	40050	50	20.9 m	0.4	5/2 <sup>+</sup> #	05	1957	$\beta^+ \approx 100$ ; $\alpha = 0.12$ 5
<sup>233</sup> Am	43260#	100#	3.2 m	0.8	5/2 <sup>-</sup> #	05	00Sa52 TD	$\beta^+$ ?; $\alpha = 4.5$ 9
<sup>233</sup> Cm	47290	70	27 s	10	3/2 <sup>+</sup> #	05	10Kh06 TD	$\alpha = 20$ 10; $\beta^+ = 80$ 10
<sup>233</sup> Bk	52860#	220#	40 s	30		15	15De22 TD	$\alpha \approx 82$ ; $\beta^+$ ?
* <sup>233</sup> U	D : ... ; <sup>28</sup> Mg < 1.3e-13							
* <sup>233</sup> Np	D : $\alpha$ observed in 50Ma14 with $\beta^+/\alpha = 1.5e5$							
* <sup>233</sup> Am	D : combining 10Kh06 $\alpha < 6$ and 00Sa52 $\alpha > 3$							
* <sup>233</sup> Cm	T : symmetrized from 23(+13-6)							
* <sup>233</sup> Bk	T : symmetrized from 15De22=21(+48-17)							
<sup>234</sup> Ra	46931	8	30 s	10	0 <sup>+</sup>	07	1990	$\beta^- = 100$ ; $\beta^-$ -SF < 1e-4
<sup>234</sup> Ac	44841	14	45 s	2	1 <sup>+</sup> #	07	08Ch.A T	$\beta^- = 100$
<sup>234</sup> Th	40613.0	2.6	24.10 d	0.03	0 <sup>+</sup>	07	1900	$\beta^- = 100$ ; $\alpha < 1e-4$
<sup>234</sup> Pa	40339	4	6.70 h	0.05	4 <sup>+</sup>	07	78Ga07 D	$\beta^- = 100$ ; SF < 3e-10
<sup>234</sup> Pa <sup>m</sup>	40417.9	2.8	79	3			78Ga07 D	$\beta^- \approx 100$ ; IT=0.16 4; SF < 1e-10
<sup>234</sup> U	38145.0	1.1	245.5 ky	0.6	0 <sup>+</sup>	07	1912	IS=0.0054 5; $\alpha = 100$ ; SF=1.64e-9 22; ...
<sup>234</sup> U <sup>m</sup>	39566.3	1.1	1421.257	0.017			1963	IT=100
<sup>234</sup> Np	39955	8	4.4 d	0.1	(0 <sup>+</sup> )	07	1949	$\beta^+ = 100$
<sup>234</sup> Pu	40350	7	8.8 h	0.1	0 <sup>+</sup>	07	1949	$\epsilon \approx 94$ ; $\alpha \approx 6$
<sup>234</sup> Am	44460#	160#	2.32 m	0.08		07	90Ha02 D	$\beta^+ \approx 100$ ; $\alpha = 0.039$ 12; $\beta^+$ -SF=0.0066 18
<sup>234</sup> Cm	46725	17	52 s	9	0 <sup>+</sup>	07	10Kh06 D	$\beta^+ \approx 71$ ; $\alpha \approx 27$ ; SF $\approx 2$
<sup>234</sup> Bk	53460#	140#	20 s	5		07	16Ka13 T	$\alpha > 80$ ; $\beta^+ < 20$
* <sup>234</sup> Ac	I : 08Ch.A reports two excited isomers with $T > 93$ s and $T = 149(+95-42)$ s							
* <sup>234</sup> Pa <sup>m</sup>	E : less than 10 keV above (3 <sup>+</sup> ) level at 73.92(0.02), see ENSDF2007							
* <sup>234</sup> U	D : ... ; <sup>28</sup> Mg=1.4e-11 3; <sup>24</sup> Ne+ <sup>26</sup> Ne=9e-12 7							
* <sup>234</sup> Am	T : also 04Sa05=3.5(1.3) not used							
* <sup>234</sup> Cm	T : average 16Ka13=49(+15-9) 01Ca.B=51(12)							
* <sup>234</sup> Bk	T : symmetrized from 16Ka13=19(+6-4) s							
<sup>235</sup> Ra	51130#	300#	3#	s	5/2 <sup>+</sup> #			$\beta^-$ ?
<sup>235</sup> Ac	47357	14	62 s	4	1/2 <sup>+</sup> #	14	08Ch.A T	$\beta^-$ ?
<sup>235</sup> Th	44018	13	7.2 m	0.1	1/2 <sup>+</sup> #	14	1969	$\beta^- = 100$
<sup>235</sup> Pa	42289	14	24.4 m	0.2	(3/2 <sup>-</sup> )	14	1950	$\beta^- = 100$
<sup>235</sup> U	40918.8	1.1	704 My	1	7/2 <sup>-</sup>	14	1935	IS=0.7204 6; $\alpha = 100$ ; ...
<sup>235</sup> U <sup>m</sup>	40918.9	1.1	0.0760	0.0004			16Ch11 T	IT=100
<sup>235</sup> U <sup>n</sup>	43420	300	2500	300			2007	SF ?
<sup>235</sup> Np	41043.1	1.4	396.1 d	1.2	5/2 <sup>+</sup>	14	1949	$\epsilon \approx 100$ ; $\alpha = 0.00260$ 13
<sup>235</sup> Pu	42182	21	25.3 m	0.5	(5/2 <sup>+</sup> )	14	1957	$\beta^+ \approx 100$ ; $\alpha = 0.0028$ 7
<sup>235</sup> Am	44630	50	10.3 m	0.6	5/2 <sup>-</sup> #	14	1996	$\beta^+ \approx 100$ ; $\alpha = 0.40$ 5
<sup>235</sup> Cm	48030#	200#	5#	m	5/2 <sup>+</sup> #	14		$\beta^+$ ?; $\alpha$ ?
<sup>235</sup> Cm <sup>p</sup>	48080#	210#	50#	50#		am		
<sup>235</sup> Bk	52700#	400#	1#	m				$\beta^+$ ?; $\alpha$ ?
* <sup>235</sup> U	D : ... ; SF=7e-9 2; <sup>20</sup> Ne=8e-10 4; <sup>25</sup> Ne $\approx$ 8e-10; <sup>28</sup> Mg=8e-10							
<sup>236</sup> Ac	51220	40	4.5 m	3.6		15	10Ch19 T	$\beta^-$ ?
<sup>236</sup> Th	46255	14	37.3 m	1.5	0 <sup>+</sup>	15	1973	$\beta^- = 100$
<sup>236</sup> Pa	45334	14	9.1 m	0.1	1 <sup>(-)</sup>	06	1963	$\beta^- = 100$ ; $\beta^-$ -SF=6e-8 4
<sup>236</sup> U	42444.6	1.1	23.42 My	0.03	0 <sup>+</sup>	06	1951	$\alpha = 100$ ; SF=9.4e-8 4
<sup>236</sup> U <sup>m</sup>	45195	3	2750	3			1969	IT=87 6; SF=13 6; $\alpha < 10$
<sup>236</sup> Np	43380	50	153 ky	5	(6 <sup>-</sup> )	06	1949	$\epsilon = 86.3$ 8; $\beta^- = 13.5$ 8; $\alpha = 0.16$ 4
<sup>236</sup> Np <sup>m</sup>	43438	7	60	50			1949	$\epsilon = 50$ 3; $\beta^- = 50$ 3
<sup>236</sup> Np <sup>p</sup>	43616	14	240	50			AD	
<sup>236</sup> Pu	42901.6	1.8	2.858 y	0.008	0 <sup>+</sup>	06	90Qg01 D	$\alpha = 100$ ; SF=1.9e-7 4; <sup>28</sup> Mg=2e-12; $2\beta^+$ ?
<sup>236</sup> Pu <sup>m</sup>	44087.0	1.8	1.2 $\mu$ s	0.3	5 <sup>-</sup>	06	2005	IT=100
<sup>236</sup> Am	46040#	110#	3.6 m	0.1	(5 <sup>-</sup> )	06	04Sa05 D	$\beta^+ = ?$ ; $\alpha = 4.0e-3$ 1
<sup>236</sup> Am <sup>m</sup>	46090#	120#	2.9 m	0.2	(1 <sup>-</sup> )	06	2004	$\beta^+ = ?$ ; $\alpha = ?$
<sup>236</sup> Cm	47855	18	6.8 m	0.8	0 <sup>+</sup>	06	10Kh06 TD	$\beta^+ = 82$ 2; $\alpha = 18$ 2; SF < 0.1
<sup>236</sup> Bk	53540#	400#	2#	m		06		$\beta^+$ ?; $\alpha$ ?
* <sup>236</sup> Ac	T : symmetrized from 10Ch19=72(+345-33) s							
* <sup>236</sup> Pa	D : $\beta^-$ -SF decay questioned in 90Ha02							
* <sup>236</sup> U	D : and Ne+Mg < 4e-10%, from 89Mi.A							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>237</sup> Ac	54020#	400#	4# m	1/2 <sup>+</sup> #				$\beta^-$ ?		
<sup>237</sup> Th	49955	16	4.8 m	0.5	5/2 <sup>+</sup> #	06	1993	$\beta^-$ =100		
<sup>237</sup> Pa	47528	13	8.7 m	0.2	(1/2 <sup>+</sup> )	06	1954	$\beta^-$ =100		
<sup>237</sup> U	45390.2	1.2	6.752 d	0.002	1/2 <sup>+</sup>	06	1940	$\beta^-$ =100		
<sup>237</sup> U <sup>m</sup>	45664.2	1.6	274.0	1.0	155 ns	6	(7/2 <sup>-</sup> )	06	IT=100	
<sup>237</sup> Np	44871.7	1.1	2.144 My	0.007	5/2 <sup>+</sup>	06	89Pr.A D	1948	$\alpha$ =100; SF<2e-10; <sup>30</sup> Mg<4e-12	
<sup>237</sup> Np <sup>m</sup>	45816.9	1.1	945.20	0.10	710 ns	40	(11/2,13/2)	06	1990	IT=100
<sup>237</sup> Pu	45091.7	1.7	45.64 d	0.04	7/2 <sup>-</sup>	06		1949	$\epsilon$ ≈100; $\alpha$ =0.0042 4	
<sup>237</sup> Pu <sup>m</sup>	45237.2	1.7	145.543	0.008	180 ms	20	1/2 <sup>+</sup>	06	1972	IT=100
<sup>237</sup> Pu <sup>n</sup>	47990	250	2900	250	1.1 $\mu$ s	0.1		06	1970	SF=?
<sup>237</sup> Am	46570#	60#	73.6 m	0.8	5/2 <sup>(-)</sup>	06		1970	$\beta^+$ ≈100; $\alpha$ =0.025 3	
<sup>237</sup> Cm	49250	70	20# m		5/2 <sup>+</sup> #	06	02As08 D	2002	$\beta^+$ ?; $\alpha$ =1.8	
<sup>237</sup> Cm <sup>p</sup>	49450#	170#	200#	150#		7/2 <sup>-</sup>				
<sup>237</sup> Bk	53190#	220#	2# m		(3/2 <sup>-</sup> )				$\beta^+$ ?; $\alpha$ ?	
<sup>237</sup> Cf	57940	90	0.8 s	0.2	5/2 <sup>+</sup> #	06	10Kh06 TD	1995	$\alpha$ =70 10; SF=30 10; $\beta^+$ ?	
* <sup>237</sup> Np	D : and cluster (Z=10-14) < 1.8e-12%, from 92Mo03								**	
* <sup>237</sup> Cm	D : partial $\alpha$ T=6.6e4 s or 1100 m								**	
* <sup>237</sup> Cf	T : others not used 95La09=2.1(0.3)								**	
<sup>238</sup> Th	52530#	280#	9.4 m	2.0	0 <sup>+</sup>	15		1999	$\beta^-$ =100	
<sup>238</sup> Pa	50894	16	2.28 m	0.09	3 <sup>-</sup> #	15	85Ba57 D	1968	$\beta^-$ =100; $\beta^-$ SF<2.6e-6	
<sup>238</sup> U	47307.8	1.5	4.468 Gy	0.006	0 <sup>+</sup>	15	91Tu02 D	1896	IS=99.2742 10; $\alpha$ =100; ...	
<sup>238</sup> U <sup>m</sup>	49865.7	1.6	2557.9	0.5	280 ns	6		1979	IT=?; SF=2.6 4; $\alpha$ <0.5	
<sup>238</sup> Np	47454.7	1.1	2.099 d	0.002	2 <sup>+</sup>	15		1949	$\beta^-$ =100	
<sup>238</sup> Np <sup>m</sup>	49760#	200#	2300#	200#	112 ns	39		1970	SF≈100; IT ?	
<sup>238</sup> Pu	46163.2	1.1	87.7 y	0.1	0 <sup>+</sup>	15	89Wa10 D	1949	$\alpha$ =100; SF=1.9e-7 1; ...	
<sup>238</sup> Am	48420	50	98 m	2	1 <sup>+</sup>	15		1950	$\beta^+$ =100; $\alpha$ =1.0e-4 4	
<sup>238</sup> Am <sup>m</sup>	50920#	210#	2500#	200#	35 $\mu$ s	18		1967	SF≈100; IT ?	
<sup>238</sup> Cm	49445	12	2.2 h	0.4	0 <sup>+</sup>	15		1994	$\epsilon$ ?; $\alpha$ =3.84 18; SF=0.048 2	
<sup>238</sup> Bk	54220#	260#	2.40 m	0.08		15		1994	$\beta^+$ ≈100; $\alpha$ ?; $\beta^+$ SF=0.048 2	
<sup>238</sup> Cf	57280#	300#	21.1 ms	1.3	0 <sup>+</sup>	15	01Og08 D	1995	SF≈100; $\alpha$ ≈0.2; $\beta^+$ ?	
* <sup>238</sup> U	D : ...; SF=5.45e-5 7; 2 $\beta^-$ =2.2e-10 7								**	
* <sup>238</sup> U	D : 2 $\beta^-$ =2.2(7)e-10% derived from 2 $\beta^-$ half-life T=2.0(0.6)Zy, in 91Tu02								**	
* <sup>238</sup> Pu	D : ...; <sup>32</sup> Si≈1.4e-14; <sup>28</sup> Mg+ <sup>30</sup> Mg≈6e-15								**	
<sup>239</sup> Th	56450#	400#	2# m		7/2 <sup>+</sup> #				$\beta^-$ ?	
<sup>239</sup> Pa	53340#	200#	1.8 h	0.5	(3/2 <sup>-</sup> ) <sup>(-#)</sup>	14		1995	$\beta^-$ =100	
<sup>239</sup> U	50572.7	1.5	23.45 m	0.02	5/2 <sup>+</sup>	14		1937	$\beta^-$ =100	
<sup>239</sup> U <sup>m</sup>	50593#	20#	20#	20#	> 250 ns		(5/2 <sup>+</sup> )	14	1994	$\beta^-$ =100
<sup>239</sup> U <sup>n</sup>	50706.5	1.5	133.7991	0.0010	780 ns	40	1/2 <sup>+</sup>	14	1975	IT=100
<sup>239</sup> Np	49311.1	1.3	2.356 d	0.003	5/2 <sup>+</sup>	14		1940	$\beta^-$ =100; $\alpha$ =5e-10#	
<sup>239</sup> Pu	48588.3	1.1	24.11 ky	0.03	1/2 <sup>+</sup>	14		1946	$\alpha$ =100; SF=3.1e-10 6	
<sup>239</sup> Pu <sup>m</sup>	48979.9	1.1	391.584	0.003	193 ns	4	7/2 <sup>-</sup>	14	1955	IT=100
<sup>239</sup> Pu <sup>n</sup>	51690	200	3100	200	7.5 $\mu$ s	1.0	(5/2 <sup>+</sup> )	14	1970	SF≈100; IT ?
<sup>239</sup> Am	49390.4	2.0	11.9 h	0.1	(5/2 <sup>-</sup> )	14		1949	$\epsilon$ ≈100; $\alpha$ =0.010 1	
<sup>239</sup> Am <sup>m</sup>	51890	200	2500	200	163 ns	12	(7/2 <sup>+</sup> )	14	1969	SF≈100; IT ?
<sup>239</sup> Cm	51150	50	2.5 h	0.4	(7/2 <sup>-</sup> )	14	02Sh.C TD	1952	$\beta^+$ ≈100; $\alpha$ =6.2e-3 14	
<sup>239</sup> Cm <sup>p</sup>	51390#	110#	240#	100#			1/2 <sup>+</sup>			
<sup>239</sup> Bk	54250#	210#	4# m		(7/2 <sup>+</sup> )	14	89Ha27 J		$\beta^+$ >99#; $\alpha$ <1; SF<1	
<sup>239</sup> Bk <sup>p</sup>	54290#	210#	41	11	AD		(3/2 <sup>-</sup> )	14	89Ha27 J	
<sup>239</sup> Cf	58270#	210#	60 s	30	5/2 <sup>+</sup> #	14		1981	$\alpha$ =?; $\beta^+$ ?	
<sup>239</sup> Es	63560#	300#	1# s						$\alpha$ ?; $\beta^+$ ?; SF ?	
* <sup>239</sup> Cf	T : symmetrized from 81Mu12=39(+37-12)								**	
<sup>240</sup> Pa	56910#	200#	2# m						$\beta^-$ ?	
<sup>240</sup> U	52715.5	2.6	14.1 h	0.1	0 <sup>+</sup>	08		1953	$\beta^-$ =100; $\alpha$ <1e-10#	
<sup>240</sup> Np	52316	17	* 61.9 m	0.2	(5 <sup>+</sup> )	08		1953	$\beta^-$ =100	
<sup>240</sup> Np <sup>m</sup>	52334	13	18	14	* 7.22 m	0.02	(1 <sup>+</sup> )	08	81Hs02 E	$\beta^-$ ≈100; IT=0.12 1
<sup>240</sup> Pu	50125.4	1.1	6.561 ky	0.007	0 <sup>+</sup>	08	13Sa65 D	1949	$\alpha$ =100; SF=5.63e-6 6; <sup>34</sup> Si<1.3e-11	
<sup>240</sup> Pu <sup>m</sup>	51434.1	1.1	1308.74	0.05	165 ns	10	(5 <sup>-</sup> )	08	1967	IT=100
<sup>240</sup> Am	51510	14	50.8 h	0.3	(3 <sup>-</sup> )	08		1949	$\beta^+$ =100; $\alpha$ ≈1.9e-4 7	
<sup>240</sup> Am <sup>m</sup>	54510	200	3000	200	940 $\mu$ s	40		1967	SF≈100; IT ?	
<sup>240</sup> Cm	51724.3	1.9	27 d	1	0 <sup>+</sup>	08		1949	$\alpha$ ≈100; $\epsilon$ <0.5; SF=3.9e-6 8	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>240</sup> Bk	55660# 150#		4.8 m 0.8			08	1980	$\beta^+$ ?; $\alpha=10\#$ ; $\beta^+$ SF=0.0020 13	*
<sup>240</sup> Bk <sup>p</sup>	55900# 180#	240# 100#		am					
<sup>240</sup> Cf	57991 19		40.3 s 0.9	0 <sup>+</sup>	08	10As.A T	1970	$\alpha=98.5$ 2; SF=1.5 2; $\beta^+$ ?	*
<sup>240</sup> Es	64200# 400#		1# s					$\alpha$ ?; $\beta^+$ ?	
* <sup>240</sup> Pu	D : SF=5.632(0.062)e-6 from SF half-life 13Sa65=116.5(1.3) Gy								**
* <sup>240</sup> Bk	D : symmetrized from $\beta^+$ SF=0.0013(+18-7)%								**
* <sup>240</sup> Cf	D : from 10Kh06; also 95La09 $\alpha\approx 98$ ; SF $\approx 2$								**
<sup>241</sup> Pa	59640# 300#		2# m	3/2 <sup>-</sup> #				$\beta^-$ ?	
<sup>241</sup> U	56200# 200#		5# m	7/2 <sup>+</sup> #	15			$\beta^-$ ?	
<sup>241</sup> Np	54260 70		13.9 m 0.2	(5/2 <sup>+</sup> )	15		1959	$\beta^-$ =100; $\alpha < 10e-6$	
<sup>241</sup> Pu	52955.2 1.1		14.329 y 0.029	5/2 <sup>+</sup>	15		1949	$\beta^- \approx 100$ ; $\alpha=0.00247$ ; SF<2.4e-14	
<sup>241</sup> Pu <sup>m</sup>	53116.9 1.1	161.6853 0.0009	880 ns 50	1/2 <sup>+</sup>	15		1975	IT=100	
<sup>241</sup> Pu <sup>m</sup>	55160 200	2200 200	20.5 $\mu$ s 2.2		15		1970	SF=100	
<sup>241</sup> Am	52934.4 1.1		432.6 y 0.6	5/2 <sup>-</sup>	15		1949	$\alpha=100$ ; SF=3.6e-10 9; 34Si<7.4e-14	
<sup>241</sup> Am <sup>m</sup>	55130 100	2200 100	1.2 $\mu$ s 0.3		15		1969	SF=100	
<sup>241</sup> Cm	53701.8 1.6		32.8 d 0.2	1/2 <sup>+</sup>	15		1952	$\epsilon=99.0$ 1; $\alpha=1.0$ 1	
<sup>241</sup> Bk	56030# 200#		4.6 m 0.4	(7/2 <sup>+</sup> )	15		2003	$\alpha$ ?; $\beta^+$ ?	
<sup>241</sup> Bk <sup>p</sup>	56080# 200#	51 3 AD		(3/2 <sup>-</sup> )	15				
<sup>241</sup> Cf	59330# 170#		2.35 m 0.18	7/2 <sup>-</sup> #	15	10As.A T	1970	$\beta^+$ ?; $\alpha\approx 25$	*
<sup>241</sup> Cf <sup>p</sup>	59480# 190#	150# 100#		(1/2 <sup>+</sup> )	15				
<sup>241</sup> Es	63860# 230#		10 s 5	(3/2 <sup>-</sup> )	15	96Ni09 TJD	1996	$\alpha=?$ ; $\beta^+$ ?	*
<sup>241</sup> Es <sup>p</sup>	64020# 300#	160# 200#		am					
<sup>241</sup> Fm	69130# 300#		730 $\mu$ s 60	5/2 <sup>+</sup> #	15		2008	SF=?; $\alpha < 14$ ; $\beta^+ < 12$	
* <sup>241</sup> Cf	T : from 10As.A=141(11) s; other 70Si19=3.78(0.70) m								**
* <sup>241</sup> Es	T : symmetrized from 96Ni09=8(+6-4)								**
<sup>242</sup> U	58620# 200#		16.8 m 0.5	0 <sup>+</sup>	02		1979	$\beta^-$ =100	
<sup>242</sup> Np	57420 200		2.2 m 0.2	(1 <sup>+</sup> )	02		1979	$\beta^-$ =100	
<sup>242</sup> Np <sup>m</sup>	57420# 210#	0# 50#	5.5 m 0.1	6 <sup>+</sup> #	02		1981	$\beta^-$ =100	
<sup>242</sup> Pu	54716.9 1.2		375 ky 2	0 <sup>+</sup>	02	13Sa65 D	1950	$\alpha=100$ ; SF=5.56e-4 7	*
<sup>242</sup> Am	55468.1 1.1		16.02 h 0.02	1 <sup>-</sup>	02		1949	$\beta^-$ =82.7 3; $\epsilon=17.3$ 3	
<sup>242</sup> Am <sup>m</sup>	55516.7 1.1	48.60 0.05	141 y 2	5 <sup>-</sup>	02		1950	IT $\approx 100$ ; $\alpha=0.45$ 2; SF<4.7e-9	
<sup>242</sup> Am <sup>m</sup>	57670 80	2200 80	14.0 ms 1.0	(2 <sup>+</sup> , 3 <sup>-</sup> )	02		1962	SF $\approx 100$ ; IT=?; $\alpha$ ?	
<sup>242</sup> Cm	54803.8 1.1		162.8 d 0.2	0 <sup>+</sup>	02		1949	$\alpha=100$ ; SF=6.2e-6 3; 34Si=1.1e-14 4; ...	*
<sup>242</sup> Cm <sup>m</sup>	57600 100	2800 100	180 ns 70		02		1971	SF ?; IT ?	
<sup>242</sup> Bk	57730# 200#		7.0 m 1.3	2 <sup>-</sup> #	02	80Ga07 D	1972	$\beta^+ \approx 100$ ; $\beta^+$ SF<3e-5; $\alpha$ ?	
<sup>242</sup> Bk <sup>m</sup>	57930# 280#	200# 200#	600 ns 100		02		1972	SF $\approx 100$ ; IT ?	
<sup>242</sup> Bk <sup>p</sup>	57980# 220#	250# 100#		4 <sup>-</sup>					
<sup>242</sup> Cf	59387 13		3.49 m 0.15	0 <sup>+</sup>	02	70Si19 T	1967	$\alpha=80$ 20; $\beta^+$ ?; SF<0.014	*
<sup>242</sup> Es	64800# 260#		17.8 s 1.6		02	10An08 TD	1994	$\alpha=57$ 3; $\beta^+=43$ 3; $\beta^+$ SF=0.6 2	*
<sup>242</sup> Fm	68400# 400#		800 $\mu$ s 200	0 <sup>+</sup>	02		1975	SF=?; $\alpha$ ?	*
* <sup>242</sup> Pu	D : SF=5.564(0.072)e-4 from SF half-life 13Sa65=67.4(0.9) Gy								**
* <sup>242</sup> Cm	D : ... ; $2\beta^+$ ? D : symmetrized from <sup>34</sup> Si=1.0(+4-3)e-14								**
* <sup>242</sup> Cf	T : average 70Si19=3.68(0.44) 67Si07=3.4(0.2) 67Fi04=3.2(0.5) 67Ii01=3.7(0.3)								**
* <sup>242</sup> Es	T : others 00Sh10=11(3) 96Ni09=16(+6-4)								**
* <sup>242</sup> Es	D : $\beta^+$ SF from 00Sh10; other 10An08=1.3(+1.2-0.7)%								**
* <sup>242</sup> Fm	T : conflicting 08Kh10 excludes 4 $\mu$ s-1s								**
<sup>243</sup> U	62360# 300#		10# m	9/2 <sup>-</sup> #				$\beta^-$ ?	
<sup>243</sup> Np	59880# 30#		1.85 m 0.15	5/2 <sup>+</sup> #	14		1979	$\beta^-$ =100	
<sup>243</sup> Np <sup>p</sup>	59926 10	50# 30#		(5/2 <sup>-</sup> )					
<sup>243</sup> Pu	57754.6 2.5		4.956 h 0.003	7/2 <sup>+</sup>	14		1951	$\beta^-$ =100	
<sup>243</sup> Pu <sup>m</sup>	58138.2 2.5	383.64 0.25	330 ns 30	(1/2 <sup>+</sup> )	14		1975	IT=100	
<sup>243</sup> Am	57175.0 1.4		7.364 ky 0.022	5/2 <sup>-</sup>	14		1950	$\alpha=100$ ; SF=3.7e-9 9	
<sup>243</sup> Am <sup>m</sup>	59480 200	2300 200	5.5 $\mu$ s 0.5		14		1970	SF $\approx 100$ ; IT ?	
<sup>243</sup> Cm	57182.0 1.5		29.1 y 0.1	5/2 <sup>+</sup>	14		1950	$\alpha\approx 100$ ; $\epsilon=0.29$ 3; SF=5.3e-9 9	
<sup>243</sup> Cm <sup>m</sup>	57269.4 1.5	87.4 0.1	1.08 $\mu$ s 0.03	1/2 <sup>+</sup>	14		1971	IT=100	
<sup>243</sup> Cm <sup>p</sup>	57279 16	97 16		(7/2 <sup>+</sup> )	14		1984	IT ?	
<sup>243</sup> Bk	58690 5		4.6 h 0.2	3/2 <sup>-</sup> #	14		1950	$\beta^+ \approx 100$ ; $\alpha\approx 0.15$	
<sup>243</sup> Bk <sup>p</sup>	58710 19	20 20		(7/2 <sup>-</sup> )					
<sup>243</sup> Cf	60990# 110#		10.7 m 0.5	(1/2 <sup>+</sup> )	14		1967	$\beta^+ \approx 86$ ; $\alpha\approx 14$	
<sup>243</sup> Es	64750# 210#		21.6 s 1.6	(7/2 <sup>+</sup> )	14		1973	$\alpha=61$ 6; $\beta^+=39$ 6; SF<1	
<sup>243</sup> Fm	69390# 220#		231 ms 9	7/2 <sup>-</sup> #	14		1981	$\alpha=91$ 3; SF=9 1; $\beta^+$ ?	*
* <sup>243</sup> Fm	D : 08Kh10 $\beta^+ < 10$								**

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)		Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
$^{244}\text{Np}$	63200#	300#			2.29	m	0.16	(7 <sup>-</sup> )	03		1987	$\beta^- = 100$	
$^{244}\text{Pu}$	59806.0	2.3			80.0	My	0.9	0 <sup>+</sup>	03	92Mo25	D	1954	$\alpha \approx 100$ ; SF=0.121 4; $2\beta^- < 7.3e-9$ *
$^{244}\text{Pu}^m$	61022	3	1216	2	1.75	s	0.12	(8 <sup>-</sup> )		16Ho13	ETJ	2016	IT=100
$^{244}\text{Am}$	59879.2	1.5			10.1	h	0.1	6 <sup>-</sup> #	03			1950	$\beta^- = 100$
$^{244}\text{Am}^m$	59968.5	1.4	89.3	1.6	26	m	1	1 <sup>+</sup>	03			1950	$\beta^- \approx 100$ ; $\epsilon = 0.0361$ 13
$^{244}\text{Am}^n$	60080#	200#	200#	200#	900	$\mu\text{s}$	150		03			1967	SF $\approx 100$ ; IT ?
$^{244}\text{Am}^p$	60080#	200#	200#	200#	6.5	$\mu\text{s}$			03			1969	SF $\approx 100$ ; IT ?
$^{244}\text{Cm}$	58451.9	1.1			18.10	y	0.02	0 <sup>+</sup>	03			1950	$\alpha = 100$ ; SF=1.37e-4 3
$^{244}\text{Cm}^m$	59492.1	1.1	1040.188	0.012	34	ms	2	6 <sup>+</sup>	03			1963	IT=100
$^{244}\text{Cm}^n$	59550#	900#	1100#	900#	> 500	ns			03			1969	SF $\approx 100$ ; IT ?
$^{244}\text{Bk}$	60714	14			5.02	h	0.03	(4 <sup>-</sup> )	03	14So17	T	1972	$\beta^+ ?$ ; $\alpha = 0.006$ 3
$^{244}\text{Bk}^m$	61210#	300#	500#	300#	820	ns	60		03			1972	SF $\approx 100$ ; IT ?
$^{244}\text{Bk}^p$	60850#	50#	140#	50#					<i>am</i>				
$^{244}\text{Cf}$	61478.2	2.6			19.4	m	0.6	0 <sup>+</sup>	03			1956	$\alpha \approx 100$ ; $\epsilon ?$
$^{244}\text{Es}$	66030#	180#			37	s	4		03			1973	$\beta^+ ?$ ; $\alpha = 5$ 3; $\beta^+$ SF=0.01 *
$^{244}\text{Es}^p$	66230#	240#	200#	150#					<i>am</i>				
$^{244}\text{Fm}$	68970#	200#			3.12	ms	0.08	0 <sup>+</sup>	03	08Kh10	TD	1967	SF $\approx 100$ ; $\beta^+ < 2$ ; $\alpha = 0.4\%$ *
* $^{244}\text{Pu}$	T : and $T(2\beta^-) > 1.1$ Ey, from 92Mo25; thus $2\beta^- < 7.3 e-9\%$ **												
* $^{244}\text{Es}$	D : symmetrized from $\alpha = 4(+3-2)\%$ **												
* $^{244}\text{Fm}$	T : other 12Sv02=3.47(0.26) **												
$^{245}\text{Np}$	65890#	300#			2#	m		5/2 <sup>+</sup> #					$\beta^- ?$
$^{245}\text{Pu}$	63178	14			10.5	h	0.1	(9/2 <sup>-</sup> )	11			1955	$\beta^- = 100$
$^{245}\text{Pu}^m$	63443	14	264.5	0.3	330	ns	20	(5/2 <sup>+</sup> )	11			2007	IT=100
$^{245}\text{Am}$	61900.5	1.9			2.05	h	0.01	(5/2 <sup>+</sup> ) <sup>+</sup>	11			1955	$\beta^- = 100$
$^{245}\text{Am}^m$	64300#	400#	2400#	400#	640	ns	60		11			1972	SF $\approx 100$ ; IT ?
$^{245}\text{Cm}$	61004.6	1.1			8.25	ky	0.07	7/2 <sup>+</sup>	11	12Ch30	T	1954	$\alpha = 100$ ; SF=6.1e-7 9
$^{245}\text{Cm}^m$	61360.5	1.1	355.92	0.10	290	ns	20	1/2 <sup>+</sup>	11			1975	IT=100
$^{245}\text{Bk}$	61813.8	1.8			4.95	d	0.03	3/2 <sup>-</sup>	11			1951	$\epsilon \approx 100$ ; $\alpha = 0.12$ 1
$^{245}\text{Bk}^p$	61860#	30#	50#	30#				(7/2 <sup>-</sup> )					
$^{245}\text{Cf}$	63385.2	2.4			45.0	m	1.5	1/2 <sup>+</sup>	11			1956	$\beta^+ ?$ ; $\alpha = 36$ 3
$^{245}\text{Es}$	66370#	200#			1.1	m	0.1	(3/2 <sup>-</sup> )	11			1967	$\beta^+ ?$ ; $\alpha = 40$ 10
$^{245}\text{Es}^p$	66650#	200#	283	15				(7/2 <sup>-</sup> )	11			2005	IT=100 *
$^{245}\text{Es}^q$	66700#	230#	330#	100#				(1/2 <sup>-</sup> )					
$^{245}\text{Fm}$	70190#	200#			4.2	s	1.3	1/2 <sup>+</sup> #	11			1967	$\alpha = ?$ ; $\beta^+ = 4.2\%$ ; SF=0.13#
$^{245}\text{Md}$	75270#	310#			400	ms	200	(7/2 <sup>-</sup> )	11	96Ni09	TJD	1996	$\alpha = ?$ ; $\beta^+ ?$ *
$^{245}\text{Md}^m$	75370#	330#	100#	100#	900	$\mu\text{s}$	250	1/2 <sup>-</sup> #	11			1996	SF $\approx ?$ ; $\alpha ?$ *
* $^{245}\text{Es}^p$	E : 253.2 keV above the 7/2 <sup>+</sup> [633] level at 30(15) keV **												
* $^{245}\text{Md}$	T : symmetrized from 96Ni09=350(+230-160) **												
$^{246}\text{Pu}$	65395	15			10.84	d	0.02	0 <sup>+</sup>	11			1955	$\beta^- = 100$
$^{246}\text{Am}$	64994#	18#			39	m	3	(7 <sup>-</sup> )	11			1955	$\beta^- = 100$
$^{246}\text{Am}^m$	65024	15	30#	10#	25.0	m	0.2	2 <sup>(-)</sup>	11			1955	$\beta^- \approx 100$ ; IT<0.02
$^{246}\text{Am}^n$	66990#	800#	2000#	800#	73	$\mu\text{s}$	10		11			1972	SF $\approx 100$ ; IT ?
$^{246}\text{Cm}$	62617.0	1.5			4.706	ky	0.040	0 <sup>+</sup>	11			1954	$\alpha \approx 100$ ; SF=0.02615 7
$^{246}\text{Cm}^m$	63796.7	1.5	1179.66	0.13	1.12	s	0.24	8 <sup>-</sup>	11	12Ta.A	ETJ	2012	IT=100
$^{246}\text{Bk}$	63970	60			1.80	d	0.02	2 <sup>(-)</sup>	11			1954	$\beta^+ \approx 100$ ; $\alpha = 0.1\%$
$^{246}\text{Cf}$	64090.3	1.5			35.7	h	0.5	0 <sup>+</sup>	11			1951	$\alpha = 100$ ; SF=2.4e-4 4; $\epsilon < 4e-3$
$^{246}\text{Es}$	67900#	220#			7.5	m	0.5	4 <sup>-</sup> #	11			1954	$\beta^+ = 90.1$ 18; $\alpha = 9.9$ 18; $\beta^+$ SF $\approx 0.003$
$^{246}\text{Es}^p$	68250#	300#	350#	200#				<i>am</i>					
$^{246}\text{Fm}$	70189	15			1.54	s	0.04	0 <sup>+</sup>	11	10An08	T	1966	$\alpha = ?$ ; SF=6.8 6; $\epsilon < 1.3$ ; $\beta^+$ SF=10 5 *
$^{246}\text{Md}$	76120#	260#			0.92	s	0.18		11	10An08	TD	1996	$\alpha = 100$ *
$^{246}\text{Md}^m$	76170#	260#	60	60	4.4	s	0.8		11			2010	$\beta^+ > 77$ ; $\beta^+$ SF>10; $\alpha < 23$ *
* $^{246}\text{Es}^p$	E : above level decaying by 152.3(0.5) keV $\gamma$ **												
* $^{246}\text{Fm}$	D : from 96Ni09 **												
* $^{246}\text{Md}$	T : average 10An08=0.9(0.2) 96Ni09=1.0(0.4) **												
$^{247}\text{Pu}$	69110#	200#			2.27	d	0.23	1/2 <sup>+</sup> #	15			1983	$\beta^- = 100$
$^{247}\text{Am}$	67150#	100#			23.0	m	1.3	5/2#	15			1967	$\beta^- = 100$
$^{247}\text{Cm}$	65533	4			15.6	My	0.5	9/2 <sup>-</sup>	15			1954	$\alpha = 100$
$^{247}\text{Cm}^m$	65760	4	227.38	0.19	26.3	$\mu\text{s}$	0.3	5/2 <sup>+</sup>	15			1968	IT=100
$^{247}\text{Cm}^n$	65938	4	404.90	0.03	100.6	ns	0.6	1/2 <sup>+</sup>	15			2003	IT=100
$^{247}\text{Bk}$	65490	5			1.38	ky	0.25	3/2 <sup>-</sup>	15			1965	$\alpha \approx 100$ ; SF ?
$^{247}\text{Cf}$	66104	15			3.11	h	0.03	7/2 <sup>+</sup> #	15			1954	$\epsilon \approx 100$ ; $\alpha = 0.035$ 5
$^{247}\text{Es}$	68578	19			4.55	m	0.26	(7/2 <sup>+</sup> )	15	89Ha27	J	1967	$\beta^+ \approx 93$ ; $\alpha \approx 7$ ; SF $\approx 9e-5$ #
$^{247}\text{Fm}$	71670#	120#			31	s	1	(7/2 <sup>+</sup> )	15			1967	$\alpha = 64$ ; $\beta^+ ?$
$^{247}\text{Fm}^m$	71720#	110#	49	8	5.1	s	0.2	(1/2 <sup>+</sup> )	15			1967	$\alpha = 88$ 2; $\beta^+ ?$ ; IT ?
$^{247}\text{Md}$	75940#	210#			1.2	s	0.1	(7/2 <sup>-</sup> )	15			1981	$\alpha \approx 100$ ; SF<0.1
$^{247}\text{Md}^m$	76200#	210#	260	40	250	ms	40	(1/2 <sup>-</sup> )	15	10An08	D	1993	$\alpha = 79$ 5; SF=21 5

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>248</sup> Am	70560#	200#	3# m		14			$\beta^-$ ?		
<sup>248</sup> Cm	67392.8	2.4	348 ky	6	0 <sup>+</sup>	14	1956	$\alpha=91.61$ 16; SF=8.39 16; $2\beta^-$ ?		
<sup>248</sup> Cm <sup>m</sup>	68850.9	2.6	146 $\mu$ s	18	(8 <sup>-</sup> )	14	12Ta.A ETJ 2012	IT=100		
<sup>248</sup> Bk	68080#	70#	> 9 y		6 <sup>+</sup> #	14	1956	$\alpha$ ?		
<sup>248</sup> Bk <sup>m</sup>	68108	21	30# 70#	*	23.7 h 0.2	1 <sup>(-)</sup>	14	$\beta^-$ =70 5; $\epsilon$ =30 5; $\alpha$ =0.001#		
<sup>248</sup> Bk <sup>p</sup>	68130	50	50# 50#			(5 <sup>-</sup> )				
<sup>248</sup> Cf	67238	5	333.5 d	2.8	0 <sup>+</sup>	14	1954	$\alpha \approx 100$ ; SF=0.0029 3		
<sup>248</sup> Es	70300#	50#	24 m	3	2 <sup>-</sup> #	14	1956	$\beta^+ \approx 100$ ; $\alpha \approx 0.25$ ; $\beta^+$ SF=3e-5		
<sup>248</sup> Fm	71898	8	34.5 s	1.2	0 <sup>+</sup>	14	1958	$\alpha=95$ 5; $\beta^+=5$ 5; SF=0.10 5		
<sup>248</sup> Fm <sup>m</sup>	73100#	100#	1200# 100#		10.1 ms 0.6		14	$\alpha$ ?; B ?		
<sup>248</sup> Md	77150#	240#	7 s	3		14	1973	$\beta^+=80$ 10; $\alpha=20$ 10; $\beta^+$ SF<0.05		
<sup>248</sup> No	80620#	220#	<2 $\mu$ s	0 <sup>+</sup>	14	03Be18 I		SF ?		
<sup>249</sup> Am	73100#	300#	1# m					$\beta^-$ ?		
<sup>249</sup> Cm	70750.7	2.4	64.15 m	0.03	(1/2 <sup>+</sup> )	11	1956	$\beta^-$ =100		
<sup>249</sup> Cm <sup>m</sup>	70799.5	2.4	23 $\mu$ s		(7/2 <sup>+</sup> )	11	1966	$\alpha=100$		
<sup>249</sup> Bk	69846.4	1.2	327.2 d	0.3	7/2 <sup>+</sup>	11	14Ch47 T 1954	$\beta^- \approx 100$ ; $\alpha=0.00145$ 8; SF=47e-9 2		
<sup>249</sup> Bk <sup>m</sup>	69855.2	1.2	300 $\mu$ s		(3/2 <sup>-</sup> )	11	1975	IT=100		
<sup>249</sup> Cf	69722.8	1.2	351 y	2	9/2 <sup>-</sup>	11	1954	$\alpha=100$ ; SF=5.0e-7 4		
<sup>249</sup> Cf <sup>m</sup>	69867.8	1.2	144.98	0.05	45 $\mu$ s	5	5/2 <sup>+</sup>	11 1967	IT=100	
<sup>249</sup> Es	71180#	30#	102.2 m	0.6	7/2 <sup>+</sup>	11	1956	$\beta^+ \approx 100$ ; $\alpha=0.57$ 8		
<sup>249</sup> Fm	73519	6	1.6 m	0.1	(7/2 <sup>+</sup> )	11	11Lo06 J 1960	$\beta^+ ?$ ; $\alpha=33$ 9		
<sup>249</sup> Md	77230#	200#	23.4 s	2.4	(7/2 <sup>-</sup> )	11	01He35 J 1973	$\alpha > 60$ ; $\beta^+ ?$		
<sup>249</sup> Md <sup>m</sup>	77330#	220#	100# 100#		1.9 s 0.9	(1/2 <sup>-</sup> )	11 01He35 TJD 2001	$\alpha=100$		
<sup>249</sup> No	81780#	280#	57 $\mu$ s	12	5/2 <sup>+</sup> #	11	03Be18 T 2003	$\beta^+ ?$ ; $\alpha ?$		
* <sup>249</sup> Fm	T : from 04He28; others 66Ak01=2.6(0.7) 59Pe27=2.5(1.0)									
* <sup>249</sup> Md	T : average 09He20=23(3) 73Es01=24(4)									
* <sup>249</sup> Md <sup>m</sup>	T : symmetrized from 1.5(+1.2-0.5)									
* <sup>249</sup> No	T : symmetrized from 54.0(+13.9-9.2)									
<sup>250</sup> Cm	72990	10	8300# y		0 <sup>+</sup>	01	1966	SF $\approx$ 74; $\alpha \approx 18$ ; $\beta^- \approx 8$		
<sup>250</sup> Bk	72950	4	3.212 h	0.005	2 <sup>-</sup>	01	1954	$\beta^-$ =100		
<sup>250</sup> Bk <sup>m</sup>	72986	4	35.59	0.10	29 $\mu$ s	1	4 <sup>+</sup>	01 08Ah02 EJ 1966	IT=100	
<sup>250</sup> Bk <sup>n</sup>	73034	4	84.1	2.1	AD	213 $\mu$ s	8	7 <sup>+</sup>	01 08Ah02 EJ 1972	IT ?
<sup>250</sup> Cf	71170.4	1.5	13.08 y	0.09	0 <sup>+</sup>	01	1954	$\alpha \approx 100$ ; SF=0.077 3		
<sup>250</sup> Es	73230#	100#	8.6 h	0.1	(6 <sup>+</sup> )	01	1956	$\beta^+ > 97$ ; $\alpha ?$		
<sup>250</sup> Es <sup>m</sup>	73430#	180#	200# 150#	*	2.22 h	0.05	1 <sup>(-)</sup>	01 1970	$\beta^+ \approx 100$ ; $\alpha ?$	
<sup>250</sup> Fm	74072	8	30.4 m	1.5	0 <sup>+</sup>	01	06Ba09 T 1954	$\alpha > 90$ ; $\epsilon < 10$ ; SF=0.0069 10		
<sup>250</sup> Fm <sup>m</sup>	75271	8	1199.2	1.0	1.92 s	0.05	(8 <sup>-</sup> )	01 08Gr17 ETJ 1973	IT $>$ 80; $\alpha < 20$ ; $\beta^+ ?$ ; SF<8.2E-5	
<sup>250</sup> Md	78630#	300#	52 s	5	01	08An16 TD 1973		$\beta^+=93$ 1; $\alpha=7$ 1; $\beta^+$ SF=0.02		
<sup>250</sup> No	81560#	200#	5.0 $\mu$ s	0.6	0 <sup>+</sup>	06	06Pe17 TD 2003	SF $\approx$ 100; $\alpha < 2.1$ ; $\beta^+ =0.00025$ #		
<sup>250</sup> No <sup>m</sup>	82610#	280#	1050# 200#		51 $\mu$ s	18	(6 <sup>+</sup> )	06 06Pe17 T 2001	SF $\approx$ 100; IT ?; $\alpha ?$	
* <sup>250</sup> Fm	T : others not used 06Fo02=18(+13-6) 66Ak01=30(3)									
* <sup>250</sup> Md	T : average 08An16=50(+10-7) 73Es01=52(6)									
* <sup>250</sup> Md	D : other recent 06Fo02 $\beta^+=91(+7-19)\%$ ; $\alpha=9(+19-7)\%$									
* <sup>250</sup> No	T : average 06Pe17=3.7(+1.1-0.8) 03Be18=5.6(+0.9-0.7)									
* <sup>250</sup> No <sup>m</sup>	T : average 06Pe17=43(+22-15) 03Be18=46(+22-14) 01Og08=36(+11-6)									
<sup>251</sup> Cm	76648	23	16.8 m	0.2	(1/2 <sup>+</sup> )	13	1978	$\beta^-$ =100		
<sup>251</sup> Bk	75228	11	55.6 m	1.1	(3/2 <sup>-</sup> )	13	1967	$\beta^-$ =100		
<sup>251</sup> Bk <sup>m</sup>	75264	11	58 $\mu$ s	4	7/2 <sup>+</sup> #	13	1966	IT=100		
<sup>251</sup> Cf	74135	4	900 y	40	1/2 <sup>+</sup>	13	1954	$\alpha \approx 100$ ; SF ?		
<sup>251</sup> Cf <sup>m</sup>	74505	4	370.47	0.03	1.3 $\mu$ s	0.1	11/2 <sup>-</sup>	13 1971	IT=100	
<sup>251</sup> Es	74512	6	33 h	1	3/2 <sup>-</sup>	13	1956	$\epsilon ?$ ; $\alpha=0.5$ 2		
<sup>251</sup> Fm	75954	15	5.30 h	0.08	(9/2 <sup>-</sup> )	13	1957	$\beta^+=98.20$ 13; $\alpha=1.80$ 13		
<sup>251</sup> Fm <sup>m</sup>	76154	15	200.00	0.10	21.1 $\mu$ s	1.9	5/2 <sup>+</sup>	13 1970	IT=100	
<sup>251</sup> Md	78967	19	4.21 m	0.23	(7/2 <sup>-</sup> )	13	06Ch52 TD 1973	$\beta^+ ?$ ; $\alpha=10$ 1		
<sup>251</sup> Md <sup>p</sup>	79020	18	53	8	AD		(1/2 <sup>-</sup> )	13 2006	IT ?	
<sup>251</sup> No	82850#	110#	800 ms	10	(7/2 <sup>+</sup> )	13	06He27 J 1967	$\alpha=83$ 16; $\beta^+ ?$ ; SF<0.3		
<sup>251</sup> No <sup>m</sup>	82960#	110#	106	6	1.02 s	0.03	(1/2 <sup>+</sup> )	13 1997	$\alpha=100$	
<sup>251</sup> No <sup>n</sup>	84600#	120#	1750	50	2 $\mu$ s		13	2006	IT ?	
<sup>251</sup> Lr	87730#	300#	150# $\mu$ s					$\beta^+ ?$ ; $\alpha ?$		
* <sup>251</sup> Fm <sup>m</sup>	T : 11As03=21.1(1.9) 06He20=21(3) 71Di03=15.2(2.3)									
* <sup>251</sup> Fm <sup>n</sup>	E : 11As03=200.09(0.11) 06He20=199.9(0.3)									
* <sup>251</sup> Md	T : average 06Ch52=4.27(0.26) 73Es01=4.0(0.5)									
* <sup>251</sup> No	D : symmetrized from 01He35 $\alpha=91(+9-22)\%$									
* <sup>251</sup> No <sup>n</sup>	E : 1699.7(0.8) + x ; x estimated 50(50)									



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>252</sup> Cm	79060# 300#		1# m <2d	0 <sup>+</sup>	06	66Rg01 I		$\beta^-$ ?
<sup>252</sup> Bk	78540# 200#		1.8 m 0.5		06	92Kr.A TD	1992	$\beta^-$ =?; $\alpha$ ?
<sup>252</sup> Cf	76034.6 2.4		2.645 y 0.008	0 <sup>+</sup>	06		1954	$\alpha$ =96.908 8; SF=3.092 8
<sup>252</sup> Es	77290 50		471.7 d 1.9	(4 <sup>+</sup> )	06	FGK12a J	1956	$\alpha$ =78 2; $\epsilon$ =22 2
<sup>252</sup> Fm	76816 5		25.39 h 0.04	0 <sup>+</sup>	06		1956	$\alpha$ ≈100; SF=0.0023 2; $2\beta^+$ ?
<sup>252</sup> Md	80510# 130#		2.3 m 0.8		06		1973	$\beta^+$ >50; $\alpha$ ?
<sup>252</sup> Md <sup>p</sup>	80550 80	40# 100#		<i>am</i>				
<sup>252</sup> No	82871 9		2.45 s 0.02	0 <sup>+</sup>	06	11Ga19 T	1967	$\alpha$ >66.7 6; SF=32.2 5; $\beta^+$ <1.1 4
<sup>252</sup> No <sup>m</sup>	84126 9	1254.5 0.7	109 ms 4	(8 <sup>-</sup> )		11Lo06 T	2007	IT=100
<sup>252</sup> Lr	88740# 240#		369 ms 75		06	08Ne01 TD	2001	$\beta^+$ =71#; $\alpha$ =?; SF<1
<sup>252</sup> Lr <sup>p</sup>	88910# 240#	170 30 AD						
* <sup>252</sup> Es	J : strong direct $\epsilon$ feeding to 3 <sup>+</sup> ; known structures in TNN							
* <sup>252</sup> No	T : average 11Ga19=2.47(0.02) 01Og08=2.44(0.04)							
* <sup>252</sup> No	T : others 12Sv02=2.3(0.1) 04He28=2.52(0.22) 03Be18=2.38(+0.26-0.22)							
* <sup>252</sup> No	D : SF 01Og08=32.2(0.5)%; other 11Ga19=29.3(0.5)%							
* <sup>252</sup> No <sup>m</sup>	E : average 08Ro21=1255(1) 07Su19=1254(1)							
* <sup>252</sup> No <sup>m</sup>	T : average 11Lo06=110(8) 08Ro21=109(6) 07Su19=110(10)							
* <sup>252</sup> No <sup>m</sup>	J : from 08Ro21 based on comparison with theory; other 07Su19=(8 <sup>+</sup> )							
* <sup>252</sup> Lr	T : average 08Ne01=270(+180-80) 01He35=360(+110-70)							
<sup>253</sup> Bk	80930# 360#		10# m		13	91Kr.A I	1991	$\beta^-$ ?
<sup>253</sup> Cf	79302 4		17.81 d 0.08	(7/2 <sup>+</sup> )	13		1954	$\beta^-$ ≈100; $\alpha$ =0.31 4
<sup>253</sup> Es	79010.5 1.2		20.47 d 0.03	7/2 <sup>+</sup>	13	05Ah03 D	1954	$\alpha$ =100; SF=10e-6 1
<sup>253</sup> Fm	79345.7 2.9		3.00 d 0.12	(1/2 <sup>+</sup> )	13		1957	$\epsilon$ =88 1; $\alpha$ =12 1
<sup>253</sup> Fm <sup>m</sup>	79697 7	351 6	560 ns 60	(11/2 <sup>-</sup> )	13	11An13 E/TJ	2011	IT=100
<sup>253</sup> Md	81170# 30#		12 m 8	(7/2 <sup>-</sup> )	13		1992	$\beta^+$ ≈100; $\alpha$ =0.6#
<sup>253</sup> Md <sup>p</sup>	81230# 40#	60 30		1/2 <sup>-</sup> #	13		1971	IT ?
<sup>253</sup> No	84359 7		1.56 m 0.02	(9/2 <sup>-</sup> )	13		1967	$\alpha$ =55 3; $\beta^+$ ?; SF=0.001#
<sup>253</sup> No <sup>m</sup>	84526 7	167.34 0.45	30.3 $\mu$ s 1.6	(5/2 <sup>+</sup> )	13	09He23 T	1973	$\alpha$ =?
<sup>253</sup> No <sup>n</sup>	85560 110	1200 110	706 $\mu$ s 24	(25/2 <sup>+</sup> )		11Lo06 TJ	2011	IT ?
<sup>253</sup> No <sup>p</sup>	85800 200	1440 200	627 $\mu$ s 5		13			
<sup>253</sup> Lr	88580# 200#		* 632 ms 46	(7/2 <sup>-</sup> )	13	01He35 TJD	1985	$\alpha$ =90 10; SF=2.6 21; $\beta^+$ =1#
<sup>253</sup> Lr <sup>m</sup>	88610# 230#	30# 100#	* 1.32 s 0.14	(1/2 <sup>-</sup> )	13	09He20 TJD	1985	$\alpha$ =90 10; SF=8 5; $\beta^+$ =1#
<sup>253</sup> Rf	93560# 410#		* 13 ms 5	(7/2) <sup>(+#)</sup>	06	95Ho.B TJ	1997	SF=?; $\alpha$ ?
<sup>253</sup> Rf <sup>m</sup>	93760# 440#	200# 150#	* 52 $\mu$ s 14	(1/2) <sup>(-#)</sup>	06	97He29 J	1995	SF=?; $\alpha$ =5#
* <sup>253</sup> Bk	I : possible identification in 91Kr.A; needs confirmation							
* <sup>253</sup> Es	D : SF=8.7(0.3)e-6% from ENSDF'99 : from $\alpha$ /SF=1.15(0.03)e7 (1965Me02)							
* <sup>253</sup> Fm <sup>m</sup>	E : 211 keV above (7/2 <sup>+</sup> ) level at 130-150 keV							
* <sup>253</sup> Md	T : symmetrized from 6.4(+11.6-3.6)							
* <sup>253</sup> No	T : average 09He23=1.56(0.02) m 09Qi04=1.57(+0.18-0.15) m 67Mi03=95(10) s							
* <sup>253</sup> No	T : and 67Gh01=105(20) s							
* <sup>253</sup> No	J : from 11Lo06 and 10St14							
* <sup>253</sup> No	D : $\epsilon/\epsilon^+$ =0.45(0.03)							
* <sup>253</sup> No <sup>m</sup>	E : average 11An13=167.5(0.5) 10St14=166.7(1.0)							
* <sup>253</sup> No <sup>m</sup>	T : average 09He23=28(3) 07Lo11=31.1(2.1) 73Be33=31.3(4.1);							
* <sup>253</sup> No <sup>m</sup>	T : others 11An13=22.7(0.5) and 10St14=24(2) disagree							
* <sup>253</sup> No <sup>n</sup>	E : greater than 1011 and less than 1380 keV							
* <sup>253</sup> No <sup>n</sup>	T : 11Lo06=706(24) 11An13=627(5) 07Lo11=970(210)							
* <sup>253</sup> No <sup>n</sup>	T : possibly two isomers with 792(43) $\mu$ s and 641(23) $\mu$ s in 11Lo06							
* <sup>253</sup> No <sup>n</sup>	T : possibly two isomers with 650(15) $\mu$ s and 552(15) $\mu$ s in 11An13							
* <sup>253</sup> No <sup>p</sup>	E : ENSDF=1440 + x, x unknown							
* <sup>253</sup> Lr	T : average 09He20=670(60) 01He35=570(+70-60)							
* <sup>253</sup> Lr	D : symmetrized from SF=1.3(+3.0-1.0)%							
* <sup>253</sup> Lr <sup>m</sup>	T : supersedes 01He35=1.49(+0.30-0.21); other 10He11=1.2(+0.7-0.4)							
* <sup>253</sup> Rf	I : the state with ≈1.8 s reported in earlier ENSDF is not confirmed							
* <sup>253</sup> Rf	T : symmetrized from 11(+6-3) I : ENSDF06 reported 253Rf ground-state and m							
* <sup>253</sup> Rf <sup>m</sup>	T : symmetrized from 48(+17-10)							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>254</sup> Bk	84390# 300#				1# m		05			$\beta^-$ ?	
<sup>254</sup> Cf	81341 11				60.5 d	0.2	0 <sup>+</sup>	05	1955	SF≈100; $\alpha=0.31$ 2; $2\beta^-$ ?	
<sup>254</sup> Es	81991 4				275.7 d	0.5	(7 <sup>+</sup> )	05	1954	$\alpha\approx 100$ ; $\epsilon=0.03\%$ ; $\beta^-=1.74e-4$ 8; SF<3e-6	
<sup>254</sup> Es <sup>m</sup>	82075 3	84.2	2.5	AD	39.3 h	0.2	2 <sup>+</sup>	05	1954	$\beta^-=98$ 2; IT<3; $\alpha=0.32$ 1; $\epsilon=0.076$ 7; ... *	
<sup>254</sup> Fm	80902.8 2.4				3.240 h	0.002	0 <sup>+</sup>	05	1954	$\alpha\approx 100$ ; SF=0.0592 3	
<sup>254</sup> Md	83450# 100#			*	10 m	3	0 <sup>-</sup> #	05	1970	$\beta^+\approx 100$ ; $\alpha$ ?	
<sup>254</sup> Md <sup>m</sup>	83500# 140#	50#	100#	*	28 m	8	3 <sup>-</sup> #	05	1970	$\beta^+\approx 100$ ; $\alpha$ ?	
<sup>254</sup> No	84723 10				51.2 s	0.4	0 <sup>+</sup>	05	06He19 T	1966	$\alpha=90$ 1; $\beta^+=10$ 1; SF=0.23 1 *
<sup>254</sup> No <sup>m</sup>	86018 10	1295	2		264.9 ms	1.4	(8 <sup>-</sup> )	05	11Lo06 T	1973	IT>80; SF=0.020 12; $\alpha=0.01$ *
<sup>254</sup> No <sup>n</sup>	87940# 300#	3220#	300#		183.8 $\mu$ s	1.6	(16 <sup>+</sup> )		10He10 ETD	2006	IT=100; SF<0.012 *
<sup>254</sup> Lr	89870# 300#				17.1 s	1.8		05	08An16 TD	1981	$\alpha=72$ 2; $\beta^+=28$ 2; SF ? *
<sup>254</sup> Lr <sup>p</sup>	89940# 310#	60	50	AD							
<sup>254</sup> Lr <sup>d</sup>	90090# 330#	220#	120#								
<sup>254</sup> Rf	93200# 280#				23.2 $\mu$ s	1.0	0 <sup>+</sup>	05	15Da12 T	1997	SF=?; $\alpha<1.5$ *
<sup>254</sup> Rf <sup>m</sup>	94500# 340#	1300#	200#		4.7 $\mu$ s	1.1	(8 <sup>-</sup> )		15Da12 JT	2015	IT=100; SF ? *
<sup>254</sup> Rf <sup>n</sup>	95200# 570#	2000#	500#		247 $\mu$ s	73	(16 <sup>+</sup> )		15Da12 JT	2015	IT=100
* <sup>254</sup> Es <sup>m</sup>	D : ... ; SF<0.045 **										
* <sup>254</sup> No	D : from 10He10 **										
* <sup>254</sup> No <sup>m</sup>	T : average 11Lo06=259(17) 10Cl01=263(2) 10He10=275(7) 06He19=266(2) **										
* <sup>254</sup> No <sup>n</sup>	T : 06Ta19=266(10); other 73Gh03=280(40) **										
* <sup>254</sup> No <sup>n</sup>	T : average 06He19=184(3) 10He10=198(13) 10Cl01=184(2) 06Ta19=171(9) **										
* <sup>254</sup> No <sup>n</sup>	E : 2917(3) + x ; x estimated 300#300; 10Cl01=2930(2) but their level **										
* <sup>254</sup> No <sup>n</sup>	E : scheme is disputed J : from 06He19 **										
* <sup>254</sup> Lr	T : average 08An16=18(2) 01Ga20=13.4(4.2); 85He22=13(+3-2) same group; other **										
* <sup>254</sup> Lr	T : 06Fo02=22(+9-6) D : not used 06Fo02 $\alpha=60(+11-15)\%$ ; $\beta^+=40(+15-11)\%$ **										
* <sup>254</sup> Rf	T : average 15Da12=23.2(1.1) 97He29=23(3); other 08Dr05=29.6(+0.7-0.6) **										
<sup>255</sup> Cf	84810# 200#				85 m	18	(7/2 <sup>+</sup> )	13		1981	$\beta^-=100$ ; SF<0.001#; $\alpha=2e-7$ #
<sup>255</sup> Es	84089 11				39.8 d	1.2	(7/2 <sup>+</sup> )	13		1954	$\beta^-=92.0$ 4; $\alpha=8.0$ 4; SF=0.0041 2
<sup>255</sup> Fm	83800 4				20.07 h	0.07	7/2 <sup>+</sup>	13		1954	$\alpha=100$ ; SF=2.4e-5 10
<sup>255</sup> Md	84843 7				27 m	2	(7/2 <sup>-</sup> )	13		1958	$\beta^+=93$ 1; $\alpha=7$ 1; SF<0.15
<sup>255</sup> Md <sup>p</sup>	84850# 70#	10#	70#				1/2 <sup>-</sup> #	13			
<sup>255</sup> No	86807 15				3.52 m	0.18	(1/2 <sup>+</sup> )	13	11As03 TJ	1967	$\beta^+=70$ 5; $\alpha=30$ 5
<sup>255</sup> No <sup>m</sup>	87020# 100#	210#	100#		1#	s	11/2 <sup>-</sup> #				
<sup>255</sup> No <sup>p</sup>	86910# 70#	100#	70#	Nm			(7/2 <sup>+</sup> )				
<sup>255</sup> Lr	89947 18				31.1 s	1.1	(1/2 <sup>-</sup> )	13	06Ch52 TJ	1971	$\alpha=99.7$ 1; $\beta^+=0.3$ 1; SF<0.1
<sup>255</sup> Lr <sup>m</sup>	89988 19	41	8	AD	2.54 s	0.05	(7/2 <sup>-</sup> )	13	06Ch52 J	2006	IT≈60; $\alpha\approx 40$
<sup>255</sup> Lr <sup>n</sup>	90741 22	794	12		< 1 $\mu$ s		(15/2 <sup>+</sup> )	13		2009	IT=100 *
<sup>255</sup> Lr <sup>p</sup>	91410 22	1463	12		1.70 ms	0.03	(25/2 <sup>+</sup> )	13		2008	IT=100; $\alpha<0.15$ *
<sup>255</sup> Rf	94330# 120#				1.66 s	0.07	(9/2 <sup>-</sup> )	13	15An05 D	1975	SF=45 3; $\alpha=48$ 3; $\beta^+<1$ *
<sup>255</sup> Rf <sup>m</sup>	94480# 120#	150	22	AD	5.0 $\mu$ s	17	(5/2 <sup>+</sup> )	13	15An05 ETJ	2015	IT=100 *
<sup>255</sup> Db	99590# 360#				17 s	0.5		13		1977	$\alpha$ ?; SF≈20 *
* <sup>255</sup> Lr <sup>n</sup>	E : 740.0 keV above 9/2 <sup>+</sup> , which is <30 above 255Lrm **										
* <sup>255</sup> Lr <sup>p</sup>	E : 1408.6 keV above 9/2 <sup>+</sup> , which is <30 above 255Lrm **										
* <sup>255</sup> Rf	T : average 06He27=1.68(0.09) 01He35=1.64(0.11) D : 15An05 SF=45 3; $\alpha=48$ 3 **										
* <sup>255</sup> Db	T : symmetrized from 1.6(+0.6-0.4) **										
<sup>256</sup> Cf	87040# 310#				12.3 m	1.2	0 <sup>+</sup>	99		1980	SF=100; $\alpha=6.2e-7$ #; $2\beta^-$ ?
<sup>256</sup> Es	87190# 100#			*	25.4 m	2.4	(1 <sup>+</sup> , 0 <sup>-</sup> )	99		1981	$\beta^-=100$
<sup>256</sup> Es <sup>m</sup>	87190# 140#	0#	100#	*	7.6 h		(8 <sup>+</sup> )	99		1976	$\beta^-\approx 100$ ; $\beta^-$ -SF=0.002
<sup>256</sup> Fm	85487 6				157.6 m	1.3	0 <sup>+</sup>	99		1955	SF=91.9 3; $\alpha=8.1$ 3
<sup>256</sup> Md	87460# 120#			*	30#	m	7 <sup>-</sup> #				$\beta^+$ ?; $\alpha$ ?; SF ?
<sup>256</sup> Md <sup>m</sup>	87620 70	160#	100#	*	77 m	2	(1 <sup>-</sup> )	99	FGK12b I	1955	$\beta^+=?$ ; $\alpha=9.2$ 7; SF<3 *
<sup>256</sup> Md <sup>p</sup>	87700# 120#	240#	140#				am				
<sup>256</sup> No	87822 8				2.91 s	0.05	0 <sup>+</sup>	99		1963	$\alpha\approx 100$ ; SF=0.53 6; $\epsilon<0.01\%$
<sup>256</sup> Lr	91750 80				27 s	3		99		1965	$\alpha=85$ 10; $\beta^+=15$ 10; SF<0.03
<sup>256</sup> Lr <sup>p</sup>	91980# 90#	230#	40#								
<sup>256</sup> Rf	94222 18				6.67 ms	0.10	0 <sup>+</sup>	14		1975	SF=?; $\alpha=0.32$ 17 *
<sup>256</sup> Rf <sup>m</sup>	95340# 100#	1120#	100#		25 $\mu$ s	2	(5 <sup>-</sup> )	14		2009	IT=100; SF ?
<sup>256</sup> Rf <sup>n</sup>	95620# 100#	1400#	100#		17 $\mu$ s	2	(8 <sup>-</sup> )	14		2009	IT=100; SF ?
<sup>256</sup> Rf <sup>p</sup>	96620# 200#	2400#	200#		27 $\mu$ s	5		14		2009	IT=100; SF ?
<sup>256</sup> Db	100500# 240#				1.7 s	0.4		16	01He35 TD	2001	$\alpha=70$ 11; $\beta^+=36$ 12; SF=? *
* <sup>256</sup> Md <sup>m</sup>	I : Following the Gallagher-Moskowsky rule, this should be the ground-state **										
* <sup>256</sup> Rf	D : other 10St14 SF=97(+2-6)% **										
* <sup>256</sup> Db	T : symmetrized from 01He35=1.6(+0.5-0.3); other 83Og.A=2.6(+1.4-0.8) **										
* <sup>256</sup> Db	D : 01He35 $\beta^+=36(12)\%$ 08Ne01 $\alpha=70(11)\%$ **										

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)		Excitation energy (keV)		Half-life		$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)			
<sup>257</sup> Es	89400#	410#			7.7	d	0.2	7/2 <sup>+</sup> #	13		1987	$\beta^- = 100; \alpha = 4e-4\#$		
<sup>257</sup> Fm	88590	4			100.5	d	0.2	(9/2 <sup>+</sup> )	13		1964	$\alpha \approx 100; SF = 0.210\ 4$		
<sup>257</sup> Md	88993.1	1.6			5.52	h	0.05	(7/2 <sup>-</sup> )	13		1965	$\epsilon = 85\ 3; \alpha = 15\ 3; SF < 1$		
<sup>257</sup> No	90247	7			24.5	s	0.5	(3/2 <sup>+</sup> )	13	02Ho11	D	1967	$\alpha = ?; \beta^+ = 15\ 8$	
<sup>257</sup> No <sup>p</sup>	90550#	110#	300#	110#				9/2 <sup>+</sup> # <i>am</i>						
<sup>257</sup> Lr	92670#	40#			6.0	s	0.4	(1/2 <sup>-</sup> )	13	10St14	T	1971	$\alpha \approx 100; \beta^+ = 0.01\#; SF = 0.001\#$	
<sup>257</sup> Lr <sup>p</sup>	92820#	110#	150#	100#				<i>am</i>	13				*	
<sup>257</sup> Rf	95866	11			4.82	s	0.13	(1/2 <sup>+</sup> )	13	FGK10a	J	1969	$\alpha = ?; \beta^+ = 19.4\ 14; SF = 1.3\ 3$	
<sup>257</sup> Rf <sup>m</sup>	95940	10	73	11	AD			(11/2 <sup>-</sup> )	13	10Be16	T	1997	$\alpha = 88\ 2; \beta^+ = 11\ 2; SF < 1.4$	
<sup>257</sup> Rf <sup>m</sup>	97022	10	1155	11	AD			(21/2 <sup>+</sup> )	13	13Ri07	TJ	2009	IT=100	
<sup>257</sup> Db	100210#	200#			*			(9/2 <sup>+</sup> )	13			1985	$\alpha > 94; SF < 6; \beta^+ = 1\#$	
<sup>257</sup> Db <sup>m</sup>	100350#	230#	140#	110#	*			(1/2 <sup>-</sup> )	99	01He35	J	1985	$\alpha > 87; SF < 13; \beta^+ = 1\#$	
<sup>257</sup> Lr	T : average 10St14=6.3(+0.9-0.7) and 5.8 (0.5)											**		
<sup>257</sup> Lr	T : others not used 97He29=3.3(+0.5-0.4) 97He29=4.3(+1.3-0.8)											**		
<sup>257</sup> Lr	T : 76Be.A=0.646(0.025) 71Es01=0.6(0.1)											**		
<sup>257</sup> Lr	J : feeding in $\epsilon$ decay of 1/2 <sup>+</sup> <sup>257</sup> Rf; and TNN trends for e-o neighbors											**		
<sup>257</sup> Rf	J : favorite $\alpha$ to the 1/2 <sup>+</sup> state at 670 keV D : also 09Qi04 SF=2(1)%											**		
<sup>257</sup> Rf	T : average 10St14=5.5(0.4) 10Be16=4.8(0.2) 09Qi04=4.7(0.3)											**		
<sup>257</sup> Rf	T : 85So03=3.8(0.8) 74Be.A=4.8(0.3) 71Gh03=4.8(0.5)											**		
<sup>257</sup> Rf <sup>m</sup>	E : 97He29=118(4) keV from direct comparison of two $\alpha$ lines											**		
<sup>257</sup> Rf <sup>m</sup>	T : average 10Be16=4.6(0.3) 08Dr05=4.1(+0.7-0.6) 97He29=3.9(0.4)											**		
<sup>257</sup> Rf <sup>m</sup>	T : 09Qi04=4.1(+2.4-1.3) maybe to a 11/2 <sup>-</sup> level in <sup>257</sup> Lr											**		
<sup>257</sup> Rf <sup>m</sup>	E : 1082(4) keV above <sup>257</sup> Rf <sup>m</sup>											**		
<sup>257</sup> Rf <sup>m</sup>	T : 10Be16=134.9 (7.7), reanalyzed in 13Ri07 to 10Be16=110(5)											**		
<sup>258</sup> Es	92700#	400#			3#	m							$\beta^- ?; \alpha ?$	
<sup>258</sup> Fm	90430#	200#			370	$\mu$ s	14	0 <sup>+</sup>	01	86Hu05	T	1971	SF $\approx$ 100; $\alpha ?$	
<sup>258</sup> Md	91687	4			*	51.5	d	0.3	8 <sup>-</sup> #	01	93Mo18	D	1970	$\alpha \approx 100; \beta^+ < 0.0015; \beta^- < 0.0015$
<sup>258</sup> Md <sup>m</sup>	91690#	200#	0#	200#	*	57.0	m	0.9	1 <sup>-</sup> #	01	93Mo18	D	1980	$\epsilon = ?; SF < 20; \beta^- < 10\#; \alpha < 1.2$
<sup>258</sup> No	91480#	100#						1.2	ms	0.2	0 <sup>+</sup>	01	1989	SF $\approx$ 100; $\alpha = 0.001\#; 2\beta^+ ?$
<sup>258</sup> Lr	94780#	100#			3.6	s	0.4		01	14Ha04	TD	1971	$\alpha = ?; \beta^+ = 2.6\ 18$	
<sup>258</sup> Lr <sup>p</sup>	95020#	140#	240#	100#				<i>am</i>					*	
<sup>258</sup> Rf	96340	30			13.8	ms	0.9	0 <sup>+</sup>	01	08Ga08	T	1969	SF=87 2; $\alpha = 13\ 2$	
<sup>258</sup> Db	101800#	310#			*	4.5	s	0.4	01	09He20	T	1981	$\alpha = 63\ 6; \beta^+ = 37\ 6; SF < 1\#$	
<sup>258</sup> Db <sup>m</sup>	101860#	320#	60#	100#	*	1.9	s	0.5	01	09He20	T	1985	$\beta^+ \approx 100; IT ?$	
<sup>258</sup> Sg	105240#	410#			2.7	ms	0.5	0 <sup>+</sup>	01	09Fo02	T	1997	SF=?; $\alpha < 20$	
<sup>258</sup> Fm	T : average 86Hu05=360(20) 71Hu03=380(20) (all 1 $\sigma$ ) ENSDF gives 3 $\sigma$											**		
<sup>258</sup> Md	D : derived from: "the sum of SF, $\epsilon$ and $\beta^-$ decay branches < 0.003%" in											**		
<sup>258</sup> Md	D : 93Mo18 and T(SF)>150000 y, from 86Lo16, thus SF<1e-4%#											**		
<sup>258</sup> Md <sup>m</sup>	D : SF<20% derived from 93Mo18 "the sum of SF and $\beta^-$ decay branches < 30%"											**		
<sup>258</sup> Lr	T : symmetrized from 14Ha04=3.54(+0.46-0.36)											**		
<sup>258</sup> Rf	T : average 08Ga08=14.7(+1.2-1.0) 85So03=13(3) 69Gh01=11(2)											**		
<sup>258</sup> Db	T : average 09He20=4.3(0.5) 06Fo02=4.8(+1.0-0.8) 01Ga20=4.3(1.1) and											**		
<sup>258</sup> Db	T : 85He22=4.4(+0.9-0.6)											**		
<sup>258</sup> Db	D : average $\beta^+ 06Fo02=39(+11-9)\%$ 85He22=33(+9-5)%											**		
<sup>258</sup> Sg	T : symmetrized from 09Fo02=2.6(+0.6-0.4); combining with earlier work											**		
<sup>259</sup> Fm	93700#	280#			1.5	s	0.2	3/2 <sup>+</sup> #	13			1980	SF=100	
<sup>259</sup> Md	93620#	200#			1.60	h	0.06	7/2 <sup>-</sup> #	13			1982	SF=?; $\alpha < 1.3$	
<sup>259</sup> No	94079	7			58	m	5	(9/2 <sup>+</sup> )	13	13As02	J	1973	$\alpha = 75\ 4; \epsilon = 25\ 4; SF < 10$	
<sup>259</sup> No <sup>p</sup>	94310#	150#	230#	150#										
<sup>259</sup> Lr	95850#	70#			6.2	s	0.3	1/2 <sup>-</sup> #	13			1971	$\alpha = 78\ 2; SF = 22\ 2; \beta^+ = 0.6\#$	
<sup>259</sup> Lr <sup>p</sup>	96200#	170#	350#	150#										
<sup>259</sup> Rf	98360#	70#			2.63	s	0.26	7/2 <sup>+</sup> #	13	08Ga08	T	1969	$\alpha = 92\ 2; SF = 8\ 2; \beta^+ = 0.3\#$	
<sup>259</sup> Rf <sup>p</sup>	98430#	100#	60	70	Nm			(3/2 <sup>+</sup> )					*	
<sup>259</sup> Rf <sup>p</sup>	98570#	110#	210	90	Nm			(9/2 <sup>+</sup> )						
<sup>259</sup> Db	101990	50			510	ms	160	9/2 <sup>+</sup> #	13	01Ga20	D	2001	$\alpha = 100$	
<sup>259</sup> Sg	106520#	120#			402	ms	56	(11/2 <sup>-</sup> )	13	15An05	TJD	1985	$\alpha = 97\ 1; SF < 3; \epsilon < 1$	
<sup>259</sup> Sg <sup>m</sup>	106610#	120#	87	22	AD			(1/2 <sup>+</sup> )		15An05	TJD		$\alpha = 97\ 1; SF > 3; \epsilon < 1$	
<sup>259</sup> Rf	T : average 08Ga08=2.5(+0.4-0.3) 94Gr08=1.7(+0.8-0.5)											**		
<sup>259</sup> Sg	D : SF=3(1)% assumed from shorter-lived isomeric state											**		
<sup>259</sup> Sg <sup>m</sup>	D : SF=3(1)% assumed from this state											**		

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>260</sup> Fm	95770# 440#	EU	1# m	0 <sup>+</sup>				SF ? *
<sup>260</sup> Md	96550# 320#		27.8 d 0.8		99	92Lo.B TD	1989	SF=?; $\alpha < 5$ ; $\epsilon < 5$ ; $\beta^- < 3.5$ *
<sup>260</sup> No	95610# 200#		106 ms 8	0 <sup>+</sup>	99		1985	SF=100 *
<sup>260</sup> Lr	98280# 120#		3.0 m 0.5		99		1971	$\alpha = 80$ 20; $\beta^+ = 20$ 20 *
<sup>260</sup> Rf	99150# 200#		21 ms 1	0 <sup>+</sup>	99		1985	SF=?; $\alpha = 2$ %; $\epsilon = 0.01$ % *
<sup>260</sup> Db	103670# 90#		1.52 s 0.13		99		1970	$\alpha > 90.4$ 6; SF < 9.6 6; $\beta^+ < 2.5$ *
<sup>260</sup> Db <sup>p</sup>	103870# 180#	200# 150#						
<sup>260</sup> Sg	106548 21		4.95 ms 0.33	0 <sup>+</sup>	99	09He20 T	1984	SF=60 30; $\alpha = 40$ 30 *
<sup>260</sup> Bh	113320# 250#		41 ms 14		16	08Ne01 TD	2008	$\alpha \approx 100$ ; $\beta^+ ?$ ; SF ? *
* <sup>260</sup> Fm	I : half-life $\approx 4$ ms and SF=100 mode were reported in the 92Lo.B internal **							
* <sup>260</sup> Fm	I : report. Not confirmed in subsequent experiment by same group (97Lo.A) **							
* <sup>260</sup> Fm	I : Discovery of this nuclide is considered unproven **							
* <sup>260</sup> Md	T : supersedes 86Hu01=31.8(0.5) of same group **							
* <sup>260</sup> Rf	T : also 08Ga08=22.2(+3.0-2.4) 08Go.A=21(+7.3,-4.3) **							
* <sup>260</sup> Db	T : also 04Mo26=1.5(+0.8-0.4) 04Ga29=0.89(+0.79-0.35) **							
* <sup>260</sup> Sg	T : supersedes 85Mu11=3.6(+0.9-0.6) **							
* <sup>260</sup> Sg	D : symmetrized from SF=50(+30-20)% and $\alpha = 50$ (+20-30)% **							
* <sup>260</sup> Bh	T : symmetrized from 08Ne01=35(+19-9) **							
<sup>261</sup> Md	98580# 510#		40# m	7/2 <sup>-</sup> #				$\alpha ?$ *
<sup>261</sup> No	98460# 200#		3# h	3/2 <sup>+</sup> #				$\alpha ?$ *
<sup>261</sup> Lr	99560# 200#		39 m 12		99		1987	SF=?; $\alpha ?$ *
<sup>261</sup> Rf	101320 50		* & 2.2 s 0.3	3/2 <sup>+</sup> #	15	11Ha13 TD	1970	SF=73 6; $\alpha = 27$ 6 *
<sup>261</sup> Rf <sup>m</sup>	101390# 110#	70# 100#	* & 20 s 4	9/2 <sup>+</sup> #	15	13Mu08 T	1970	$\alpha = ?$ ; $\beta^+ < 15$ ; SF < 10 *
<sup>261</sup> Rf <sup>p</sup>	101620# 110#	300# 100#						
<sup>261</sup> Db	104310# 110#		4.7 s 1.0		99	13Su04 TD	1970	SF=73 11; $\alpha = ?$ *
<sup>261</sup> Db <sup>p</sup>	104610# 230#	300# 200#						
<sup>261</sup> Sg	108005 18		183 ms 5	(3/2 <sup>+</sup> )	99	10St14 TJD	1984	$\alpha = 98.1$ 4; $\beta^+ = 1.3$ 3; SF=0.6 2 *
<sup>261</sup> Sg <sup>m</sup>	108110# 50#	100# 50#	9.3 $\mu$ s 1.8	(11/2 <sup>-</sup> )	99	10Be16 TJ	2010	IT=100 *
<sup>261</sup> Bh	113130# 210#		12.8 ms 3.2	(5/2 <sup>-</sup> )	99	10He11 TJD	1989	$\alpha = 95$ 5; SF < 5 *
* <sup>261</sup> Rf	T : average 12Ha05=2.6(+0.7-0.5) 11Ha13=1.9(0.4) 08Go.A=2.2(+0.9-0.5) **							
* <sup>261</sup> Rf	T : others 08Dv02=3(1) 08Mo09 2 events at 2.97 and 8.3s 02Ho11=4.2(+3.4-1.3) **							
* <sup>261</sup> Rf	T : 13Mu08=3.9(3.0) using SF events **							
* <sup>261</sup> Rf	D : SF others 12Ha05=82(9)%, 11Ha13=73(6)%, 08Dv02=91% for 11 events **							
* <sup>261</sup> Rf <sup>m</sup>	T : symmetrized from 19(+5-3); others 08Dv02=20(+110-10) 02Ho11=78(+11-6) **							
* <sup>261</sup> Db	T : average 13Su04=4.7(+3.6-1.4) 10St14=4.1(+1.4-0.8) **							
* <sup>261</sup> Db	D : observed 11 SF and 4 $\alpha$ decays; uncertainty evaluated by NUBASE **							
* <sup>261</sup> Sg	T : average 10St14=184(5) 10Be16=178(14) **							
* <sup>261</sup> Sg <sup>m</sup>	T : symmetrized from 9.0(+2.0-1.5) **							
* <sup>261</sup> Bh	T : symmetrized from 10He11=11.8(+3.9-2.4); others not used 06Fo02=10(+14-5) **							
* <sup>261</sup> Bh	T : and 08Ne08=6.7(+3.8-1.8) **							
<sup>262</sup> Md	101630# 500#		3# m					SF ?; $\alpha ?$ *
<sup>262</sup> No	100100# 360#		5 ms	0 <sup>+</sup>			1989	SF $\approx 100$ ; $\alpha ?$ *
<sup>262</sup> Lr	102100# 200#		4 h				1987	$\beta^+ = ?$ ; SF < 10; $\alpha ?$ *
<sup>262</sup> Rf	102390# 220#		* 250 ms 100	0 <sup>+</sup>	01	08Go.A TD	1985	SF $\approx 100$ *
<sup>262</sup> Rf <sup>m</sup>	103390# 460#	1000# 400#	* 47 ms 5	high		96La11 I	1978	SF=100 *
<sup>262</sup> Db	106250# 140#		34 s 4		01	14Ha04 TD	1971	SF=52 4; $\alpha = ?$ ; $\beta^+ = 3$ ## *
<sup>262</sup> Db <sup>p</sup>	106300# 160#	50# 70#						$\alpha ?$ *
<sup>262</sup> Sg	108370 40		10.9 ms 2.3	0 <sup>+</sup>	01	06Gr24 TD	2001	SF $\approx 100$ ; $\alpha ?$ *
<sup>262</sup> Sg <sup>p</sup>	109220 90	860 90 AD						
<sup>262</sup> Bh	114540# 310#		84 ms 11		01	09He20 T	1981	$\alpha = ?$ ; SF < 20 *
<sup>262</sup> Bh <sup>m</sup>	114760# 310#	210 50 AD	9.5 ms 1.6		01	06Fo02 T	1981	$\alpha = ?$ ; SF < 10 *
* <sup>262</sup> Rf	T : symmetrized from 08Go.A=210(+128-58) ms; 7 SF events **							
* <sup>262</sup> Rf	T : conflicting 96La11=2.1(0.2) 94La22=1.2(+1.0-0.5) **							
* <sup>262</sup> Rf	T : 11Ha13 and 08Go.A suggest these activities belong to <sup>261</sup> Rf **							
* <sup>262</sup> Rf	D : this suggestion contradicts 96La11 $\alpha < 0.8$ ; not adopted by NUBASE **							
* <sup>262</sup> Rf <sup>m</sup>	I : assigned in 96La11 to K-isomeric state T : 85So03=47(5) **							
* <sup>262</sup> Db	T : symmetrized from 14Ha04=33.8(+4.4-3.5) **							
* <sup>262</sup> Sg	T : 06Gr24=15(+5-3) 01Ho06=6.9(+3.8-1.8) D : no $\alpha$ observed $\alpha < 16$ % **							
* <sup>262</sup> Bh	T : average 09He20=83(14) 06Fo02=84(+21-16) **							
* <sup>262</sup> Bh	T : other 08Ne08(10 events)=120(+55-29) not used **							
* <sup>262</sup> Bh <sup>m</sup>	T : 06Fo02=9.6(+3.6-2.4) 97Ho14(11 events)=12.2(+5.5-2.8) 89Mu09=8.0(2.1) **							
* <sup>262</sup> Bh <sup>m</sup>	T : also 09He20=22(4) 08Ne08(4 events)=16(+14-5) not used **							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>263</sup> No	103130#	490#	20# m					$\alpha$ ?; SF ?	
<sup>263</sup> Lr	103730#	280#	5# h					$\alpha$ ?	
<sup>263</sup> Rf	104760#	150#	11 m	3	3/2 <sup>+</sup> #	99	93Gr.C TD	2003 SF=?; $\alpha$ =30	
<sup>263</sup> Rf <sup>p</sup>	105060#	250#	300# 200#						
<sup>263</sup> Db	107110#	170#	29 s	9		99	92Kr01 D	1992 SF=56 14; $\alpha$ =?; $\beta^+$ =6.9 16	
<sup>263</sup> Db <sup>p</sup>	107370#	260#	200#						
<sup>263</sup> Sg	110190#	100#	940 ms	140	7/2 <sup>+</sup> #	99	06Gr24 TD	1974 $\alpha$ =87 8; SF=13 8	
<sup>263</sup> Sg <sup>m</sup>	110240#	100#	51 19 Nm *	420 ms	100	3/2 <sup>+</sup> #	99	04Fo08 T	1995 $\alpha$ =?; IT ?
<sup>263</sup> Sg <sup>p</sup>	110290#	100#	100 30 AD						
<sup>263</sup> Bh	114500#	310#	200# ms			99		$\alpha$ ?	
<sup>263</sup> Hs	119680#	130#	760 $\mu$ s	40	3/2 <sup>+</sup> #	99	09Dr02 TD	2009 $\alpha$ =?; SF<8.4	
<sup>263</sup> Hs <sup>m</sup>	120000#	130#	320 70 AD	760 $\mu$ s	40	11/2 <sup>-</sup> #	09Dr02 TD	2009 $\alpha$ =?; SF ?	
* <sup>263</sup> Rf	T : average 03Kr20=24(+19-7) m 93Gr.C=500(+300-200) s 92Cz.A=600(+300-200) s								
* <sup>263</sup> Rf	T : also one SF event 08Dv02=8(+40-4) s								
* <sup>263</sup> Db	D : SF from 92Kr01=57(+13-15)%; $\beta^+$ average 03Kr20=3(+4-1)% 93Gr.C=8(2)%								
* <sup>263</sup> Db	T : Possibly a candidate for the 54(+98-21) s SF decay observed in 98Ik02								
* <sup>263</sup> Db	T : symmetrized from 27(+10-7)								
* <sup>263</sup> Sg	T : average 06Gr24=820(+370-190) 94Gr08=553(+336-152) 74Gh04=900(200); all								
* <sup>263</sup> Sg	T : produced via direct production mechanisms								
* <sup>263</sup> Sg <sup>m</sup>	T : average 04Fo08=290(+170-90) 04Mo40=549(+300-143) ms 03Gi05=222(+404-87)								
* <sup>263</sup> Sg <sup>m</sup>	T : and 98Ho13=310(+160-80) ms; all produced via $\alpha$ decay of parent								
* <sup>263</sup> Sg <sup>m</sup>	T : also 10Ni14 at $\tau$ =702ms via $\alpha$ -decay of parent, but with low energy								
* <sup>263</sup> Hs	T : symmetrized from 740(+48-21) 6 events for both states (low statistics)								
* <sup>263</sup> Hs	D : 09Dr02 no SF observed								
<sup>264</sup> No	105010#	590#	1# m		0 <sup>+</sup>			$\alpha$ ?; SF ?	
<sup>264</sup> Lr	106380#	440#	10# h					$\alpha$ ?; SF ?	
<sup>264</sup> Rf	106080#	360#	1# h		0 <sup>+</sup>			$\alpha$ ?	
<sup>264</sup> Db	109360#	240#	3# m					$\alpha$ ?	
<sup>264</sup> Sg	110780#	280#	47 ms	20	0 <sup>+</sup>	06		2006 SF $\approx$ 100; $\alpha$ ?	
<sup>264</sup> Bh	116060#	180#	1.07 s	0.21		99	04Mo26 TD	1995 $\alpha$ =86; SF=14; $\beta^+$ ?	
<sup>264</sup> Bh <sup>p</sup>	116290#	230#	230# 150#						
<sup>264</sup> Hs	119563	29	540 $\mu$ s	300	0 <sup>+</sup>	99	95Ho.B T	1986 $\alpha$ $\approx$ 50; SF $\approx$ 50	
* <sup>264</sup> Sg	T : symmetrized from 37(+27-11); also 10Ni14(1 event)=86.4 ms								
* <sup>264</sup> Sg	D : no $\alpha$ observed $\alpha$ <36%								
* <sup>264</sup> Bh	T : average 04Mo26=0.9(+0.3-0.2) 04Ga29=1.17(+0.88-0.44) and								
* <sup>264</sup> Bh	T : 02Ho11=1.02(+0.69-0.29)								
* <sup>264</sup> Hs	T : 95Ho.B (2 events 76 $\mu$ s and 825 $\mu$ s) 87Mu15 (1 event 80 $\mu$ s); average of the								
* <sup>264</sup> Hs	T : 3 events: 327(+448-120) $\mu$ s, see 84Sc13								
<sup>265</sup> Lr	108230#	550#	10# h					$\alpha$ ?; SF ?	
<sup>265</sup> Rf	108690#	360#	1.6 m	0.8	3/2 <sup>+</sup> #	15	15Ut02 T	2010 SF $\approx$ 100; $\alpha$ ?	
<sup>265</sup> Db	110480#	220#	15# m					$\alpha$ ?	
<sup>265</sup> Sg	112790#	120#	& 9.2 s	1.6	9/2 <sup>+</sup> #	15	12Ha05 T	1994 $\alpha$ >50; SF ?	
<sup>265</sup> Sg <sup>m</sup>	112860#	120#	60# 160#	& 16.4 s	2.4	3/2 <sup>+</sup> #	15	13Su04 T	1994 $\alpha$ >65 16; SF ?
<sup>265</sup> Bh	116420#	230#	1.19 s	0.52		99	04Ga29 TD	2004 $\alpha$ =?	
<sup>265</sup> Hs	120900	24	1.96 ms	0.16	3/2 <sup>+</sup> #	99	09He20 T	1984 $\alpha$ $\approx$ 100; SF<1	
<sup>265</sup> Hs <sup>m</sup>	121130	24	229 22 AD	360 $\mu$ s	150	9/2 <sup>+</sup> #	99	09He20 T	1995 $\alpha$ $\approx$ 100; IT ?
<sup>265</sup> Mt	126680#	450#	2# ms					$\alpha$ ?	
* <sup>265</sup> Rf	T : symmetrized from 15Ut02=1.0(+1.2-0.3)								
* <sup>265</sup> Sg	T : average 12Ha05=8.5(+2.6,-1.6) 08Du09=8.9(+2.7-1.9)								
* <sup>265</sup> Sg <sup>m</sup>	T : average 13Su04=20(+15-6) 12Ha05=14.4(+3.7,-2.5) 08Du09=16.2(+4.7-3.5)								
* <sup>265</sup> Bh	T : symmetrized from 0.94(+0.70-0.31)								
* <sup>265</sup> Hs	T : average 09He20=1.9(0.2) 99He11=2.0(+0.3-0.2)								
* <sup>265</sup> Hs <sup>m</sup>	T : symmetrized from 300(+200-100); other 99He11=750(+170-120)								
<sup>266</sup> Lr	111620#	580#	21 h	14		14		2014 SF=100	
<sup>266</sup> Rf	110080#	470#	4# h		0 <sup>+</sup>			$\alpha$ ?; SF ?	
<sup>266</sup> Db	112740#	280#	80 m	70		07	07Og02 T	2007 $\alpha$ ?; SF ?; $\beta^+$ ?	
<sup>266</sup> Sg	113620#	250#	390 ms	110	0 <sup>+</sup>	05	13Og03 TD	2006 SF=100	
<sup>266</sup> Bh	118100#	160#	2.5 s	1.6		05	08Mo09 T	2000 $\alpha$ $\approx$ 100; $\beta^+$ ?; SF ?	
<sup>266</sup> Hs	121140	40	3.02 ms	0.54	0 <sup>+</sup>	05	11Ac.A T	2001 $\alpha$ =?; SF $\approx$ 1.4#	
<sup>266</sup> Hs <sup>m</sup>	122240	80	1100 70 AD	280 ms	220	9 <sup>-</sup> #	11Ac.A T	2011 $\alpha$ =?	

... A-group is continued on next page ...

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
... A-group continued ...									
<sup>266</sup> Mt	127960# 310#		1.2 ms 0.4		05	97Ho14	T	1982	$\alpha$ ?; SF<5.5
<sup>266</sup> Mt <sup>m</sup>	129100# 310#	1140 80 AD	6 ms 3			97Ho14	TD	1984	$\alpha$ =100
* <sup>266</sup> Lr	T: symmetrized from 14Kh04=11(+21-5)								
* <sup>266</sup> Db	T: one event at 31.74 m, yields 22(+105-10), see 84Sc13								
* <sup>266</sup> Sg	T: average 13Og03=280(+190-80) ms 08Dv02=360(+250-100) ms								
* <sup>266</sup> Sg	T: 08Dv02 supersedes 06Dv01=444(+444-148)								
* <sup>266</sup> Sg	I: 98Tu01=21(+20-12) s 94La22=10-30 s with 18%< $\alpha$ <50% 50%<SF<82% re-assigned								
* <sup>266</sup> Sg	I: to <sup>265</sup> Sg, see 08Dv02; 10Gr04 one SF event after 23 ms, not trusted								
* <sup>266</sup> Bh	T: 2 events at 2.469 and 1.31 s; other 06Qi03=0.66(+0.59-0.26)								
* <sup>266</sup> Hs	T: average 11Ac.A=2.97(+0.78-0.51) 01Ho06=2.3(+1.3-0.6)								
* <sup>266</sup> Hs <sup>m</sup>	T: symmetrized from 11Ac.A=74(+354-34); the possibility in 01Ho06 that								
* <sup>266</sup> Hs <sup>m</sup>	T: 01Ho06=6.3(+8.6-2.3) is ruled out by the 11Ac.A result								
* <sup>266</sup> Mt	T: 10 events yielding 1.01(+0.47-0.24), see 84Sc13								
* <sup>266</sup> Mt <sup>m</sup>	T: 3 events at 7.8, 2.0 and 5.0 yield 3.4(+4.7-1.3), see 84Sc13								
<sup>267</sup> Rf	113440# 580#		2.5 h 1.5		05	06Og05	TD	2004	SF=100
<sup>267</sup> Rf <sup>p</sup>	113660# 580#	220# 100#							
<sup>267</sup> Db	114070# 410#		100 m 60		05	13Ru11	T	2004	SF=100
<sup>267</sup> Sg	115810# 260#		1.8 m 0.7			08Dv02	TD	2008	SF=83; $\alpha$ =17
<sup>267</sup> Sg <sup>p</sup>	115880# 280#	70# 100#							
<sup>267</sup> Bh	118770# 260#		22 s 10		05			2000	$\alpha$ =100
<sup>267</sup> Hs	122650# 100#		55 ms 11	5/2 <sup>+</sup> #	05			1995	$\alpha$ >80; SF?
<sup>267</sup> Hs <sup>m</sup>	122690# 100#	39 24 AD	990 $\mu$ s 90		05	04Fo08	TD	2004	$\alpha$ ?; IT?
<sup>267</sup> Mt	127790# 500#		10# ms						$\alpha$ ?
<sup>267</sup> Ds	133880# 140#		10 $\mu$ s 8	3/2 <sup>+</sup> #	05	95Gh04	T	1995	$\alpha$ =100
* <sup>267</sup> Rf	T: symmetrized from 1.3(+2.3-0.5); supersedes 04Og12 one event at 2.3 h								
* <sup>267</sup> Db	T: 13Ru11 one event at 30.6 m and 04Og03 one event at 73								
* <sup>267</sup> Sg	T: symmetrized from 80(+60-20) s; other 99Og.B=19 ms not trusted								
* <sup>267</sup> Bh	T: symmetrized from 00Wi15=17(+14-6)								
* <sup>267</sup> Hs	T: symmetrized from 52(+13-8)								
* <sup>267</sup> Hs <sup>m</sup>	T: 04Fo08(2 events)=940(+120-45); other not trusted 04Mo40 (1 event)=803 ms								
* <sup>267</sup> Ds	T: one single event, $\tau$ =4 $\mu$ s, thus $T$ =2.8(+13.0-1.3), see 84Sc13								
<sup>268</sup> Rf	115480# 660#		1# h	0 <sup>+</sup>					$\alpha$ ?; SF?
<sup>268</sup> Db	117060# 530#		29 h 4		05	13Ru11	T	2004	SF $\approx$ 100; $\beta$ <sup>+</sup> ?
<sup>268</sup> Db <sup>p</sup>	117210# 530#	150 70							
<sup>268</sup> Sg	116800# 470#		2# m	0 <sup>+</sup>					$\alpha$ ?; SF?
<sup>268</sup> Bh	120810# 380#		25# s						$\alpha$ ?; SF?
<sup>268</sup> Hs	122830# 280#		1.42 s 1.13	0 <sup>+</sup>		10Ni14	TD	2010	$\alpha$ $\approx$ 100
<sup>268</sup> Mt	129150# 230#		27 ms 6	5 <sup>+</sup> #, 6 <sup>+</sup> #	05	04Mo26	T	1995	$\alpha$ =100
<sup>268</sup> Ds	133650# 300#		100# $\mu$ s	0 <sup>+</sup>					$\alpha$ ?
* <sup>268</sup> Db	T: average 13Ru11=26(+7-5) 13Og01=25.9(+6.2-4.2) 07St18=28(+11-4)								
* <sup>268</sup> Db	T: 13Og01 supersedes 12Og02=27.9(+7.8-5.0) 05Og02=29(+9-6) 04Og03=16(+19-6)								
* <sup>268</sup> Hs	T: symmetrized from 0.38(+1.8-0.17)								
* <sup>268</sup> Mt	T: mean lifetime of 14 events in 04Mo26=30 ms and 6 events in 02Ho11=60 ms								
<sup>269</sup> Db	119150# 620#		3# h						$\alpha$ ?; SF?
<sup>269</sup> Sg	119760# 360#		5 m 3		15	15Ut02	T	2010	$\alpha$ $\approx$ 100; SF?
<sup>269</sup> Bh	121480# 370#		1# m						$\alpha$ ?
<sup>269</sup> Hs	124560# 120#		16 s 6	9/2 <sup>+</sup> #	05	13Su04	T	1996	$\alpha$ =100
<sup>269</sup> Mt	129370# 460#		100# ms						$\alpha$ ?
<sup>269</sup> Ds	134830 30		230 $\mu$ s 110	9/2 <sup>+</sup> #	05	95Ho03	T	1995	$\alpha$ =100
* <sup>269</sup> Sg	T: symmetrized from 15Ut02=3.1(+3.7-1.1)								
* <sup>269</sup> Hs	T: average 13Su04=12(+9-4) and 02Ho11 2 events at 19.7s 22.0s, see 84Sc13								
* <sup>269</sup> Ds	T: symmetrized from 170(+160-60)								
<sup>270</sup> Db	122310# 620#		2.0 h 1.3		10	14Kh04	TD	2010	SF=100
<sup>270</sup> Sg	121490# 560#		3# m	0 <sup>+</sup>					$\alpha$ ?; SF?
<sup>270</sup> Bh	124230# 290#		3.8 m 3.0		07	07Og02	TD	2007	$\alpha$ =100
<sup>270</sup> Bh <sup>p</sup>	124920# 350#	690# 200#							
<sup>270</sup> Hs	125110# 250#		9 s 4	0 <sup>+</sup>	05	13Og03	T	2003	$\alpha$ =100; SF?
<sup>270</sup> Mt	130710# 170#		6.3 ms 1.5		05			2004	$\alpha$ $\approx$ 100
... A-group is continued on next page ...									

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
... A-group continued ...								
<sup>270</sup> Ds	134680	50	205 $\mu$ s	48	0 <sup>+</sup>	05 11Ac.A	T 2001	$\alpha \approx 100$ ; SF<0.2
<sup>270</sup> Ds <sup>m</sup>	136070	60	1390	60	AD	10 ms 6	(10) <sup>(-#)</sup>	05 2001 $\alpha=?$ ; IT?
* <sup>270</sup> Db	T: symmetrized from 14Kh04=1.0(+1.9-0.4); other not used 13Og04=17(+15-6)							
* <sup>270</sup> Bh	T: symmetrized from 61(+292-28)s							
* <sup>270</sup> Hs	T: symmetrized from 13Og03=7.6(+4.9-2.2); other estimated 03Tu05=3.6(+0.8-1.4)							
* <sup>270</sup> Mt	T: symmetrized from 5.0(+2.4-0.3)							
* <sup>270</sup> Ds	T: average 11Ac.A=200(+70-40) 01Ho06=100(+140-40)							
* <sup>270</sup> Ds <sup>m</sup>	T: symmetrized from 6.0(+8.2-2.2)							
<sup>271</sup> Sg	124760#	590#	3.1 m	1.6		06 06Og05	TD 2004	$\alpha=70$ ; SF=30
<sup>271</sup> Bh	125920#	420#	10 m	8		05 13Ru11	TD	$\alpha=100$
<sup>271</sup> Hs	127740#	280#	10# s				2008	$\alpha?$ ; SF?
<sup>271</sup> Mt	131100#	330#	400# ms					$\alpha?$
<sup>271</sup> Ds	135950#	100#	90 ms	40	13/2 <sup>-#</sup>	05	1998	$\alpha=100$
<sup>271</sup> Ds <sup>m</sup>	136020#	100#	68	27	AD	* & 1.7 ms 0.4	9/2 <sup>+#</sup>	05 1995 $\alpha=100$
* <sup>271</sup> Sg	T: symmetrized from 1.9(+2.4-0.6); supersedes 04Og12=2.4(4.3-1.0) $\alpha=50$ ; SF=50							
* <sup>271</sup> Bh	T: 13Ru11 one event at 2.6 m							
* <sup>271</sup> Ds	T: symmetrized from 69(+56-21)							
* <sup>271</sup> Ds <sup>m</sup>	T: symmetrized from 1.63(+0.44-0.29)							
<sup>272</sup> Sg	126580#	730#	4# m		0 <sup>+</sup>			$\alpha?$ ; SF?
<sup>272</sup> Bh	128790#	530#	11.3 s	1.8		05 13Ru11	T 2004	$\alpha \approx 100$
<sup>272</sup> Hs	129010#	510#	10# s		0 <sup>+</sup>			$\alpha?$ ; SF?
<sup>272</sup> Mt	133580#	490#	400# ms					$\alpha?$ ; SF?
<sup>272</sup> Ds	136020#	410#	200# ms		0 <sup>+</sup>			SF?
<sup>272</sup> Rg	142770#	230#	4.5 ms	1.0	5 <sup>+#</sup> , 6 <sup>+#</sup>	05 04Mo26	T 1995	$\alpha=100$
* <sup>272</sup> Bh	T: average 13Ru11=9.2(+3.1-1.8) 13Og01=12.0(+3.1-2.1)							
* <sup>272</sup> Bh	T: 13Og01 supersedes 12Og02=8.2(+2.5-1.6)s 04Og03=9.8(+11.7-3.5)s							
* <sup>272</sup> Rg	T: mean lifetime of 14 events in 04Mo26=5.5 ms and 6 events in 02Ho11=2.3							
<sup>273</sup> Sg	130020#	500#	5# m					SF?
<sup>273</sup> Bh	130630#	690#	1# m					$\alpha?$ ; SF?
<sup>273</sup> Hs	131890#	370#	1060 ms	500	3/2 <sup>+#</sup>	15 15Ut02	T 2010	$\alpha \approx 100$
<sup>273</sup> Hs <sup>p</sup>	132000#	380#	110#	100#				$\alpha?$ ; SF?
<sup>273</sup> Mt	134710#	420#	800# ms					$\alpha?$ ; SF?
<sup>273</sup> Ds	138360#	130#	240 $\mu$ s	80	13/2 <sup>-#</sup>	05 13Su04	T 1996	$\alpha=100$
<sup>273</sup> Ds <sup>m</sup>	138560#	130#	198	20	EU	120 ms	3/2 <sup>+#</sup>	05 1996 $\alpha=100$
<sup>273</sup> Rg	142700#	530#	2# ms					$\alpha?$
* <sup>273</sup> Hs	T: symmetrized from 15Ut02=760(+710-240)							
* <sup>273</sup> Ds	T: average 13Su04=190(+140-60) ENSDF=170(+170-60) for 4 events:							
* <sup>273</sup> Ds	T: 08Mo09 2 events at 520 and 40 $\mu$ s; 02Ho11 at 310; 96Ho13 at 110 $\mu$ s							
<sup>274</sup> Bh	133680#	620#	60 s	30		10 14Kh04	TD 2010	$\alpha=100$
<sup>274</sup> Hs	133490#	590#	500# ms		0 <sup>+</sup>			$\alpha?$ ; SF?
<sup>274</sup> Mt	137250#	350#	850 ms	540		07 07Og02	TD 2007	$\alpha=100$
<sup>274</sup> Ds	139200#	390#	10# ms		0 <sup>+</sup>			$\alpha?$ ; SF?
<sup>274</sup> Rg	144610#	180#	29 ms	18		05 08Mo09	TD 2004	$\alpha \approx 100$
* <sup>274</sup> Bh	T: average 14Kh04=30(+54-12) 13Og04=54(+65-19)s							
* <sup>274</sup> Mt	T: symmetrized from 440(+810-170)ms							
* <sup>274</sup> Rg	T: 2 events at 9.26 and 34.3 ms							
<sup>275</sup> Bh	135690#	600#	5# m					SF?
<sup>275</sup> Hs	136620#	590#	290 ms	150		05 06Og05	TD 2004	$\alpha=100$
<sup>275</sup> Hs <sup>p</sup>	136860#	600#	240#	100#				
<sup>275</sup> Mt	138830#	420#	117 ms	74		05 13Ru11	T 2004	$\alpha=100$
<sup>275</sup> Ds	141570#	410#	10# ms					$\alpha?$ ; SF?
<sup>275</sup> Rg	145300#	520#	5# ms					$\alpha?$
* <sup>275</sup> Hs	T: symmetrized from 190(+220-70)ms; supersedes 04Og12=150(+270-60)							
* <sup>275</sup> Mt	T: 13Ru11 one event at 51.3 ms and 04Og03 one event at 9.7							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)				Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>276</sup> Hs	138290# 750#					100# ms	0 <sup>+</sup>				$\alpha$ ?; SF ?
<sup>276</sup> Mt	141320# 530#				*	630 ms 100		05	13Ru11 T	2004	$\alpha=100$ *
<sup>276</sup> Mt <sup>m</sup>	141570# 540#	250	80	AD	*	10 s 5			13Og01 TD	2012	$\alpha=100$ *
<sup>276</sup> Ds	142540# 550#					100# ms	0 <sup>+</sup>				$\alpha$ ?; SF ?
<sup>276</sup> Rg	147490# 630#					10# ms					$\alpha$ ?; SF ?
<sup>276</sup> Cn	150350# 600#					100# $\mu$ s	0 <sup>+</sup>				$\alpha$ ?; SF ?
* <sup>276</sup> Mt	T : average 13Ru11=750(+250-150) 13Og01=540(+140-90)										
* <sup>276</sup> Mt	T : 13Og01 supersedes 12Og02=680(+200-120)ms 04Og03=720(+870-250)ms										
* <sup>276</sup> Mt <sup>m</sup>	T : symmetrized from 6(+8-2) supersedes 12Og02										
<sup>277</sup> Hs	141490# 540#					11 ms 9	3/2 <sup>+</sup> #	14	10Du06 TD	2010	SF=100 *
<sup>277</sup> Hs <sup>m</sup>	141590# 550#	100#	100#			110 s 70		14	12Ho12 TD	2012	SF=100 *
<sup>277</sup> Hs <sup>p</sup>	142150# 580#	660#	200#								
<sup>277</sup> Mt	142970# 700#					9 s 6		14	13Og04 TD	2013	SF=100; $\alpha$ ? *
<sup>277</sup> Ds	145140# 380#					6 ms 3	11/2 <sup>+</sup> #	15	15U02 T	2010	$\alpha \approx 100$ ; SF ? *
<sup>277</sup> Rg	148340# 520#					10# ms					$\alpha$ ?; SF ?
<sup>277</sup> Cn	152400# 140#					850 $\mu$ s 280	3/2 <sup>+</sup> #	05	13Su04 T	1996	$\alpha=100$ *
* <sup>277</sup> Hs	T : symmetrized from 3.0(+14.4-1.4); 99Og10 one SF event at 16.5m, not trusted										
* <sup>277</sup> Hs <sup>m</sup>	T : (SF 1 event) symmetrized from $\tau=34(+164-16)$ s										
* <sup>277</sup> Mt	T : symmetrized from 13Og04=5(+9-2) s										
* <sup>277</sup> Ds	T : symmetrized from 15U02=4.1(+3.7-1.3)										
* <sup>277</sup> Cn	T : average 13Su04=610(+460-180) and 4 events : 08Mo09 at 1100 and 1220 $\mu$ s,										
* <sup>277</sup> Cn	T : 02Ho11 at 1406 $\mu$ s and 96Ho13 at 280 $\mu$ s										
<sup>278</sup> Mt	145740# 620#					7 s 3		10	14Kh04 T	2010	$\alpha=100$ *
<sup>278</sup> Mt <sup>p</sup>	146210# 650#	470#	200#								
<sup>278</sup> Ds	146380# 630#					270# ms	0 <sup>+</sup>				$\alpha$ ?; SF ?
<sup>278</sup> Rg	150520# 360#					8 ms 5		07	07Og02 TD	2007	$\alpha=100$ *
<sup>278</sup> Cn	152930# 440#					2# ms	0 <sup>+</sup>				$\alpha$ ?; SF ?
<sup>278</sup> Ed	158890# 180#					2.3 ms 1.3		05	12Mo25 TD	2004	$\alpha \approx 100$ *
* <sup>278</sup> Mt	T : average 14Kh04=3.6(+6.5-1.4) 13Og04=5.2(+6.2-1.8)s										
* <sup>278</sup> Rg	T : symmetrized from 4.2(+7.5-1.7)										
* <sup>278</sup> Ed	T : 3 events at 0.344, 4.930 and 0.667 ms; supersedes 08Mo09										
<sup>279</sup> Mt	147500# 670#					30# s					$\alpha$ ?; SF ?
<sup>279</sup> Ds	149130# 600#					210 ms 50		05	06Og05 TD	2004	SF=90; $\alpha=10$ *
<sup>279</sup> Ds <sup>p</sup>	149410# 610#	280#	100#								
<sup>279</sup> Rg	151780# 420#					180 ms 110		05	13Ru11 T	2004	$\alpha=100$ *
<sup>279</sup> Rg <sup>p</sup>	151910# 430#	130#	100#								
<sup>279</sup> Cn	155030# 460#					5# ms					$\alpha$ ?; SF ?
<sup>279</sup> Ed	159240# 700#					1# ms					$\alpha$ ?; SF ?
* <sup>279</sup> Ds	T : symmetrized from 200(+50-40); supersedes 04Og12=180(+50-30) and										
* <sup>279</sup> Ds	T : 04Og07=290(+350-100);										
* <sup>279</sup> Ds	T : others : 09St21 one SF event at 185 ms, 07Ei02 one SF event at 536 ms										
* <sup>279</sup> Rg	T : 13Ru11 one event at 16.1 ms and 04Og03 one event at 170										
<sup>280</sup> Ds	150520# 780#					11 s 6	0 <sup>+</sup>	05	01Og01 TD	1999	SF=100 *
<sup>280</sup> Rg	153890# 530#					4.3 s 0.7		05	13Ru11 T	2004	$\alpha=100$ *
<sup>280</sup> Cn	155700# 580#					5# ms	0 <sup>+</sup>				$\alpha$ ?; SF ?
<sup>280</sup> Ed	161140# 400#					10# ms					$\alpha$ ?; SF ?
* <sup>280</sup> Ds	T : 3 events at 6.93, 14.3 and 7.4 yield 6.6(+9-2.4), see 84Sc13										
* <sup>280</sup> Rg	T : average 13Ru11=6.4(+2.1-1.3) 13Og01=3.61(+0.90-0.60);										
* <sup>280</sup> Rg	T : 13Og01 supersedes 12Og02=3.53(+0.99-0.63)ms 04Og03=3.6(+4.3-1.3)ms										
<sup>281</sup> Ds	153430# 580#					14 s 4	3/2 <sup>+</sup> #	05	10Du06 TD	2004	SF=85 12; $\alpha=15$ 12 *
<sup>281</sup> Ds <sup>m</sup>	153470# 550#	40#	240#			0.9 s 0.7			12Ho12 TD	2012	$\alpha=100$ *
<sup>281</sup> Rg	155300# 810#					24 s 8		10	16Fo16 T	2010	SF=100 *
<sup>281</sup> Cn	158020# 390#					180 ms 80	3/2 <sup>+</sup> #	15	15U02 T	2010	$\alpha \approx 100$ ; SF ? *
<sup>281</sup> Ed	161810# 300#					100# ms					$\alpha$ ?; SF ?
* <sup>281</sup> Ds	T : average 10Du06=20(+20-7) 07Og01=11.1(+5.0-2.7); supersedes										
* <sup>281</sup> Ds	T : 04Og07=9.6(+5.0-2.5); 99Og10 one $\alpha$ event at 1.6 m, not trusted										
* <sup>281</sup> Ds	D : symmetrized from SF=91(+7-16)%; $\alpha=9(+16-7)$ %										
* <sup>281</sup> Ds <sup>m</sup>	T : symmetrized from 0.25(+1.18-0.11) s										
* <sup>281</sup> Rg	T : symmetrized from 16Fo16=21(+10-5), reanalyzed data of 13Og04=17(+6-3),										
* <sup>281</sup> Rg	T : 12Og06=26(+25-8), 10Og01=26(+25-8)										
* <sup>281</sup> Cn	T : symmetrized from 15U02=130(+120-40)										



**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>282</sup> Rg	157800# 650#		1.6 m 0.7			05 14Kh04	TD 2010	$\alpha=100$
<sup>282</sup> Cn	158980# 660#		900 $\mu$ s 240	0 <sup>+</sup>	05	06Og05	TD 2004	SF=100
<sup>282</sup> Ed	163730# 360#		140 ms 90		07	07Og02	TD 2007	$\alpha=100$
* <sup>282</sup> Rg	T : average 14Kh04=3.1(+5.7-1.2)m 13Og04=59(+55-19)s							
* <sup>282</sup> Cn	T : symmetrized from SF=820(+300-180); supersedes 04Og12=500(+330-140)							
* <sup>282</sup> Cn	T : also 10E106 one SF event at 522 $\mu$ s; 09St21 one SF at 3600 $\mu$ s							
* <sup>282</sup> Ed	T : symmetrized from 73(+134-29)							
<sup>283</sup> Rg	159280# 700#		30# s					$\alpha ?$ ; SF ?
<sup>283</sup> Cn	161490# 610#		4.1 s 1.0		06	06Og05	TD 2004	$\alpha=?$ ; SF<10
<sup>283</sup> Ed	164710# 440#		160 ms 100		05	13Ru11	T 2004	$\alpha=100$
* <sup>283</sup> Cn	T : symmetrized from 3.8(+1.2-0.7); supersedes 04Og12=4.0(+1.3-0.7) and							
* <sup>283</sup> Cn	T : 04Og07=6.1(+7.2-2.2); other 07Ho18=6.9(+6.9-2.3), SF=50							
* <sup>283</sup> Cn	T : 09St21 one $\alpha$ event at 1.92 s							
* <sup>283</sup> Cn	T : Four SF events at 99Og07=9.3 m, 3.8 m, 99Og05=3.0 m, 0.9 m, not trusted							
* <sup>283</sup> Ed	T : 13Ru11 one event at 68.4 ms and 04Og03 one event at 100							
<sup>284</sup> Cn	162550# 810#		104 ms 20	0 <sup>+</sup>	05	10Du06	TD 2004	SF=100
<sup>284</sup> Ed	166590# 530#		930 ms 140		05	13Ru11	T 2004	$\alpha=100$
<sup>284</sup> Fl	168920# 660#		3.3 ms 1.4	0 <sup>+</sup>	15	15Ut02	TD 2015	SF $\approx$ 100; $\alpha ?$
* <sup>284</sup> Cn	T : average 10Du06=101(+50-25) 07Og01=97(+31-19); supersedes							
* <sup>284</sup> Cn	T : 04Og12=101(+41-22) and 04Og07=98(+41-23)							
* <sup>284</sup> Cn	TD : 01Og01 3 $\alpha$ 's at 53.9 s, 10.3 s, 18.0 s, not trusted							
* <sup>284</sup> Ed	T : average 13Ru11=810(+230-150) 13Og01=970(+250-170);							
* <sup>284</sup> Ed	T : 13Og01 supersedes 12Og02=940(+290-180)ms 04Og03=480(+580-170)ms							
* <sup>284</sup> Fl	T : 5 events at 0.555, 8.588, 0.857, 7.246 and 0.529 ms							
<sup>285</sup> Cn	165170# 580#		32 s 9	5/2 <sup>+</sup> #	05	10Du06	TD 2004	$\alpha=100$
<sup>285</sup> Cn <sup>m</sup>	165740# 560# 570# 250#		15 s 12			12Ho12	TD 2012	$\alpha=100$
<sup>285</sup> Ed	167730# 810#		3.3 s 1.1		10	16Fo16	T 2010	$\alpha=100$
<sup>285</sup> Fl	171000# 390#		210 ms 100		15	15Ut02	T 2010	$\alpha\approx$ 100; SF ?
* <sup>285</sup> Cn	T : average 10Du06=30(+30-10) 07Og01=29(+13-7); supersedes							
* <sup>285</sup> Cn	T : 04Og07=34(+17-9); 99Og10 one event at 15.4 m, not trusted							
* <sup>285</sup> Cn <sup>m</sup>	T : symmetrized from 4.0(+19.1-1.8) s							
* <sup>285</sup> Ed	T : symmetrized from 16Fo16=2.9(+1.4-0.7), reanalyzed data of							
* <sup>285</sup> Ed	T : 13Og04=4.2(+1.4-0.8), 12Og06=4.9(+6.7-1.8), 10Og01=5.5(+5.0-1.8)							
* <sup>285</sup> Fl	T : symmetrized from 15Ut02=150(+140-50)							
<sup>286</sup> Ed	170010# 660#		7 s 3		10	14Kh04	T 2010	$\alpha=100$
<sup>286</sup> Fl	171770# 660#		140 ms 30	0 <sup>+</sup>	05	06Og05	TD 2004	SF $\approx$ 60; $\alpha\approx$ 40
* <sup>286</sup> Ed	T : average 14Kh04=2.9(+5.3-1.1) 13Og04=13(+12-4)							
* <sup>286</sup> Fl	T : symmetrized from 130(+40-20); supersedes 04Og12=160(+70-30) and							
* <sup>286</sup> Fl	T : 04Og07=290(+540-110); also one $\alpha$ each 10E106=76 ms, 09St21=301 ms							
<sup>287</sup> Ed	171250# 730#		2# m					$\alpha ?$ ; SF ?
<sup>287</sup> Fl	174070# 610#		520 ms 130		05	06Og05	TD 2004	$\alpha=100$
<sup>287</sup> Ef	177900# 440#		95 ms 60		05	13Ru11	T 2004	$\alpha=100$
* <sup>287</sup> Fl	T : symmetrized from 480(+160-90); supersedes 04Og12=510(+180-100)							
* <sup>287</sup> Fl	T : supersedes 04Og07=1.1(+1.3-0.4); 99Og07 2 evts 1.32, 14.4 s not trusted							
* <sup>287</sup> Fl	T : also 09St21 one $\alpha$ event at 815 ms							
* <sup>287</sup> Ef	T : 13Ru11 one event at 67.6 ms and 04Og03 one event at 32 ms							
<sup>288</sup> Fl	175040# 810#		750 ms 140	0 <sup>+</sup>	05	11Ga19	TD 2004	$\alpha=100$
<sup>288</sup> Ef	179770# 540#		170 ms 25		05	13Ru11	T 2004	$\alpha=100$
* <sup>288</sup> Fl	T : average 11Ga19=520(+220-130) 07Og01=800(+270-160); supersedes							
* <sup>288</sup> Fl	T : 10Du06=470(+240-120); 04Og12=800(+320-180) and 04Og07=630(+270-140)							
* <sup>288</sup> Fl	T : 01Og01=1800(+2100-600) re-assigned to <sup>289</sup> Fl							
* <sup>288</sup> Ef	T : average 13Ru11=150(+43-28) 13Og01=171(+42-28);							
* <sup>288</sup> Ef	T : 13Og01 supersedes 12Og02=173(+52-32) ms and 04Og03=87(+105-30) ms							

**Table I. The NUBASE2016 table (continued, Explanation of Table on page 18)**

Nuclide	Mass excess (keV)	Excitation energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
<sup>289</sup> F1	177560# 580#		2.4 s	0.6	5/2 <sup>+</sup> #	05 10Du06	TD 2004	$\alpha=100$ *
<sup>289</sup> F1 <sup>m</sup>	178330# 560# 770# 260#		1.1 s	0.8		12Ho12	TD 2012	$\alpha=100$ *
<sup>289</sup> Ef	180670# 810#		310 ms	90		10 16Fo16	T 2010	$\alpha=100$ *
<sup>289</sup> Lv	184530# 490#	RN	2# ms		5/2 <sup>+</sup> #	00 02Ni10	I	$\alpha ?$ *
* <sup>289</sup> F1	T : average 10Du06=0.97(+0.97-0.32) 07Og01=2.6(+1.2-0.7);							
* <sup>289</sup> F1	T : supersedes 04Og07=2.7(+1.4-0.7);							
* <sup>289</sup> F1	T : 99Og10 one event at 30.4 s, not trusted							
* <sup>289</sup> F1 <sup>m</sup>	T : symmetrized from 0.28(+1.35-0.13) s							
* <sup>289</sup> Ef	T : symmetrized from 16Fo16=270(+120-60), reanalyzed data of							
* <sup>289</sup> Ef	T : 13Og04=330(+120-80), 12Og06=430(+590-160), 10Og01=220(+260-80)							
* <sup>289</sup> Lv	T : 99Ni03=600(+860-300) $\alpha$ decay retracted by authors in 02Ni10							
<sup>290</sup> Ef	182890# 660#		410 ms	190		10 13Og04	T 2010	$\alpha=100$ *
<sup>290</sup> Lv	185200# 660#		8 ms	3	0 <sup>+</sup>	05 06Og05	TD 2004	$\alpha=100$ *
* <sup>290</sup> Ef	T : symmetrized from 13Og04=240(+280-90); other 14Kh04=1300(+2300-500)							
* <sup>290</sup> Lv	T : symmetrized from 7.1(+3.2-1.7); supersedes 04Og07=15(+26-6)							
<sup>291</sup> Ef	183990# 780#		1# s					$\alpha ?$ ; SF ? *
<sup>291</sup> Lv	187390# 610#		28 ms	15		05 06Og05	TD 2004	$\alpha=100$ *
<sup>291</sup> Eh	191800# 590#		2# ms					$\alpha ?$ ; SF ? *
* <sup>291</sup> Lv	T : symmetrized from 18(+22-6); supersedes 04Og07=6.3(+11.6-2.5)							
<sup>292</sup> Lv	188240# 810#		24 ms	12	0 <sup>+</sup>	05 04Og12	TD 2004	$\alpha=100$ *
<sup>292</sup> Eh	193580# 670#		10# ms					$\alpha ?$ ; SF ? *
* <sup>292</sup> Lv	T : symmetrized from 18(+16-6)							
* <sup>292</sup> Lv	T : 01Og01 reported one event at 46.9 ms, re-assigned to next isotope							
<sup>293</sup> Lv	190670# 590#		80 ms	40		05 07Og01	TD 2004	$\alpha=100$ *
<sup>293</sup> Lv <sup>m</sup>	191410# 560# 740# 270#		80 ms	60		12Ho12	TD 2012	$\alpha=100$ *
<sup>293</sup> Eh	194390# 810#		21 ms	6		10 16Fo16	T 2010	$\alpha=100$ *
<sup>293</sup> Ei	198870# 700#	RN	1# ms		1/2 <sup>+</sup> #	00 02Ni10	I	$\alpha ?$ *
* <sup>293</sup> Lv	T : symmetrized from 61(+57-20); supersedes 04Og07=53(+62-19)							
* <sup>293</sup> Lv <sup>m</sup>	T : symmetrized from 20(+96-9) ms							
* <sup>293</sup> Eh	T : symmetrized from 16Fo16=18(+8-4), reanalyzed data of							
* <sup>293</sup> Eh	T : 13Og04=22(+8-4), 12Og06=27(+12-6), 10Og01=14(+11-4)							
* <sup>293</sup> Ei	T : 99Ni03=120(+180-60) $\alpha$ decay retracted by authors in 02Ni10							
<sup>294</sup> Eh	196520# 660#		70 ms	30		10 14Kh04	T 2010	$\alpha=100$ *
<sup>294</sup> Ei	199460# 660#		1.15 ms	0.47	0 <sup>+</sup>	05 12Og06	T 2006	$\alpha=100$ *
* <sup>294</sup> Eh	T : average 14Kh04=51(+94-20) 13Og04=50(+60-18)							
* <sup>294</sup> Ei	T : 12Og06=0.135 ms (1 event) 06Og05=0.89 (4 events) 04Og12=1.8 ms (1 event)							
<sup>295</sup> Ei	201510# 640#		10# ms			04Og05	TD	$\alpha ?$ *
* <sup>295</sup> Ei	T : 04Og05 reports one $\alpha$ event at 2.55 ms ; re-assigned to <sup>294</sup> Ei							