

WEEKLY REPORT – WASTEWATER-BASED EPIDEMIOLOGICAL SURVEILLANCE OF THE SARS-COV-2

— RESULTS OF 08/11/2021

Authors: R. Janssens¹, H. Maloux¹, S. Hanoteaux¹, B. Verhaegen², K. Van Hoorde², K. Dierick², S. Quoilin¹, K. Blot¹, M. Lesenfans^{1*}

¹ Sciensano, Service Epidemiology of infectious diseases

² Sciensano, Service Foodborne pathogens

*Project responsible and contact: marie.lesenfans@sciensano.be

In collaboration with:



TABLE OF CONTENTS

TABLE OF CONTENTS	2
1. Summary	3
2. Introduction	4
3. Methodology	4
3.1. SAMPLE COLLECTION AND ANALYSIS	4
3.2. WASTEWATER RESULTS	5
3.3. ALERTING INDICATORS	5
3.4. CASES RESULTS.....	6
3.5. CAUTION POINTS FOR THE RESULTS INTERPRETATION.....	6
4. Results	6
4.1. REGIONS.....	6
4.2. PROVINCES	9
4.3. CATCHMENT AREAS.....	10
5. Appendix – Areas classified by alerting indicator	13

1. Summary

The circulation of the SARS-CoV-2 virus in the environment is assessed in the present work based on three indicators. The analysis of wastewater samples collected in 42 wastewater treatment plants covers 45% of the Belgian population, with increased accuracy in the urban areas. The results of the wastewater surveillance is a source of complementary information to the infection cases number as the populations represented are different. Indeed, the wastewater results do notably include all asymptomatic persons, and are independent of the testing strategy.

Here are the conclusions based on the latest results of November 08th 2021:

- At the national level: The numbers of areas for which the indicators "high circulation" and "fast increase" are fulfilled has reached their highest values since the beginning of the measurements (see Figure 1). Concretely, this means that more than half of the areas covered by this surveillance had reached at least 50% of the viral circulation level observed during the 3rd wave (= high circulation indicator);
- At the provincial level: all provinces are positive to at least one of the three alerting indicators. The provinces of Hainaut, Limburg, Namur, Oost-Vlaanderen, are of particular concern as 3 indicators are fulfilled in these provinces.
- At the covered areas level: Amongst the 40 areas covered¹, the number of treatment plants with at least one alerting indicator positive is 32, 22 belonging to the "High circulation" indicator, 26 belonging to the "Fast increase" indicator, and 14 belonging to the "Increasing trend" indicator. Last week (results of November 03th 2021²), 20 areas had at least one alerting indicator positive.
- Attention point for the covered areas of Froyennes, Genk, Hasselt, Houthalen-Centrum, Vallée du Hain (L'Orchis) and Wasmuel, where all three indicators are fulfilled.

The wastewater situation can be followed on a weekly basis on:

- The graphics available on the public [COVID-19 dashboard](#)
- The Risk Assessment Group (RAG) updating the weekly epidemiological situation through a report validated by the Risk Management Group (RMG) and published every Friday in [French](#) and [Dutch](#).
- Further details on the methodology applied for the wastewater surveillance can be found in [the Appendix Methodology document \(access online\)](#).

¹ Due to the recent flooding events, the treatment plants from Wegnez (Verviers) and Grosses-Battes (Liège) are temporarily out of order. These two areas had therefore not been screened for SARS-CoV-2. Fortunately, Liège is still well represented within this surveillance, however Verviers is not. Since the 13th of October, the covered area of Soumagne is added to the surveillance. No samples were taken in the area of Tessenderlo.

² No samples were taken on the 10th of November. Hence, the data presented here (from the 8th of November) are compared to the last available previous data (from the 3rd November 2021).

2. Introduction

In mid-September 2020 started the SARS-CoV-2 national wastewater surveillance project. The present report is an outcome of this project aiming to assess weekly the wastewater-based epidemiological situation of Belgium.

The surveillance is based on the analysis of water samples collected twice per week from the influent of 42 WasteWater Treatment Plants (WWTPs) spread over Belgium. The evolution of the SARS-CoV-2 viral concentrations measured over time in the different treatment plants is analyzed at different levels: regional, provincial, and the catchment area covered by the individual treatment plants. Also, alerting indicators were developed to highlight areas of concern regarding the high circulation, the fast evolution, and the increasing trend of the observed viral concentrations.

In this report, the weekly wastewater-based epidemiological situation is presented and discussed at the above-mentioned levels based on the three categories of alerting indicators. Moreover, the remaining sources of uncertainties are discussed together with their expected impacts on the wastewater results interpretation.

3. Methodology

3.1. SAMPLE COLLECTION AND ANALYSIS

The surveillance project, which started in mid-September 2020 covers around 45% of the Belgian population. The population covered is 40% in the Flemish region, 35% in the Walloon region, and nearly 100% in the Brussels region. Figure 1 shows the catchment areas covered by the 42 WWTPs included in the project, which are located in high population density areas. A catchment area is defined by the area delimiting the population covered by a specific wastewater sample, collected at the inlet of the WWTPs. Further coverage details can be found in Table 1 by province (see also Tables A4, A5 and A6) and on the [Sciensano public dashboard](#).

In practice, 24-hour composite samples are collected twice a week on Mondays and Wednesdays from the influent of WWTP and are analysed for the presence of SARS-CoV-2 RNA by three different laboratories. The resulting SARS-CoV-2 concentrations (3 targeted genes) are delivered to Sciensano within 2 days for data analysis purpose. Concretely, the results which are representative of Mondays and Wednesdays are respectively available on Wednesdays and Fridays.




Figure 1: The population located in the areas covered by the wastewater treatment plants (highlighted in yellow) and the population density for each municipality (indicated by the blue scale). Note that due to the recent flooding, the treatment plant of Wegnez is out of order. This area is, hence, not considered anymore in the wastewater surveillance and has been replaced by the treatment plant of Soumagne.

3.2. WASTEWATER RESULTS

The quantification of SARS-CoV-2 in RNA copies/ ml (concentration) was generalized in mid-February 2021 to all the treatment plants investigated. Before this date, the quantitative values were estimated based on the retrospective application of the quantification method (see details in Appendix) from mid-September 2020 to mid-February 2021 for Flanders and Brussels. In Wallonia, the quantitative results were available since the start.

Preliminary results allowed for estimating the limit of quantification of the analytical method at 20 copies/ml.

3.3. ALERTING INDICATORS

To highlight the areas of possible concerns, the three following types of alerting indicators are assessed twice a week, based on the viral concentration (RNA copies/ml) measured for the three targeted gene fragments (E, N1, and N2):

1. The **High circulation** indicator highlights the catchment areas where the viral concentrations are high. It corresponds to a situation where the viral concentrations exceed half of the highest value recorded since mid-February 2021.
2. The **Fast increase** indicator highlights the catchment areas where the viral concentrations have rapidly increased for the last week. It corresponds to a situation where the moving average on the past 7 days of the viral concentration has increased faster than 10% per day if being above the estimated limit of quantification. The increasing slope is normalized for each treatment plant.

3. The **Increasing trend** indicator highlights the catchment areas where the viral concentrations have been increasing for more than 6 days. It corresponds to a situation where the moving average on the past two weeks of the viral concentration has increased during more than 6 days.

3.4. CASES RESULTS

The cases number data presented in this report come from the COVID-19 laboratory-confirmed cases database centralized by Sciensano. The cases number, used to compute the 14 days-incidence, only accounts for the physical areas covered by the wastewater project (see Figure 1).

3.5. CAUTION POINTS FOR THE RESULTS INTERPRETATION

Only the trends, observed through the alerting indicators, should be assessed for the comparison of different areas. Absolute values should not be compared as the concentration values differ from an WWTP to another, notably due to the differences in population sizes covered. Additionally, the situation comparison between the regions should be considered with caution. The degree of comparability is not yet known and depends on the comparability of the results between the different laboratories performing the analysis (see [the Appendix Methodology document \(access online\)](#)).

Moreover, the wastewater concentrations and the cases numbers presented in this report do not originate from the same population, even though the positive cases are selected only for the municipalities covered by the wastewater surveillance. For instance, the wastewater results account for all infected persons (whatever age or symptomatic condition) while the cases include only the persons tested clinically positive. Likewise, an infected person covered by the wastewater results could be associated with another area in the clinical surveillance as the person's postal code is used for clinical statistics (mobility bias). Therefore, the correlation between the wastewater concentrations and the cases number varies according to the area considered. The wastewater results are thus complementary and independent information to the results of the cases.

The correlation between the wastewater viral concentrations and the cases numbers could also be influenced by the vaccination campaign and the circulation of variants.

Finally, all values below the limit of quantification (< 20 RNA copies/ml) should be considered as non-quantitatively reliable values. A reported value lower than 20 copies/ml only shows that SARS-CoV-2 has been detected in the sample at an undetermined concentration.

4. Results


4.1. REGIONS

Figure 2 presents the quantitative SARS-CoV-2 RNA concentration in the wastewaters and the 14 days incidence for each region, compiling the incidence data of the area covered by the wastewater surveillance. The estimated viral concentrations can also be seen for the period before mid-February (see 2.2. in [the Appendix Methodology document \(access online\)](#) for more details).

The second wave peak occurring in November 2020 can be seen in the three regions in Figure 2 below. This remains true for the third wave, but to a lower extent in Wallonia. Several hypotheses could explain this: (1) the sizes of the treatment plants in Wallonia are smaller than the ones located in the two other regions, affecting the viral concentrations measured; (2) the extent to which the results are comparable between the different laboratories is unknown (see [Methodology](#) for more details)

More recently, the viral concentration in the wastewater was increasing since the beginning of July in Wallonia, Brussels and Flanders, with a newly strong increasing period observed at the beginning of October, illustrating the fourth wave (Figure 2).

The results indicate that the viral concentrations in the three regions are quickly increasing since beginning of October. It should be noted that the last viral concentrations in wastewaters are not displayed in the figure as the regional corresponding trends are still to be validated with the next week results³.



³ The trends of SARS-CoV-2 circulation in wastewater, given by the dark line on the graphs, corresponds to a 14 days centred mean of the concentrations measured.

Figure 2: The SARS-CoV-2 RNA estimated and quantified concentrations expressed as copies/ml (two weeks centered moving average applied on the linear interpolation), the estimated limit of quantification of 20 SARS-CoV-2 RNA copies/ ml, and the 14 days incidence in the population covered by the wastewater surveillance since the surveillance starting date (graph set above) and mid-February 2021 (graph set below).

4.2. PROVINCES

Table 1 shows, for each Province, the results associated with the samples of Monday November 08th 2021, for the three alerting indicators:

- Last week on Monday November 03th 2021, 4 provinces had at least one alerting indicator positive: Antwerpen, Brabant Wallon, Hainaut, and Limburg.
- This week on Monday November 08th 2021, all provinces had at least one alerting indicator positive. The provinces of Hainaut, Limburg, Namur, Oost-Vlaanderen, are of particular concern as 3 indicators are fulfilled in these provinces.
- It should be noted that although the 50% threshold of viral circulation is not exceeded in the province of Brabant Wallon (in comparison of the highest level observed since the 3rd wave), the viral concentration is still high (45%).

Table 1: Alerting indicators fulfilled (1) or not (0) on November 08th 2021 and the percentage of population covered by the provinces. Columns represent the population coverage of the WWTP within the Province (Pop. coverage), the High Circulation indicator (High), the Fast increase indicator (Fast) and the Increasing trend indicator (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table. Missing data is indicated with a “/”.

Province	Pop. coverage	High	Fast	Incr.	Norm. viral cc. (%) ¹	Mean viral. cc. (c./ml) ²	Norm. evol. (%/day) ³	Incr. days ⁴
Antwerpen	39%	0	0	1	32	250	10	21
Brabant Wallon	33%	0	0	1	45	122	6	14
Brussels	100%	1	1	0	92	8679	28	5
Hainaut	29%	1	1	1	59	110	19	7
Liège	43%	1	1	0	70	288	15	4
Limburg	26%	1	1	1	100	432	30	19
Luxembourg	10%	1	1	0	55	107	13	3
Namur	23%	1	1	1	78	249	15	8
Oost-Vlaanderen	38%	1	1	1	66	1249	50	9
Vlaams-Brabant	49%	1	1	0	77	2544	32	5
West-Vlaanderen	52%	0	1	0	38	1299	12	5

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area since mid-February 2021.

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/day) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration

4.3. CATCHMENT AREAS

Table 2 shows, for each catchment area, the values of the three alerting indicators obtained based on the results of last Wednesday's sample. The number of treatment plants with at least one alerting indicator positive is 32 (out of 40 areas covered). Last week (results of November 03th 2021), 20 areas had at least one alerting indicator positive.

- The indicator "High circulation" is fulfilled in 22 covered areas. The full list of these areas can be found in Table 2 and in Appendix A1. In several areas, the measured viral concentration exceeded the maximal concentration registered during the 3rd wave: Amay, Basse Wavre (Dyle), Brussels-North, Froyennes, Genk, Grimbergen, Harelbeke, Hasselt, Houthalen-Centrum, Leuven, Liège Oupeye, Liège Sclessin, Montignies-sur-Sambre, Namur-Brumagne, Soumagne, Vallée du Hain (L'Orchis) and Wasmuel.
- The indicator "Fast increase" is fulfilled in 26 covered areas. Amongst these covered areas, an increase of more than 50% per day of the normalized viral concentration is observed for 3 areas: Destelbergen (106% increase per day), Leuven (66% increase per day), and Gent (61% increase per day). Further details can be seen in Appendix A2.
- The indicator "Increasing trend" is fulfilled in 14 covered areas. Amongst these areas, the viral concentration is continually increasing since two weeks in 11 covered areas. The full list of these areas can be found in Table 2 and in Appendix A3.
- Attention point for the covered areas of Froyennes, Genk, Hasselt, Houthalen-Centrum, Vallée du Hain (L'Orchis), and Wasmuel, where all three indicators are fulfilled.

The wastewater results can be accessed online for each area on the [COVID-19 dashboard](#).

Table 2 is a snapshot of the number of areas highlighted by the indicators for the last results obtained, which correspond to the 24h-representative samples of last Monday November 08th 2021.

Table 2: Alerting indicators fulfilled (1) or not (0) on November 08th 2021 for the catchment areas covered by the wastewater treatment plants. Columns represent the population coverage of the WWTP within the Province (Pop. coverage), the High Circulation indicator(High), the Fast increase indicator (Fast) and the Increasing trend indicator (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ²	Norm evol. (%/d) ³	Incr days ⁴
Oost-Vlaanderen	Aalst	1	1	0	76	1557	32	5
Antwerpen	Aartselaar	0	1	1	43	257	22	26
Antwerpen	Antwerpen-North	1	0	1	84	267	5	28
Antwerpen	Antwerpen-South	0	0	1	12	243	2	33
Luxembourg	Arlon	1	1	0	66	127	22	5
Vlaams-Brabant	Beersel	0	1	0	27	1737	24	5
West-Vlaanderen	Brugge	1	1	0	56	533	14	5
Brussels	Brussels-North	1	1	0	100	9189	26	5
Brussels	Brussel-South	1	1	0	66	6970	34	5
Oost-Vlaanderen	Dendermonde	0	1	1	30	108	19	14
Oost-Vlaanderen	Destelbergen	0	1	0	44	2343	106	5
Antwerpen	Deurne	0	0	1	4	257	5	12

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ²	Norm evol. (%/d) ³	Incr days ⁴
Hainaut	Froyennes	1	1	1	100	209	20	19
Limburg	Genk	1	1	1	100	466	45	12
Oost-Vlaanderen	Gent	1	1	0	88	1448	61	5
Vlaams-Brabant	Grimbergen	1	1	0	100	3104	13	5
Limburg	Hasselt	1	1	1	100	451	44	26
Limburg	Houthalen-Centrum	1	1	1	100	252	20	33
Vlaams-Brabant	Leuven	1	1	0	100	3199	66	5
Vlaams-Brabant	Liedekerke	1	1	0	60	1710	18	5
Liège	Liège Oupeye	1	1	0	73	219	25	5
Liège	Liège Sclessin	1	0	0	80	494	6	5
Antwerpen	Mechelen-Noord	1	1	0	68	248	32	5
Namur	Mornimont	0	0	1	24	38	3	14
Namur	Namur-Brumagne	1	1	0	100	337	20	5
West-Vlaanderen	Oostende	0	1	0	47	1256	16	5
West-Vlaanderen	Roeselare	0	1	0	13	1758	13	5
Oost-Vlaanderen	Sint-Niklaas	0	1	1	30	325	25	28
Liège	Soumagne	1	1	0	78	338	49	5
Antwerpen	Turnhout	1	0	1	55	215	7	26
Brabant Wallon	Vallée du Hain (L'Orchis)	1	1	1	100	266	36	33
Hainaut	Wasmuel	1	1	1	100	161	39	12

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area since mid-February 2021.

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/day) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

Figure 3 was developed to offer a dynamic view of the three indicators over time.

The numbers of areas for which the indicators "high circulation" and "fast increase" are fulfilled has reached their highest values since the beginning of the measurements (see Figure 1). Concretely, this means that more than half of the areas covered by this surveillance had reached at least 50% of the viral circulation level observed during the 3rd wave (= high circulation indicator);



Figure 3: The number of areas (amongst the 40 covered by the wastewater surveillance this week and the 42 normally considered), with positive alerting indicators (latest results on November 08th 2021).

5. Appendix – Areas classified by alerting indicator

A 1: Covered areas (22 out of 40 on November 08th 2021) characterized as “High circulation” sorted in the descending order of importance.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/d) ³	Incr days ⁴
Brussels	Brussels-North	1	1	0	100	9189	26	5
Hainaut	Froyennes	1	1	1	100	209	20	19
Limburg	Genk	1	1	1	100	466	45	12
Vlaams-Brabant	Grimbergen	1	1	0	100	3104	13	5
Limburg	Hasselt	1	1	1	100	451	44	26
Limburg	Houthalen-Centrum	1	1	1	100	252	20	33
Vlaams-Brabant	Leuven	1	1	0	100	3199	66	5
Namur	Namur-Brumagne	1	1	0	100	337	20	5
Brabant Wallon	Vallée du Hain (L'Orchis)	1	1	1	100	266	36	33
Hainaut	Wasmuel	1	1	1	100	161	39	12
Oost-Vlaanderen	Gent	1	1	0	88	1448	61	5
Antwerpen	Antwerpen-North	1	0	1	84	267	5	28
Liège	Liège Sclessin	1	0	0	80	494	6	5
Liège	Soumagne	1	1	0	78	338	49	5
Oost-Vlaanderen	Aalst	1	1	0	76	1557	32	5
Liège	Liège Oupeye	1	1	0	73	219	25	5
Antwerpen	Mechelen-Noord	1	1	0	68	248	32	5
Luxembourg	Arlon	1	1	0	66	127	22	5
Brussels	Brussel-South	1	1	0	66	6970	34	5
Vlaams-Brabant	Liedekerke	1	1	0	60	1710	18	5
West-Vlaanderen	Brugge	1	1	0	56	533	14	5
Antwerpen	Turnhout	1	0	1	55	215	7	26

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area since mid-February 2021.

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/day) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the third wave.

A 2: Covered areas (26 out of 40 on the November 08th 2021) characterized as “Fast increase” sorted in the descending order of importance.

Province	WWTP	High	Fast	Incr.	Norm viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/d) ³	Incr days ⁴
Oost-Vlaanderen	Destelbergen	0	1	0	44	2343	106	5
Vlaams-Brabant	Leuven	1	1	0	100	3199	66	5
Oost-Vlaanderen	Gent	1	1	0	88	1448	61	5
Liège	Soumagne	1	1	0	78	338	49	5
Limburg	Genk	1	1	1	100	466	45	12
Limburg	Hasselt	1	1	1	100	451	44	26
Hainaut	Wasmuel	1	1	1	100	161	39	12
Brabant Wallon	Vallée du Hain (L'Orchis)	1	1	1	100	266	36	33
Brussels	Brussel-South	1	1	0	66	6970	34	5
Oost-Vlaanderen	Aalst	1	1	0	76	1557	32	5
Antwerpen	Mechelen-Noord	1	1	0	68	248	32	5
Brussels	Brussels-North	1	1	0	100	9189	26	5
Liège	Liège Oupeye	1	1	0	73	219	25	5
Oost-Vlaanderen	Sint-Niklaas	0	1	1	30	325	25	28
Vlaams-Brabant	Beersel	0	1	0	27	1737	24	5
Antwerpen	Aartselaar	0	1	1	43	257	22	26
Luxembourg	Arlon	1	1	0	66	127	22	5
Hainaut	Froyennes	1	1	1	100	209	20	19
Limburg	Houthalen-Centrum	1	1	1	100	252	20	33
Namur	Namur-Brumagne	1	1	0	100	337	20	5
Oost-Vlaanderen	Dendermonde	0	1	1	30	108	19	14
Vlaams-Brabant	Liedekerke	1	1	0	60	1710	18	5
West-Vlaanderen	Oostende	0	1	0	47	1256	16	5
West-Vlaanderen	Brugge	1	1	0	56	533	14	5
Vlaams-Brabant	Grimbergen	1	1	0	100	3104	13	5
West-Vlaanderen	Roeselare	0	1	0	13	1758	13	5

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area since mid-February 2021.

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/day) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the third wave.

A 3: Covered areas (15 out of 40 on the November 08th 2021) characterized as “Increasing trend” sorted in the descending order of importance.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/d) ³	Incr days ⁴
Antwerpen	Antwerpen-South	0	0	1	12	243	2	33
Limburg	Houthalen-Centrum	1	1	1	100	252	20	33
Brabant Wallon	Vallée du Hain (L'Orchis)	1	1	1	100	266	36	33
Antwerpen	Antwerpen-North	1	0	1	84	267	5	28
Oost-Vlaanderen	Sint-Niklaas	0	1	1	30	325	25	28
Antwerpen	Aartselaar	0	1	1	43	257	22	26
Limburg	Hasselt	1	1	1	100	451	44	26
Antwerpen	Turnhout	1	0	1	55	215	7	26
Hainaut	Froyennes	1	1	1	100	209	20	19
Oost-Vlaanderen	Dendermonde	0	1	1	30	108	19	14
Namur	Mornimont	0	0	1	24	38	3	14
Antwerpen	Deurne	0	0	1	4	257	5	12
Limburg	Genk	1	1	1	100	466	45	12
Hainaut	Wasmuel	1	1	1	100	161	39	12

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area since mid-February 2021.

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/day) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the third wave.

A 4: Covered areas (8 out of 40 on the November 08th 2021) in which no alerting indicator are fulfilled.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/d) ³	Incr days ⁴
Liège	Amay	0	0	0	15	27	-11	0
Brabant Wallon	Basse Wavre (Dyle)	0	0	0	4	16	-16	0
West-Vlaanderen	Harelbeke	0	0	0	32	2114	8	5
Luxembourg	Marche-en-Famenne	0	0	0	31	61	-9	0
Hainaut	Marchienne-au-Pont	0	0	0	12	26	-0	0
West-Vlaanderen	Menen	0	0	0	7	1592	7	5

Province	WWTP	High	Fast	Incr.	Norm - viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/d) ³	Incr days ⁴
Hainaut	Montignies-sur-Sambre	0	0	0	23	65	3	0
Hainaut	Mouscron versant Espierres	0	0	0	14	26	6	0

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area since mid-February 2021.

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/day) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the third wave.