

WEEKLY REPORT – WASTEWATER-BASED EPIDEMIOLOGICAL SURVEILLANCE OF THE SARS-COV-2

— RESULTS OF 08/12/2021

Authors: R. Janssens¹, H. Maloux¹, S. Hanoteaux¹, B. Verhaegen², K. Van Hoorde², K. Dierick², S. Quoilin¹, K. Blot¹, M. Lesenfans^{1*}

¹ Sciensano, Service Epidemiology of infectious diseases

² Sciensano, Service Foodborne pathogens

*Project responsible and contact: marie.lesenfans@sciensano.be

In collaboration with:



TABLE OF CONTENTS

TABLE OF CONTENTS	2
1. Summary	3
2. Introduction	4
3. Methodology	4
3.1. SAMPLE COLLECTION AND ANALYSIS	4
3.2. WASTEWATER RESULTS	5
3.3. ALERTING INDICATORS	5
3.4. CASES RESULTS.....	6
3.5. CAUTION POINTS FOR THE RESULTS INTERPRETATION.....	6
4. Results	7
4.1. REGIONS.....	7
4.2. PROVINCES	9
4.3. CATCHMENT AREAS.....	11
4.4. VARIANTS	Error! Bookmark not defined.
5. Appendix – Areas classified by alerting indicator	14

1. Summary

The circulation of the SARS-CoV-2 virus in the environment is assessed in the present work based on three indicators. The analysis of wastewater samples collected in 42 wastewater treatment plants covers 45% of the Belgian population, with increased accuracy in the urban areas. The results of the wastewater surveillance is a source of complementary information to the infection cases number as the populations represented are different. Indeed, the wastewater results do notably include all asymptomatic persons, and are independent of the testing strategy.

Here are the conclusions based on the latest results of December 08th 2021:

- At the national level: The number of "High circulation" areas is high and stable since four weeks. The rainy event recorded on 01/12 has impacted the 3 indicators to different extends: the High circulation areas were underestimated last week, while the fast and increasing trend areas' number are overestimated this week. Overall, a high circulation of the virus is observed but with a decreasing tendency.
- At the provincial level: Brabant Wallon, Brussels, Hainaut, Liège, Limburg, Luxembourg, Namur, Oost-Vlaanderen, Vlaams-Brabant, and West-Vlaanderen are the provinces positive to at least one of the three alerting indicators. Also, the provinces of Brussels, Limburg, Oost-Vlaanderen, and Vlaams-Brabant are of particular concern as the conditions for 2 indicators are met in these provinces. Brussels' region shows this week an Increasing trend and a High circulation, which was not the case last week.
- At the covered areas level: among the 41 areas covered¹, the number of treatment plants with at least one alerting indicator positive is 31, 26 belonging to the High circulation indicator, 17 belonging to the "Fast increase" indicator, and 7 belonging to the "Increasing trend" indicator. Last week (results of December 01th 2021), 12 areas (out of 33 areas) had at least one alerting indicator positive.
- An alerting situation is evidenced for the covered areas of Amay, Brussels-North, Froyennes, Grimbergen, Leuven, and Mouscron versant Espierres as all three indicators are fulfilled.
- The appearance of the Omicron Variant of Concern is unlikely to have a substantial impact on the wastewater surveillance (see 4.4 Variants). Deeper investigation on the impact of this variant is currently performed.

The wastewater situation can be followed on a weekly basis on:

- The graphics available on the public [COVID-19 dashboard](#)
- The Risk Assessment Group (RAG) updating the weekly epidemiological situation through a report validated by the Risk Management Group (RMG) and published every Friday in [French](#) and [Dutch](#).
- Further details on the methodology applied for the wastewater surveillance can be found in [the Appendix Methodology document \(access online\)](#).

¹ Due to the recent flooding events, the treatment plants from Wegnez (Verviers) and Grosses-Battes (Liège) are temporarily out of order. These two areas had therefore not been screened for SARS-CoV-2. Liège is still well represented within this surveillance, however Verviers is not. Since the 13th of October, the covered area of Soumagne is added to the surveillance.

2. Introduction

In mid-September 2020 started the SARS-CoV-2 national wastewater surveillance project. The present report is an outcome of this project aiming to assess weekly the wastewater-based epidemiological situation of Belgium.

The surveillance is based on the analysis of water samples collected twice per week from the influent of 42 WasteWater Treatment Plants (WWTPs) spread over Belgium. The evolution of the SARS-CoV-2 viral concentrations measured over time in the different treatment plants is analyzed at different levels: regional, provincial, and the catchment area covered by the individual treatment plants. Also, alerting indicators were developed to highlight areas of concern regarding the high circulation, the fast evolution, and the increasing trend of the observed viral concentrations.

In this report, the weekly wastewater-based epidemiological situation is presented and discussed at the above-mentioned levels based on the three categories of alerting indicators. Moreover, the remaining sources of uncertainties are discussed together with their expected impacts on the wastewater results interpretation.

3. Methodology

3.1. SAMPLE COLLECTION AND ANALYSIS

The surveillance project, which started in mid-September 2020 covers around 45% of the Belgian population. The population covered is 40% in the Flemish region, 35% in the Walloon region, and nearly 100% in the Brussels region. Figure 1 shows the catchment areas covered by the 42 WWTPs included in the project, which are located in high population density areas. A catchment area is defined by the area delimiting the population covered by a specific wastewater sample, collected at the inlet of the WWTPs. Further coverage details can be found in Table 1 by province (see also Table A1 in the [Appendix Methodology document \(access online\)](#)) and on the [Sciensano public dashboard](#).

In practice, 24-hour composite samples are collected twice a week on Mondays and Wednesdays from the influent of WWTP and are analysed for the presence of SARS-CoV-2 RNA by three different laboratories. The resulting SARS-CoV-2 concentrations (3 targeted genes) are delivered to Sciensano within 2 days for data analysis purpose. Concretely, the results which are representative of Mondays and Wednesdays are respectively available on Wednesdays and Fridays.




Figure 1: The population located in the areas covered by the wastewater treatment plants (highlighted in yellow) and the population density for each municipality (indicated by the blue scale). Note that due to the recent flooding, the treatment plant of Wegnez is out of order. This area is, hence, not considered anymore in the wastewater surveillance and has been replaced by the treatment plant of Soumagne.

3.2. WASTEWATER RESULTS

The quantification of SARS-CoV-2 in RNA copies/ ml (concentration) was generalized in mid-February 2021 to all the treatment plants investigated. Before this date, the quantitative values were estimated based on the retrospective application of the quantification method (see details in Appendix) from mid-September 2020 to mid-February 2021 for Flanders and Brussels. In Wallonia, the quantitative results were available since the start.

Preliminary results allowed for estimating the limit of quantification of the analytical method at 20 copies/ml.

3.3. ALERTING INDICATORS

To highlight the areas of possible concerns, the three following types of alerting indicators are assessed twice a week, based on the viral concentration (RNA copies/ml) measured for the three targeted gene fragments (E, N1, and N2):

1. The **High circulation** indicator highlights the catchment areas where the viral concentrations are high. It corresponds to a situation where the viral concentrations exceed half of the highest value recorded during the third wave (i.e. from mid-February 2021 till begin of May).
2. The **Fast increase** indicator highlights the catchment areas where the viral concentrations have rapidly increased for the last week. It corresponds to a situation where the moving average on the past 7 days of the viral concentration has increased faster than 70% per week if being above the estimated limit of quantification. The increasing slope is normalized for each treatment plant.

3. The **Increasing trend** indicator highlights the catchment areas where the viral concentrations have been increasing for more than 6 days. The indicator is computed based on the moving average on the past two weeks of the viral concentration has increased during more than 6 days.

The indicators were developed in order to be able to track the different phases of an outbreak. Typically, when the viral concentrations in wastewaters start to increase in an area, the indicator Increasing trend will be the first fulfilled. If the concentrations increase quickly, the Fast increase indicator will then be fulfilled. Finally, after the initial increasing phase, the concentrations in an area will be sufficiently high to result in the High circulation to be fulfilled.

3.4. CASES RESULTS

The cases number data presented in this report come from the COVID-19 laboratory-confirmed cases database centralized by Sciensano. The cases number, used to compute the 14 days-incidence, only accounts for the physical areas covered by the wastewater project (see Figure 1).

3.5. CAUTION POINTS FOR THE RESULTS INTERPRETATION

Only the trends, observed through the alerting indicators, should be assessed for the comparison of different areas. Absolute values should not be compared as the concentration values differ from an WWTP to another, notably due to the differences in population sizes covered. Additionally, the situation comparison between the regions should be considered with caution. The degree of comparability is not yet known and depends on the comparability of the results between the different laboratories performing the analysis (see [the Appendix Methodology document \(access online\)](#)).

Moreover, the wastewater concentrations and the cases numbers presented in this report do not originate from the same population, even though the positive cases are selected only for the municipalities covered by the wastewater surveillance. For instance, the wastewater results account for all infected persons (whatever age or symptomatic condition) while the cases include only the persons tested clinically positive. Likewise, an infected person covered by the wastewater results could be associated with another area in the clinical surveillance as the person's postal code is used for clinical statistics (mobility bias). Therefore, the correlation between the wastewater concentrations and the cases number varies according to the area considered. The wastewater results are thus complementary and independent information to the results of the cases.

The correlation between the wastewater viral concentrations and the cases numbers could also be influenced by the vaccination campaign and the circulation of variants.

Finally, all values below the limit of quantification (< 20 RNA copies/ml) should be considered as non-quantitatively reliable values. A reported value lower than 20 copies/ml only shows that SARS-CoV-2 has been detected in the sample at an undetermined concentration.

4. Results

4.1. REGIONS

Figure 2 presents the quantitative SARS-CoV-2 RNA concentration in the wastewaters and the 14 days incidence for each region, compiling the incidence data of the area covered by the wastewater surveillance. The estimated viral concentrations can also be seen for the period before mid-February (see 2.2. in [the Appendix Methodology document \(access online\)](#) for more details).

The second wave peak occurring in November 2020 can be seen in the three regions in Figure 2 below. This remains true for the third wave, but to a lower extent in Wallonia. Several hypotheses could explain this: (1) the sizes of the treatment plants in Wallonia are smaller than the ones located in the two other regions, affecting the viral concentrations measured; (2) the extent to which the results are comparable between the different laboratories is unknown (see [Methodology](#) for more details)

Recently, the viral concentrations in the wastewater were increasing since the beginning of October, illustrating the fourth wave (Figure 2). The last results seem to indicate that the viral concentrations in Wallonia and Flanders are in a decreasing trend, while the concentrations may increase back in Brussels.

It should be noted that the last viral concentrations in wastewaters are not displayed in the figure as the regional corresponding trends are still to be validated with the next week results².

² The trends of SARS-CoV-2 circulation in wastewater, given by the dark line on the graphs, corresponds to a 14 days centred mean of the concentrations measured.




Figure 2: The SARS-CoV-2 RNA estimated and quantified concentrations expressed as copies/ml (two weeks centered moving average applied on the linear interpolation), the estimated limit of quantification of 20 SARS-CoV-2 RNA copies/ ml, and the 14 days incidence in the population covered by the wastewater surveillance since the surveillance starting date (graph set above) and mid-February 2021 (graph set below).

4.2. PROVINCES

Table 1 shows, for each Province, the results associated with the samples of Wednesday December 08th 2021, for the three alerting indicators:

- Last week on Wednesday December 01th 2021, 6 provinces had at least one alerting indicator positive: Brabant Wallon, Brussels, Hainaut, Liège, Luxembourg, and Namur.
- This week on Wednesday December 08th 2021, 10 provinces had at least one alerting indicator positive: Brabant Wallon, Brussels, Hainaut, Liège, Limburg, Luxembourg, Namur, Oost-Vlaanderen, Vlaams-Brabant, and West-Vlaanderen. The provinces of Brussels, Limburg, Oost-Vlaanderen, and Vlaams-Brabant are of particular concern as 2 indicators are fulfilled in these provinces.
- Brussels' region shows this week an "Increasing trend" and a "High circulation", which was not the case last week.
- It should be noted that although the 50% threshold of viral circulation is not exceeded (in comparison of the highest level observed since the 3rd wave), the viral concentrations in the province of West-Vlaanderen are still high (43%).

Table 1 allows to track the changes between the situation as of today (December 08th 2021) and the situation as of last week (December 01th 2021). Hereby, three distinct cases are taken into account:

1. If a province has at least one indicator fulfilled this week and it was not the case last week, its **name is displayed in bold in the table**;
2. If a province has at least one indicator fulfilled this week and if it also was the case last week, any change in indicator fulfilment (i.e. if the value for any indicator has changed from 0 to 1 or from 1 to 0) is indicated in **coloured bold text**;

Table 1: Alerting indicators fulfilled (1) or not (0) on December 08th 2021 and the percentage of population covered by the provinces. Columns represent the population coverage of the WWTP within the Province (Pop. coverage), the "High Circulation" indicator (High), the "Fast increase" indicator (Fast) and the "Increasing trend" indicator (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table. Missing data is indicated with a "/".

Province	Pop. coverage	High	Fast	Incr.	Norm. viral cc. (%) ¹	Mean viral. cc. (c./ml) ²	Norm. evol. (%/w) ³	Incr. days ⁴
Antwerpen	41%	0	0	0	28	275	8	0
Brabant Wallon	33%	1	0	0	211	274	-52	0
Brussels	100%	1	0	1	180	14705	60	7
Hainaut	29%	1	0	0	152	288	19	2
Liège	50%	1	0	0	242	530	10	2
Limburg	26%	1	1	0	66	162	75	0
Luxembourg	10%	1	0	0	114	221	-42	0
Namur	23%	1	0	0	55	126	17	0
Oost-Vlaanderen	38%	1	1	0	80	1513	197	0
Vlaams-Brabant	61%	1	1	0	102	3025	141	4

Province	Pop. coverage	High	Fast	Incr.	Norm. viral cc. (%) ¹	Mean viral. cc. (c./ml) ²	Norm. evol. (%/w) ³	Incr. days ⁴
West-Vlaanderen	52%	0	1	0	43	1003	370	0

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area during the third wave (i.e. from mid-February 2021 till begin of July).

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/week) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration

4.3. CATCHMENT AREAS

Table 2 shows, for each catchment area, the values of the three alerting indicators obtained based on the results of last Wednesday's sample. The number of treatment plants with at least one alerting indicator positive is 31 (out of 41 areas covered). Last week (results of December 01th 2021), 12 areas (out of 41 area covered) had at least one alerting indicator positive.

- The indicator "High circulation" is fulfilled in 26 covered areas. The full list of these stations can be found in Table 2 and in Appendix A1. In several areas, the measured viral concentration exceeded the maximal concentration registered during the 3rd wave. The full list of these stations can be found in the tables of the different Appendices.
- The indicator "Fast increase" is fulfilled in 17 covered areas. Among these covered areas, an increase of more than 300% per week of the normalized viral concentration (i.e. the measured concentrations have more than quadrupled during last week) is observed for 5 areas: Brugge (623% increase per week), Oostende (329% increase per week), Gent (323% increase per week), Destelbergen (320% increase per week) and Leuven (312% increase per week). Further details can be found in Appendix A2.
- The indicator "Increasing trend" is fulfilled in 7 covered areas. Among these areas, the viral concentration is continually increasing since two weeks in 1 covered areas: Amay (16 days). Further details can be found in Appendix A3.
- An alerting situation is evidenced for the covered areas of Amay, Brussels-North, Froyennes, Grimbergen, Leuven, and Mouscron versant Espierres as all three indicators are fulfilled.
- As mentioned last week, on the 01st of December, the numbers of areas in which the different indicators were fulfilled were impacted by the high precipitation and the lack of data for 8 covered areas.

The wastewater results can be accessed online for each area on the [COVID-19 dashboard](#).

Table 2 is a snapshot of the number of areas highlighted by the indicators for the last results obtained, which correspond to the 24h-representative samples of last Wednesday December 08th 2021.

Table 2 allows to track the changes between the situation as of today (December 08th 2021) and the situation as of last week (December 01th 2021). Hereby, three distinct cases are taken into account:

1. If an area has at least one indicator fulfilled this week and it was not the case last week, its **name is displayed in bold** in the table;
2. If an area has at least one indicator fulfilled this week and if it also was the case last week, any change in indicator fulfilment (i.e. if the value for any indicator has changed from 0 to 1 or from 1 to 0) is indicated in **coloured bold text**;
3. Any area which had at least one indicator fulfilled last week but not this week is listed below Table 1.

Table 2: Alerting indicators fulfilled (1) or not (0) on December 08th 2021 for the catchment areas covered by the wastewater treatment plants. Columns represent the population coverage of the WWTP within the Province (Pop. coverage), the "High circulation indicator" (High), the "Fast increase" indicator (Fast) and the "Increasing trend" indicator (Incr.). The specifications of the four last columns are explained in the footnotes 1-4 below the table.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ²	Norm. evol. (%/w) ³	Incr days ⁴
Oost-Vlaanderen	Aalst	1	1	0	74	1515	107	0

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ²	Norm evol. (%/w) ³	Incr days ⁴
Liège	Amay	1	1	1	168	246	113	16
Antwerpen	Antwerpen-North	1	0	0	69	217	-46	0
Luxembourg	Arlon	1	0	0	121	233	-38	0
Brabant Wallon	Basse Wavre (Dyle)	1	0	0	217	299	-61	0
West-Vlaanderen	Brugge	1	1	0	59	554	623	0
Brussels	Brussels-North	1	1	1	208	16374	83	7
Brussels	Brussel-South	1	0	1	87	9109	-16	7
Oost-Vlaanderen	Destelbergen	0	1	0	40	2146	320	2
Hainaut	Froyennes	1	1	1	378	492	91	7
Limburg	Genk	1	1	0	63	237	189	0
Oost-Vlaanderen	Gent	1	1	0	129	2120	323	0
Vlaams-Brabant	Grimbergen	1	1	1	163	4256	166	7
West-Vlaanderen	Harelbeke	0	1	0	48	1184	238	0
Limburg	Hasselt	1	0	0	89	104	23	0
Limburg	Houthalen-Centrum	0	1	0	45	96	125	2
Vlaams-Brabant	Leuven	1	1	1	148	4489	312	7
Liège	Liège Oupeye	1	0	0	209	474	-19	0
Liège	Liège Sclessin	1	0	0	334	729	24	0
Luxembourg	Marche-en-Famenne	1	0	0	97	193	-52	0
Hainaut	Marchienne-au-Pont	1	0	0	98	217	-48	0
Antwerpen	Mechelen-Noord	1	0	0	52	134	1	0
West-Vlaanderen	Menen	0	1	0	6	1256	151	0
Hainaut	Montignies-sur-Sambre	1	0	0	149	379	-21	0
Namur	Mornimont	1	1	0	83	134	190	0
Hainaut	Mouscron versant Espierres	1	1	1	101	186	79	7
West-Vlaanderen	Oostende	1	1	0	56	1498	329	0
West-Vlaanderen	Roeselare	0	1	0	5	682	219	0
Liège	Soumagne	1	0	0	101	439	61	0
Brabant Wallon	Vallée du Hain (L'Orchis)	1	0	0	203	241	-40	0
Hainaut	Wasmuel	1	0	0	130	208	48	2

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area during the third wave (i.e. from mid-February 2021 till begin of July).

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/week) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

Figure 3 was developed to offer a dynamic view of the three indicators over time. For further insights on the dynamic of the different indicators, see Section 3.3.

The number of "High circulation" areas is high and stable since four weeks. The rainy event recorded on 01/12 has impacted the 3 indicators to different extends: the "High circulation" areas were underestimated last week, while the fast and increasing trend areas' number are overestimated this week. Overall, a high circulation of the virus is observed but with a decreasing tendency.



Figure 3: The number of areas (among the 41 covered by the wastewater surveillance this week and the 42 normally considered), with positive alerting indicators (latest results on December 08th 2021).

4.4. VARIANTS

Mutations are present in the Omicron variant on the N1 and E genes which could potentially lead to a drop out of those two genes leading to an underestimation of the viral concentrations in wastewaters. However, as the Omicron variant only represented a very small proportion of all sequences reported in Belgium³ and as the correlations between the measured concentrations on the N1, E and N2 genes remain high and significant over the last two weeks, it is unlikely that this variant would cause substantial impact on the wastewater surveillance. A regular investigation of the possible impact of Omicron is done in order to screen the evolution in time.

³ Source : <https://datastudio.google.com/embed/u/0/reporting/c14a5cfc-cab7-4812-848c-0369173148ab/page/urrUC>

5. Appendix – Areas classified by alerting indicator

A 1: Covered areas (26 out of 41 on December 08th 2021) characterized as High circulation sorted in the descending order of importance.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/w) ₃	Incr days ⁴	Date Max cc ⁵
Hainaut	Froyennes	1	1	1	378	492	91	7	08/12/2021
Liège	Liège Sclessin	1	0	0	334	729	24	0	24/11/2021
Brabant Wallon	Basse Wavre (Dyle)	1	0	0	217	299	-61	0	29/11/2021
Liège	Liège Oupeye	1	0	0	209	474	-19	0	15/11/2021
Brussels	Brussels-North	1	1	1	208	16374	83	7	08/12/2021
Brabant Wallon	Vallée du Hain (L'Orchis)	1	0	0	203	241	-40	0	24/11/2021
Liège	Amay	1	1	1	168	246	113	16	06/12/2021
Vlaams-Brabant	Grimbergen	1	1	1	163	4256	166	7	08/12/2021
Hainaut	Montignies-sur-Sambre	1	0	0	149	379	-21	0	15/11/2021
Vlaams-Brabant	Leuven	1	1	1	148	4489	312	7	17/11/2021
Hainaut	Wasmuel	1	0	0	130	208	48	2	15/11/2021
Oost-Vlaanderen	Gent	1	1	0	129	2120	323	0	17/11/2021
Luxembourg	Arlon	1	0	0	121	233	-38	0	22/11/2021
Hainaut	Mouscron versant Espierres	1	1	1	101	186	79	7	06/12/2021
Liège	Soumagne	1	0	0	101	439	61	0	15/11/2021
Hainaut	Marchienne-au-Pont	1	0	0	98	217	-48	0	15/11/2021
Luxembourg	Marche-en-Famenne	1	0	0	97	193	-52	0	29/11/2021
Limburg	Hasselt	1	0	0	89	104	23	0	08/11/2021
Brussels	Brussel-South	1	0	1	87	9109	-16	7	29/11/2021
Namur	Mornimont	1	1	0	83	134	190	0	15/11/2021
Oost-Vlaanderen	Aalst	1	1	0	74	1515	107	0	15/11/2021
Antwerpen	Antwerpen-North	1	0	0	69	217	-46	0	29/11/2021
Limburg	Genk	1	1	0	63	237	189	0	15/11/2021
West-Vlaanderen	Brugge	1	1	0	59	554	623	0	17/11/2021
West-Vlaanderen	Oostende	1	1	0	56	1498	329	0	17/11/2021
Antwerpen	Mechelen-Noord	1	0	0	52	134	1	0	24/11/2021

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area during the third wave (i.e. from mid-February 2021 till begin of July).

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/week) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the measurements. If the date was between 15th February and 15th May 2021, the date is considered to be during the third wave and mentioned as such.

A 2: Covered areas (17 out of 41 on the December 08th 2021) characterized as Fast increase sorted in the descending order of importance.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/w) ₃	Incr days ⁴	Date Max cc ⁵
West-Vlaanderen	Brugge	1	1	0	59	554	623	0	17/11/2021
West-Vlaanderen	Oostende	1	1	0	56	1498	329	0	17/11/2021
Oost-Vlaanderen	Gent	1	1	0	129	2120	323	0	17/11/2021
Oost-Vlaanderen	Destelbergen	0	1	0	40	2146	320	2	3 rd wave
Vlaams-Brabant	Leuven	1	1	1	148	4489	312	7	17/11/2021
West-Vlaanderen	Harelbeke	0	1	0	48	1184	238	0	18/10/2021
West-Vlaanderen	Roeselare	0	1	0	5	682	219	0	3 rd wave
Namur	Mornimont	1	1	0	83	134	190	0	15/11/2021
Limburg	Genk	1	1	0	63	237	189	0	15/11/2021
Vlaams-Brabant	Grimbergen	1	1	1	163	4256	166	7	08/12/2021
West-Vlaanderen	Menen	0	1	0	6	1256	151	0	3 rd wave
Limburg	Houthalen-Centrum	0	1	0	45	96	125	2	08/11/2021
Liège	Amay	1	1	1	168	246	113	16	06/12/2021
Oost-Vlaanderen	Aalst	1	1	0	74	1515	107	0	15/11/2021
Hainaut	Froyennes	1	1	1	378	492	91	7	08/12/2021
Brussels	Brussels-North	1	1	1	208	16374	83	7	08/12/2021
Hainaut	Mouscron versant Espierres	1	1	1	101	186	79	7	06/12/2021

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area during the third wave (i.e. from mid-February 2021 till begin of July).

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/week) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the measurements. If the date was between 15th February and 15th May 2021, the date is considered to be during the third wave and mentioned as such.

A 3: Covered areas (7 out of 41 on the December 08th 2021) characterized as Increasing trend sorted in the descending order of importance.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/w) ₃	Incr days ⁴	Date Max cc ⁵
Liège	Amay	1	1	1	168	246	113	16	06/12/2021
Brussels	Brussels-North	1	1	1	208	16374	83	7	08/12/2021
Brussels	Brussel-South	1	0	1	87	9109	-16	7	29/11/2021
Hainaut	Froyennes	1	1	1	378	492	91	7	08/12/2021
Vlaams-Brabant	Grimbergen	1	1	1	163	4256	166	7	08/12/2021
Vlaams-Brabant	Leuven	1	1	1	148	4489	312	7	17/11/2021
Hainaut	Mouscron versant Espierres	1	1	1	101	186	79	7	06/12/2021

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area during the third wave (i.e. from mid-February 2021 till begin of July).

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/week) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the measurements. If the date was between 15th February and 15th May 2021, the date is considered to be during the third wave and mentioned as such.

A 4: Covered areas (10 out of 41 on the December 08th 2021) in which no alerting indicator are fulfilled.

Province	WWTP	High	Fast	Incr.	Norm. viral cc (%) ¹	Mean viral cc (c./ml) ₂	Norm evol. (%/w) ₃	Incr days ⁴	Date Max cc ⁵
Antwerpen	Aartselaar	0	0	0	36	217	-47	2	29/11/2021
Antwerpen	Antwerpen-South	0	0	0	18	363	16	0	3 rd wave
Vlaams-Brabant	Beersel	0	0	0	20	1274	-4	0	3 rd wave
Oost-Vlaanderen	Dendermonde	0	0	0	6	23	-36	0	22/11/2021
Antwerpen	Deurne	0	0	0	6	327	49	0	3 rd wave
Vlaams-Brabant	Liedekerke	0	0	0	40	1147	7	0	3 rd wave
Namur	Namur-Brumagne	0	0	0	43	122	-54	0	15/11/2021
Oost-Vlaanderen	Sint-Niklaas	0	0	0	38	422	30	0	3 rd wave
Limburg	Tessenderlo	0	0	0	45	177	-21	0	22/11/2021
Antwerpen	Turnhout	0	0	0	41	160	-32	0	24/11/2021

¹ : the viral concentration normalized with the maximum viral concentration measured in the corresponding catchment area during the third wave (i.e. from mid-February 2021 till begin of July).

² : the viral concentration computed on the replicate of the three targeted gene fragments.

³ : the slope (%/week) of the past 7 days moving average of the viral concentration if being above the estimated limit of quantification.

⁴ : the cumulative number of days of increase of the past 14 days moving average of the viral concentration.

⁵ : date at which the measured viral concentration was the highest since the beginning of the measurements. If the date was between 15th February and 15th May 2021, the date is considered to be during the third wave and mentioned as such.